

Topological games

Gary Gruenhage
Auburn University

May 31, 2010

The general setup

I'll mostly talk about games of length ω of the following type:

The general setup

I'll mostly talk about games of length ω of the following type:
There are two players, I and II.

The general setup

I'll mostly talk about games of length ω of the following type:

There are two players, I and II.

In the n^{th} round, I chooses a set I_n (in some restricted collection of sets defined by the “rules” of the game). II responds by choosing a set J_n .

The general setup

I'll mostly talk about games of length ω of the following type:

There are two players, I and II.

In the n^{th} round, I chooses a set I_n (in some restricted collection of sets defined by the "rules" of the game). II responds by choosing a set J_n .

I: I_0 I_1 . . . I_n . . .

II: J_0 J_1 . . . J_n . . .

The general setup

I'll mostly talk about games of length ω of the following type:

There are two players, I and II.

In the n^{th} round, I chooses a set I_n (in some restricted collection of sets defined by the "rules" of the game). II responds by choosing a set J_n .

I: I_0 I_1 . . . I_n . . .

II: J_0 J_1 . . . J_n . . .

There is a round for every $n \in \omega$, and then the game is over.

The general setup

I'll mostly talk about games of length ω of the following type:

There are two players, I and II.

In the n^{th} round, I chooses a set I_n (in some restricted collection of sets defined by the “rules” of the game). II responds by choosing a set J_n .

I: I_0 I_1 . . . I_n . . .

II: J_0 J_1 . . . J_n . . .

There is a round for every $n \in \omega$, and then the game is over.

I *wins* the game if the sequence

$$I_0, J_0, I_1, J_1, \dots$$

of plays of the game satisfies a certain condition (e.g., $\bigcap_{n \in \omega} J_n = \emptyset$); otherwise II wins.

The general setup

I'll mostly talk about games of length ω of the following type:

There are two players, I and II.

In the n^{th} round, I chooses a set I_n (in some restricted collection of sets defined by the "rules" of the game). II responds by choosing a set J_n .

I: I_0 I_1 . . . I_n . . .

II: J_0 J_1 . . . J_n

There is a round for every $n \in \omega$, and then the game is over.

I *wins* the game if the sequence

$$I_0, J_0, I_1, J_1, \dots$$

of plays of the game satisfies a certain condition (e.g., $\bigcap_{n \in \omega} J_n = \emptyset$); otherwise II wins.

In a topological game, the sets I_n and J_n of course are topological objects, e.g., points in a space X , closed subsets of a space, an open cover of a space, etc.

A *strategy for Player I* is a function σ whose domain is the set of finite initial segments $I_0, J_0, \dots, I_n, J_n$ of plays of the game ending with a play by Player II.

A *strategy for Player I* is a function σ whose domain is the set of finite initial segments $I_0, J_0, \dots, I_n, J_n$ of plays of the game ending with a play by Player II.

The strategy σ is a *winning strategy for Player I* if I wins every sequence

$$I_0, J_0, I_1, J_1, \dots$$

of legal plays of the game in which $I_n = \sigma(I_0, J_0, \dots, I_{n-1}, J_{n-1})$ for every $n \in \omega$.

A *strategy for Player I* is a function σ whose domain is the set of finite initial segments $I_0, J_0, \dots, I_n, J_n$ of plays of the game ending with a play by Player II.

The strategy σ is a *winning strategy for Player I* if I wins every sequence

$$I_0, J_0, I_1, J_1, \dots$$

of legal plays of the game in which $I_n = \sigma(I_0, J_0, \dots, I_{n-1}, J_{n-1})$ for every $n \in \omega$.

Wlog, a strategy for Player I may be considered to be a function whose domain is the set of finite sequences J_0, J_1, \dots of plays by Player II, since given a strategy σ as above, and J_0, \dots, J_n , there is a unique way to fill in the plays $I_0 = \sigma(\emptyset)$, $I_1 = \sigma(I_0, J_0)$, etc. of Player I.

A **strategy for Player I** is a function σ whose domain is the set of finite initial segments $I_0, J_0, \dots, I_n, J_n$ of plays of the game ending with a play by Player II.

The strategy σ is a **winning strategy for Player I** if I wins every sequence

$$I_0, J_0, I_1, J_1, \dots$$

of legal plays of the game in which $I_n = \sigma(I_0, J_0, \dots, I_{n-1}, J_{n-1})$ for every $n \in \omega$.

Wlog, a strategy for Player I may be considered to be a function whose domain is the set of finite sequences J_0, J_1, \dots of plays by Player II, since given a strategy σ as above, and J_0, \dots, J_n , there is a unique way to fill in the plays $I_0 = \sigma(\emptyset)$, $I_1 = \sigma(I_0, J_0)$, etc. of Player I.

A **(winning) strategy for Player II** is defined mutatis mutandis.

A **strategy for Player I** is a function σ whose domain is the set of finite initial segments $I_0, J_0, \dots, I_n, J_n$ of plays of the game ending with a play by Player II.

The strategy σ is a **winning strategy for Player I** if I wins every sequence

$$I_0, J_0, I_1, J_1, \dots$$

of legal plays of the game in which $I_n = \sigma(I_0, J_0, \dots, I_{n-1}, J_{n-1})$ for every $n \in \omega$.

Wlog, a strategy for Player I may be considered to be a function whose domain is the set of finite sequences J_0, J_1, \dots of plays by Player II, since given a strategy σ as above, and J_0, \dots, J_n , there is a unique way to fill in the plays $I_0 = \sigma(\emptyset)$, $I_1 = \sigma(I_0, J_0)$, etc. of Player I.

A **(winning) strategy for Player II** is defined mutatis mutandis.

Obviously, I and II cannot both have a winning strategy, and it is possible that neither has. A game is **determined** if one of the players has a winning strategy, otherwise it is **undetermined**.

Banach-Mazur game

Banach-Mazur game $BM(X)$ on X :

Banach-Mazur game

Banach-Mazur game $BM(X)$ on X :

Players E and NE alternately choose nonempty open sets in X :

E: U_0 U_1

...

NE: V_0 V_1

Banach-Mazur game

Banach-Mazur game $BM(X)$ on X :

Players E and NE alternately choose nonempty open sets in X :

E: U_0 U_1

...

NE: V_0 V_1

such that $U_0 \supset V_0 \supset U_1 \supset V_1 \dots$

Banach-Mazur game

Banach-Mazur game $BM(X)$ on X :

Players E and NE alternately choose nonempty open sets in X :

E: U_0 U_1

...

NE: V_0 V_1

such that $U_0 \supset V_0 \supset U_1 \supset V_1 \dots$

E *wins* if $\bigcap_{n \in \omega} U_n = \emptyset$.

Banach-Mazur game

Banach-Mazur game $BM(X)$ on X :

Players E and NE alternately choose nonempty open sets in X :

E: U_0 U_1

...

NE: V_0 V_1

such that $U_0 \supset V_0 \supset U_1 \supset V_1 \dots$

E *wins* if $\bigcap_{n \in \omega} U_n = \emptyset$.

Banach-Mazur game

Banach-Mazur game $BM(X)$ on X :

Players E and NE alternately choose nonempty open sets in X :

E: U_0 U_1

...

NE: V_0 V_1

such that $U_0 \supset V_0 \supset U_1 \supset V_1 \dots$

E *wins* if $\bigcap_{n \in \omega} U_n = \emptyset$.

Theorem (Oxtoby)

A space X is a Baire space iff E has no winning strategy in $BM(X)$.

Banach-Mazur game

Banach-Mazur game $BM(X)$ on X :

Players E and NE alternately choose nonempty open sets in X :

E: U_0 U_1

...

NE: V_0 V_1

such that $U_0 \supset V_0 \supset U_1 \supset V_1 \dots$

E *wins* if $\bigcap_{n \in \omega} U_n = \emptyset$.

Theorem (Oxtoby)

A space X is a Baire space iff E has no winning strategy in $BM(X)$.

If $G(X)$ is a game on X , we'll write " $I \uparrow G(X)$ " to mean "Player I has a winning strategy in $G(X)$ "

Banach-Mazur game

Banach-Mazur game $BM(X)$ on X :

Players E and NE alternately choose nonempty open sets in X :

E: U_0 U_1

...

NE: V_0 V_1

such that $U_0 \supset V_0 \supset U_1 \supset V_1 \dots$

E *wins* if $\bigcap_{n \in \omega} U_n = \emptyset$.

Theorem (Oxtoby)

A space X is a Baire space iff E has no winning strategy in $BM(X)$.

If $G(X)$ is a game on X , we'll write " $I \uparrow G(X)$ " to mean "Player I has a winning strategy in $G(X)$ "

So: X is Baire iff $E \not\uparrow BM(X)$

A space X in which $\text{NE} \uparrow \text{BM}(X)$ is called *weakly α -favorable*

A space X in which $\text{NE} \uparrow \text{BM}(X)$ is called *weakly α -favorable*

X completely metrizable (or Čech complete) $\Rightarrow \text{NE} \uparrow \text{BM}(X) \Rightarrow \text{Baire}$

A space X in which $\text{NE} \uparrow \text{BM}(X)$ is called *weakly α -favorable*

X completely metrizable (or Čech complete) $\Rightarrow \text{NE} \uparrow \text{BM}(X) \Rightarrow \text{Baire}$

There are metric Baire spaces X, Y with $X \times Y$ not Baire

A space X in which $\text{NE} \uparrow \text{BM}(X)$ is called *weakly α -favorable*

X completely metrizable (or Čech complete) $\Rightarrow \text{NE} \uparrow \text{BM}(X) \Rightarrow \text{Baire}$

There are metric Baire spaces X, Y with $X \times Y$ not Baire

$\text{NE} \uparrow \text{BM}(X_\alpha)$ for each $\alpha \Rightarrow \text{NE} \uparrow \text{BM}(\prod_{\alpha \in \kappa} X_\alpha)$

A space X in which $\text{NE} \uparrow \text{BM}(X)$ is called *weakly α -favorable*

X completely metrizable (or Čech complete) $\Rightarrow \text{NE} \uparrow \text{BM}(X) \Rightarrow \text{Baire}$

There are metric Baire spaces X, Y with $X \times Y$ not Baire

$\text{NE} \uparrow \text{BM}(X_\alpha)$ for each $\alpha \Rightarrow \text{NE} \uparrow \text{BM}(\prod_{\alpha \in \kappa} X_\alpha)$

A space X in which $\text{NE} \uparrow \text{BM}(X)$ is called *weakly α -favorable*

X completely metrizable (or Čech complete) $\Rightarrow \text{NE} \uparrow \text{BM}(X) \Rightarrow \text{Baire}$

There are metric Baire spaces X, Y with $X \times Y$ not Baire

$\text{NE} \uparrow \text{BM}(X_\alpha)$ for each $\alpha \Rightarrow \text{NE} \uparrow \text{BM}(\prod_{\alpha \in \kappa} X_\alpha)$

Easy: $\text{NE} \uparrow \text{BM}(X) \Rightarrow X^\kappa$ with **box topology** Baire $\forall \kappa$

A space X in which $NE \uparrow BM(X)$ is called *weakly α -favorable*

X completely metrizable (or Čech complete) $\Rightarrow NE \uparrow BM(X) \Rightarrow$ Baire

There are metric Baire spaces X, Y with $X \times Y$ not Baire

$NE \uparrow BM(X_\alpha)$ for each $\alpha \Rightarrow NE \uparrow BM(\prod_{\alpha \in \kappa} X_\alpha)$

Easy: $NE \uparrow BM(X) \Rightarrow X^\kappa$ with **box topology** Baire $\forall \kappa$

Conjecture (Galvin)

Converse is true: $NE \uparrow BM(X) \iff X^\kappa$ with box topology Baire $\forall \kappa$

A space X in which $NE \uparrow BM(X)$ is called *weakly α -favorable*

X completely metrizable (or Čech complete) $\Rightarrow NE \uparrow BM(X) \Rightarrow$ Baire

There are metric Baire spaces X, Y with $X \times Y$ not Baire

$NE \uparrow BM(X_\alpha)$ for each $\alpha \Rightarrow NE \uparrow BM(\prod_{\alpha \in \kappa} X_\alpha)$

Easy: $NE \uparrow BM(X) \Rightarrow X^\kappa$ with **box topology** Baire $\forall \kappa$

Conjecture (Galvin)

Converse is true: $NE \uparrow BM(X) \iff X^\kappa$ with box topology Baire $\forall \kappa$

Conjecture is consistent if there is a proper class of measurables;

A space X in which $NE \uparrow BM(X)$ is called *weakly α -favorable*

X completely metrizable (or Čech complete) $\Rightarrow NE \uparrow BM(X) \Rightarrow$ Baire

There are metric Baire spaces X, Y with $X \times Y$ not Baire

$NE \uparrow BM(X_\alpha)$ for each $\alpha \Rightarrow NE \uparrow BM(\prod_{\alpha \in \kappa} X_\alpha)$

Easy: $NE \uparrow BM(X) \Rightarrow X^\kappa$ with **box topology** Baire $\forall \kappa$

Conjecture (Galvin)

Converse is true: $NE \uparrow BM(X) \iff X^\kappa$ with box topology Baire $\forall \kappa$

Conjecture is consistent if there is a proper class of measurables;
true in ZFC?

Stationary strategies

A strategy of a player is *stationary* if that player's next move depends only on the previous move of his opponent.

Stationary strategies

A strategy of a player is *stationary* if that player's next move depends only on the previous move of his opponent.

Galvin and Telgarsky obtained a result which says that for games in a certain class, call it *STAT*, if Player I has a winning strategy, then she has a stationary winning strategy.

Stationary strategies

A strategy of a player is *stationary* if that player's next move depends only on the previous move of his opponent.

Galvin and Telgarsky obtained a result which says that for games in a certain class, call it *STAT*, if Player I has a winning strategy, then she has a stationary winning strategy.

$BM(X)$ in this class. If E has a winning strategy in the $BM(X)$, then E has a stationary winning strategy.

Idea of proof:

Idea of proof:

Let σ be a winning strategy for E.

Idea of proof:

Let σ be a winning strategy for E.

Fix a **well-order** \prec of the collection of all **open sets**.

Idea of proof:

Let σ be a winning strategy for E.

Fix a **well-order** \prec of the collection of all **open sets**.

Suppose NE plays $V_n = V$ in round n.

Idea of proof:

Let σ be a winning strategy for E.

Fix a **well-order** \prec of the collection of all **open sets**.

Suppose NE plays $V_n = V$ in round n.

Look at the set $\mathcal{P}(V)$ of all **partial legal plays** $(U_0, V_0, \dots, U_k, V_k)$ of the game with E using σ and **with** $V_k = V$.

Idea of proof:

Let σ be a winning strategy for E.

Fix a **well-order** \prec of the collection of all **open sets**.

Suppose NE plays $V_n = V$ in round n.

Look at the set $\mathcal{P}(V)$ of all **partial legal plays** $(U_0, V_0, \dots, U_k, V_k)$ of the game with E using σ and **with** $V_k = V$.

One may check:

(i) The lexicographic order on $\mathcal{P}(V)$ is a well-order.

Idea of proof:

Let σ be a winning strategy for E.

Fix a **well-order** \prec of the collection of all **open sets**.

Suppose NE plays $V_n = V$ in round n .

Look at the set $\mathcal{P}(V)$ of all **partial legal plays** $(U_0, V_0, \dots, U_k, V_k)$ of the game with E using σ and **with** $V_k = V$.

One may check:

(i) The lexicographic order on $\mathcal{P}(V)$ is a well-order.

(ii) Given $V_n = V$ is NE's move in round n , let

$\tau(V) = \sigma(V_0, V_1, \dots, V_k = V)$, where $(V_0, V_1, \dots, V_k = V)$ is the **least element** of $\mathcal{P}(V)$ ending in V . Then τ defines a stationary winning strategy for E.

Example (Debs)

$\exists X$ with $NE \uparrow BM(X)$ but NE has no stationary winning strategy.

Example (Debs)

$\exists X$ with $NE \uparrow BM(X)$ but NE has no stationary winning strategy.

NE has strategy depending on last two moves of E.

Example (Debs)

$\exists X$ with $NE \uparrow BM(X)$ but NE has no stationary winning strategy.

NE has strategy depending on last two moves of E.

Open: must they all?

Example (Debs)

$\exists X$ with $NE \uparrow BM(X)$ but NE has no stationary winning strategy.

NE has strategy depending on last two moves of E.

Open: must they all?

Question (Telgarsky)

$\exists X$ with NE having winning strategy based on last 3 moves of E but not last 2?

Example (Debs)

$\exists X$ with $NE \uparrow BM(X)$ but NE has no stationary winning strategy.

NE has strategy depending on last two moves of E.

Open: must they all?

Question (Telgarsky)

$\exists X$ with NE having winning strategy based on last 3 moves of E but not last 2?

Remark: Galvin and Telgarsky, Debs: $NE \uparrow BM(X) \Rightarrow$ NE has winning strategy based on last move of opponent and his own last move.

Telgársky's game

Let X be a space, and let \mathbb{K} be a **closed hereditary** class of spaces.

Telgársky's game

Let X be a space, and let \mathbb{K} be a **closed hereditary** class of spaces. We define the game $G(\mathbb{K}, X)$. There are two players, I and II.

Telgársky's game

Let X be a space, and let \mathbb{K} be a **closed hereditary** class of spaces.

We define the game $G(\mathbb{K}, X)$. There are two players, I and II.

Player I begins by choosing a nonempty closed subset A_0 of X such that $A_0 \in \mathbb{K}$.

Telgársky's game

Let X be a space, and let \mathbb{K} be a **closed hereditary** class of spaces.

We define the game $G(\mathbb{K}, X)$. There are two players, I and II.

Player I begins by choosing a nonempty closed subset A_0 of X such that $A_0 \in \mathbb{K}$.

II responds by choosing a **closed** set $B_0 \subset X \setminus A_0$.

Telgársky's game

Let X be a space, and let \mathbb{K} be a **closed hereditary** class of spaces.

We define the game $G(\mathbb{K}, X)$. There are two players, I and II.

Player I begins by choosing a nonempty closed subset A_0 of X such that $A_0 \in \mathbb{K}$.

II responds by choosing a **closed** set $B_0 \subset X \setminus A_0$.

Player I then chooses a nonempty closed $A_1 \subset B_0$ with $A_1 \in \mathbb{K}$.

Telgársky's game

Let X be a space, and let \mathbb{K} be a **closed hereditary** class of spaces.

We define the game $G(\mathbb{K}, X)$. There are two players, I and II.

Player I begins by choosing a nonempty closed subset A_0 of X such that $A_0 \in \mathbb{K}$.

II responds by choosing a **closed** set $B_0 \subset X \setminus A_0$.

Player I then chooses a nonempty closed $A_1 \subset B_0$ with $A_1 \in \mathbb{K}$.

II chooses a closed $B_1 \subset B_0 \setminus A_1$

etc.

I: $A_0 \in \mathbb{K}$ $A_1 \subset B_0, A_1 \in \mathbb{K}$. . . $A_n \subset B_{n-1}, A_n \in \mathbb{K}$

II: $B_0 \subset X \setminus A_0$ $B_1 \subset B_0 \setminus A_1$. . . $B_n \subset B_{n-1} \setminus A_n$

I: $A_0 \in \mathbb{K}$ $A_1 \subset B_0, A_1 \in \mathbb{K}$. . . $A_n \subset B_{n-1}, A_n \in \mathbb{K}$

II: $B_0 \subset X \setminus A_0$ $B_1 \subset B_0 \setminus A_1$. . . $B_n \subset B_{n-1} \setminus A_n$

We say **I wins** the game if $\bigcap_{n \in \omega} B_n = \emptyset$; otherwise II wins.

I: $A_0 \in \mathbb{K}$ $A_1 \subset B_0, A_1 \in \mathbb{K}$. . . $A_n \subset B_{n-1}, A_n \in \mathbb{K}$

II: $B_0 \subset X \setminus A_0$ $B_1 \subset B_0 \setminus A_1$. . . $B_n \subset B_{n-1} \setminus A_n$

We say **I wins** the game if $\bigcap_{n \in \omega} B_n = \emptyset$; otherwise II wins.

The space X is said to be **\mathbb{K} -like** if Player I has a winning strategy in $G(\mathbb{K}, X)$ (i.e., if $I \uparrow G(\mathbb{K}, X)$).

I: $A_0 \in \mathbb{K}$ $A_1 \subset B_0, A_1 \in \mathbb{K}$. . . $A_n \subset B_{n-1}, A_n \in \mathbb{K}$

II: $B_0 \subset X \setminus A_0$ $B_1 \subset B_0 \setminus A_1$. . . $B_n \subset B_{n-1} \setminus A_n$

We say **I wins** the game if $\bigcap_{n \in \omega} B_n = \emptyset$; otherwise II wins.

The space X is said to be **\mathbb{K} -like** if Player I has a winning strategy in $G(\mathbb{K}, X)$ (i.e., if $I \uparrow G(\mathbb{K}, X)$).

Trivially, \mathbb{K} is contained in the class of \mathbb{K} -like spaces.

Let \mathbb{DC} be the class of all topological sums (discrete unions) of compact sets.

Let \mathbb{DC} be the class of all topological sums (discrete unions) of compact sets.

If $X \in \mathbb{DC}$, then $X \times Y$ is paracompact \forall paracompact Y

Let \mathbb{DC} be the class of all topological sums (discrete unions) of compact sets.

If $X \in \mathbb{DC}$, then $X \times Y$ is paracompact \forall paracompact Y
(because compact \times paracompact is paracompact)

Let \mathbb{DC} be the class of all topological sums (discrete unions) of compact sets.

If $X \in \mathbb{DC}$, then $X \times Y$ is paracompact \forall paracompact Y
(because compact \times paracompact is paracompact)

Theorem (Telgarsky)

If X is paracompact and \mathbb{DC} -like, then $X \times Y$ is paracompact for all paracompact spaces Y .

Let \mathbb{DC} be the class of all topological sums (discrete unions) of compact sets.

If $X \in \mathbb{DC}$, then $X \times Y$ is paracompact \forall paracompact Y
(because compact \times paracompact is paracompact)

Theorem (Telgarsky)

If X is paracompact and \mathbb{DC} -like, then $X \times Y$ is paracompact for all paracompact spaces Y .

(Sub)paracompact scattered spaces, more generally \mathbb{C} -scattered (every closed subspace has a point of local compactness), and spaces with a σ -closure-preserving cover by compact sets, are \mathbb{DC} -like.

Telgarsky's Conjecture

$X \times Y$ is paracompact \forall paracompact Y iff I has a winning strategy in $G(\mathbb{DC}, X)$ (i.e., X is \mathbb{DC} -like)

Telgarsky's Conjecture

$X \times Y$ is paracompact \forall paracompact Y iff I has a winning strategy in $G(\mathbb{DC}, X)$ (i.e., X is \mathbb{DC} -like)

Theorem(Alster, 2006)

Telgarsky's Conjecture holds if X has a base of cardinality $\leq \aleph_1$

Telgärsky's game is in STAT. Player I has a winning strategy iff she has a stationary winning strategy.

Telgärsky's game is in STAT. Player I has a winning strategy iff she has a stationary winning strategy.

Thus, X is \mathbb{K} -like iff there is a function $\sigma : \mathcal{C}(X) \rightarrow \mathbb{K}$, where $\mathcal{C}(X)$ is the collection of nonempty closed subsets of X , such that

Telgärsky's game is in STAT. Player I has a winning strategy iff she has a stationary winning strategy.

Thus, X is \mathbb{K} -like iff there is a function $\sigma : \mathcal{C}(X) \rightarrow \mathbb{K}$, where $\mathcal{C}(X)$ is the collection of nonempty closed subsets of X , such that

① $\sigma(C) \subset C$;

Telgärsky's game is in STAT. Player I has a winning strategy iff she has a stationary winning strategy.

Thus, X is \mathbb{K} -like iff there is a function $\sigma : \mathcal{C}(X) \rightarrow \mathbb{K}$, where $\mathcal{C}(X)$ is the collection of nonempty closed subsets of X , such that

- 1 $\sigma(C) \subset C$;
- 2 $\sigma(C) \in \mathbb{K}$;

Telgärsky's game is in STAT. Player I has a winning strategy iff she has a stationary winning strategy.

Thus, X is \mathbb{K} -like iff there is a function $\sigma : \mathcal{C}(X) \rightarrow \mathbb{K}$, where $\mathcal{C}(X)$ is the collection of nonempty closed subsets of X , such that

- 1 $\sigma(C) \subset C$;
- 2 $\sigma(C) \in \mathbb{K}$;
- 3 If

$$X = B_{-1} \supset B_0 \supset B_1 \cdots \supset B_n \supset \dots$$

is a decreasing sequence of closed sets such that for each $n \in \omega$, $B_n \cap \sigma(B_{n-1}) = \emptyset$, then $\bigcap_{n \in \omega} B_n = \emptyset$.

A space X is a *D-space* if, given an open nbhd $N(x)$ for each $x \in X$, there is a closed discrete $D \subset X$ such that $N(D) = \{N(x) : x \in D\}$ covers X .

A space X is a *D-space* if, given an open nbhd $N(x)$ for each $x \in X$, there is a closed discrete $D \subset X$ such that $N(D) = \{N(x) : x \in D\}$ covers X .

Compact or σ -compact implies D .

A space X is a *D-space* if, given an open nbhd $N(x)$ for each $x \in X$, there is a closed discrete $D \subset X$ such that $N(D) = \{N(x) : x \in D\}$ covers X .

Compact or σ -compact implies D .

Open question:

Do any of the other standard covering properties (e.g., (Lindelöf, paracompact, metacompact, submetacompact,...) imply D ?

X is said to be *Menger* if, given open covers $\mathcal{U}_0, \mathcal{U}_1, \dots$, there are finite $\mathcal{F}_n \subset \mathcal{U}_n$ such that $\bigcup_{n \in \omega} \mathcal{F}_n$ covers X .

X is said to be *Menger* if, given open covers $\mathcal{U}_0, \mathcal{U}_1, \dots$, there are finite $\mathcal{F}_n \subset \mathcal{U}_n$ such that $\bigcup_{n \in \omega} \mathcal{F}_n$ covers X .

σ -compact \Rightarrow Menger \Rightarrow Lindelöf

X is said to be *Menger* if, given open covers $\mathcal{U}_0, \mathcal{U}_1, \dots$, there are finite $\mathcal{F}_n \subset \mathcal{U}_n$ such that $\bigcup_{n \in \omega} \mathcal{F}_n$ covers X .

σ -compact \Rightarrow Menger \Rightarrow Lindelöf

Irrationals are not Menger.

X is said to be *Menger* if, given open covers $\mathcal{U}_0, \mathcal{U}_1, \dots$, there are finite $\mathcal{F}_n \subset \mathcal{U}_n$ such that $\bigcup_{n \in \omega} \mathcal{F}_n$ covers X .

σ -compact \Rightarrow Menger \Rightarrow Lindelöf

Irrationals are not Menger.

$MA(\kappa) \Rightarrow$ any $X \subset \mathbb{R}$ with $|X| \leq \kappa$ is Menger.

X is said to be *Menger* if, given open covers $\mathcal{U}_0, \mathcal{U}_1, \dots$, there are finite $\mathcal{F}_n \subset \mathcal{U}_n$ such that $\bigcup_{n \in \omega} \mathcal{F}_n$ covers X .

σ -compact \Rightarrow Menger \Rightarrow Lindelöf

Irrationals are not Menger.

$MA(\kappa) \Rightarrow$ any $X \subset \mathbb{R}$ with $|X| \leq \kappa$ is Menger.

Hurewicz: analytic + Menger \Rightarrow σ -compact

X is said to be *Menger* if, given open covers $\mathcal{U}_0, \mathcal{U}_1, \dots$, there are finite $\mathcal{F}_n \subset \mathcal{U}_n$ such that $\bigcup_{n \in \omega} \mathcal{F}_n$ covers X .

σ -compact \Rightarrow Menger \Rightarrow Lindelöf

Irrationals are not Menger.

$MA(\kappa) \Rightarrow$ any $X \subset \mathbb{R}$ with $|X| \leq \kappa$ is Menger.

Hurewicz: analytic + Menger \Rightarrow σ -compact

(Fremlin-Miller)ZFC $\Rightarrow \exists$ non- σ -compact Menger $X \subset \mathbb{R}$

X is said to be *Menger* if, given open covers $\mathcal{U}_0, \mathcal{U}_1, \dots$, there are finite $\mathcal{F}_n \subset \mathcal{U}_n$ such that $\bigcup_{n \in \omega} \mathcal{F}_n$ covers X .

σ -compact \Rightarrow Menger \Rightarrow Lindelöf

Irrationals are not Menger.

$MA(\kappa) \Rightarrow$ any $X \subset \mathbb{R}$ with $|X| \leq \kappa$ is Menger.

Hurewicz: analytic + Menger \Rightarrow σ -compact

(Fremlin-Miller)ZFC $\Rightarrow \exists$ non- σ -compact Menger $X \subset \mathbb{R}$

Theorem(Aurichi)

Menger spaces are D -spaces.

X is said to be *Menger* if, given open covers $\mathcal{U}_0, \mathcal{U}_1, \dots$, there are finite $\mathcal{F}_n \subset \mathcal{U}_n$ such that $\bigcup_{n \in \omega} \mathcal{F}_n$ covers X .

σ -compact \Rightarrow Menger \Rightarrow Lindelöf

Irrationals are not Menger.

$MA(\kappa) \Rightarrow$ any $X \subset \mathbb{R}$ with $|X| \leq \kappa$ is Menger.

Hurewicz: analytic + Menger \Rightarrow σ -compact

(Fremlin-Miller)ZFC $\Rightarrow \exists$ non- σ -compact Menger $X \subset \mathbb{R}$

Theorem(Aurichi)

Menger spaces are D -spaces.

Not clear how to do a direct proof. A game characterization of Menger, due to Hurewicz, provides an easy proof.

Menger game $M(X)$:

Menger game $M(X)$:

In round n , Player I chooses open cover \mathcal{U}_n of X .

Menger game $M(X)$:

In round n , Player I chooses open cover \mathcal{U}_n of X .

II responds with finite $\mathcal{F}_n \subset \mathcal{U}_n$.

Menger game $M(X)$:

In round n , Player I chooses open cover \mathcal{U}_n of X .

II responds with finite $\mathcal{F}_n \subset \mathcal{U}_n$.

II *wins* if $\bigcup_{n \in \omega} \mathcal{F}_n$ covers X .

Menger game $M(X)$:

In round n , Player I chooses open cover \mathcal{U}_n of X .

II responds with finite $\mathcal{F}_n \subset \mathcal{U}_n$.

II *wins* if $\bigcup_{n \in \omega} \mathcal{F}_n$ covers X .

Equivalently, I's cover is closed under finite unions, and II chooses $U_n \in \mathcal{U}_n$. II wins if $\bigcup_{n \in \omega} U_n = X$.

Menger game $M(X)$:

In round n , Player I chooses open cover \mathcal{U}_n of X .

II responds with finite $\mathcal{F}_n \subset \mathcal{U}_n$.

II *wins* if $\bigcup_{n \in \omega} \mathcal{F}_n$ covers X .

Equivalently, I's cover is closed under finite unions, and II chooses $U_n \in \mathcal{U}_n$. II wins if $\bigcup_{n \in \omega} U_n = X$.

Easy: $I \nVdash M(X) \Rightarrow X$ Menger

Menger game $M(X)$:

In round n , Player I chooses open cover \mathcal{U}_n of X .

II responds with finite $\mathcal{F}_n \subset \mathcal{U}_n$.

II *wins* if $\bigcup_{n \in \omega} \mathcal{F}_n$ covers X .

Equivalently, I's cover is closed under finite unions, and II chooses $U_n \in \mathcal{U}_n$. II wins if $\bigcup_{n \in \omega} U_n = X$.

Easy: $I \nexists M(X) \Rightarrow X$ Menger

Theorem (Hurewicz)

X is Menger iff $I \nexists M(X)$

Proof of Aurichi's theorem

Proof of Aurichi's theorem

Assume X Menger.

Proof of Aurichi's theorem

Assume X Menger.

Let N be a neighborhood assignment on X .

Proof of Aurichi's theorem

Assume X Menger.

Let N be a neighborhood assignment on X .

Let Player I's first play be $\{N(x) : x \in X\}$.

Proof of Aurichi's theorem

Assume X Menger.

Let N be a neighborhood assignment on X .

Let Player I's first play be $\{N(x) : x \in X\}$.

Player II responds with $\{N(x) : x \in F_0\}$, $F_0 \in [X]^{<\omega}$.

Proof of Aurichi's theorem

Assume X Menger.

Let N be a neighborhood assignment on X .

Let Player I's first play be $\{N(x) : x \in X\}$.

Player II responds with $\{N(x) : x \in F_0\}$, $F_0 \in [X]^{<\omega}$.

Let $V_0 = \cup\{N(x) : x \in F_0\}$, and let I then play

$$\{V_0 \cup N(x) : x \in X \setminus V_0\}.$$

Proof of Aurichi's theorem

Assume X Menger.

Let N be a neighborhood assignment on X .

Let Player I's first play be $\{N(x) : x \in X\}$.

Player II responds with $\{N(x) : x \in F_0\}$, $F_0 \in [X]^{<\omega}$.

Let $V_0 = \cup\{N(x) : x \in F_0\}$, and let I then play

$$\{V_0 \cup N(x) : x \in X \setminus V_0\}.$$

Then similarly, if II's reply is $\{V_0 \cup N(x) : x \in F_1\}$, where $F_1 \in [X \setminus V_0]^{<\omega}$,

Proof of Aurichi's theorem

Assume X Menger.

Let N be a neighborhood assignment on X .

Let Player I's first play be $\{N(x) : x \in X\}$.

Player II responds with $\{N(x) : x \in F_0\}$, $F_0 \in [X]^{<\omega}$.

Let $V_0 = \cup\{N(x) : x \in F_0\}$, and let I then play

$$\{V_0 \cup N(x) : x \in X \setminus V_0\}.$$

Then similarly, if II's reply is $\{V_0 \cup N(x) : x \in F_1\}$, where $F_1 \in [X \setminus V_0]^{<\omega}$,

let $V_1 = V_0 \cup \cup\{N(x) : x \in F_1\}$ and let I play

$$\{V_1 \cup N(x) : x \in X \setminus V_1\},$$

Proof of Aurichi's theorem

Assume X Menger.

Let N be a neighborhood assignment on X .

Let Player I's first play be $\{N(x) : x \in X\}$.

Player II responds with $\{N(x) : x \in F_0\}$, $F_0 \in [X]^{<\omega}$.

Let $V_0 = \cup\{N(x) : x \in F_0\}$, and let I then play

$$\{V_0 \cup N(x) : x \in X \setminus V_0\}.$$

Then similarly, if II's reply is $\{V_0 \cup N(x) : x \in F_1\}$, where $F_1 \in [X \setminus V_0]^{<\omega}$,

let $V_1 = V_0 \cup \cup\{N(x) : x \in F_1\}$ and let I play

$$\{V_1 \cup N(x) : x \in X \setminus V_1\},$$

and so on.

Proof of Aurichi's theorem

Assume X Menger.

Let N be a neighborhood assignment on X .

Let Player I's first play be $\{N(x) : x \in X\}$.

Player II responds with $\{N(x) : x \in F_0\}$, $F_0 \in [X]^{<\omega}$.

Let $V_0 = \cup\{N(x) : x \in F_0\}$, and let I then play

$$\{V_0 \cup N(x) : x \in X \setminus V_0\}.$$

Then similarly, if II's reply is $\{V_0 \cup N(x) : x \in F_1\}$, where $F_1 \in [X \setminus V_0]^{<\omega}$,

let $V_1 = V_0 \cup \cup\{N(x) : x \in F_1\}$ and let I play

$$\{V_1 \cup N(x) : x \in X \setminus V_1\},$$

and so on.

This defines a strategy for Player I.

X is Menger, so this can't be a winning strategy.

X is Menger, so this can't be a winning strategy.

Therefore there is some play of the game with I using this strategy such that, if F_0, F_1, \dots code the plays of II, then

$$X = \bigcup_{n \in \omega} V_n = \bigcup \{N(x) : x \in \bigcup_{n \in \omega} F_n\}.$$

X is Menger, so this can't be a winning strategy.

Therefore there is some play of the game with I using this strategy such that, if F_0, F_1, \dots code the plays of II, then

$$X = \bigcup_{n \in \omega} V_n = \bigcup \{N(x) : x \in \bigcup_{n \in \omega} F_n\}.$$

Let $D = \bigcup_{n \in \omega} F_n$. Then $N(D)$ covers X .

X is Menger, so this can't be a winning strategy.

Therefore there is some play of the game with I using this strategy such that, if F_0, F_1, \dots code the plays of II, then

$$X = \bigcup_{n \in \omega} V_n = \bigcup \{N(x) : x \in \bigcup_{n \in \omega} F_n\}.$$

Let $D = \bigcup_{n \in \omega} F_n$. Then $N(D)$ covers X .

Since for each n , we have $F_n \subset V_n$ and $F_{n+1} \cap V_n = \emptyset$, it is easy to check that D is a closed discrete subset of X . Hence X is a D -space. \square

Proof of Hurewicz's theorem

To show: Menger \Rightarrow $I \not\uparrow M(X)$

Proof of Hurewicz's theorem

To show: Menger \Rightarrow I \nexists $M(X)$

Suppose X Menger, and consider any fixed strategy by I. We show it can be defeated.

Proof of Hurewicz's theorem

To show: Menger \Rightarrow I \nexists $M(X)$

Suppose X Menger, and consider any fixed strategy by I. We show it can be defeated.

X is Lindelöf, so wlog, I always chooses a countable open cover.

Proof of Hurewicz's theorem

To show: Menger \Rightarrow I \nexists $M(X)$

Suppose X Menger, and consider any fixed strategy by I. We show it can be defeated.

X is Lindelöf, so wlog, I always chooses a countable open cover.

Since II is choosing a finite subcollection, it doesn't hurt I to be restricted to increasing open covers.

Proof of Hurewicz's theorem

To show: Menger \Rightarrow I \nexists $M(X)$

Suppose X Menger, and consider any fixed strategy by I. We show it can be defeated.

X is Lindelöf, so wlog, I always chooses a countable open cover.

Since II is choosing a finite subcollection, it doesn't hurt I to be restricted to increasing open covers.

In this case, it doesn't harm II to be restricted to choosing a single element of the cover.

Proof of Hurewicz's theorem

To show: Menger \Rightarrow I \nexists $M(X)$

Suppose X Menger, and consider any fixed strategy by I. We show it can be defeated.

X is Lindelöf, so wlog, I always chooses a countable open cover.

Since II is choosing a finite subcollection, it doesn't hurt I to be restricted to increasing open covers.

In this case, it doesn't harm II to be restricted to choosing a single element of the cover.

Finally, if II chooses U , I may as well make the first member of the increasing open cover that is his response contain U .

Proof of Hurewicz's theorem

To show: Menger \Rightarrow I \nexists $M(X)$

Suppose X Menger, and consider any fixed strategy by I. We show it can be defeated.

X is Lindelöf, so wlog, I always chooses a countable open cover.

Since II is choosing a finite subcollection, it doesn't hurt I to be restricted to increasing open covers.

In this case, it doesn't harm II to be restricted to choosing a single element of the cover.

Finally, if II chooses U , I may as well make the first member of the increasing open cover that is his response contain U .

To summarize: I chooses countable increasing open cover, each member of which contains II's previous move. II chooses a member of I's cover.

Proof of Hurewicz's theorem

To show: Menger \Rightarrow I \nVdash M(X)

Suppose X Menger, and consider any fixed strategy by I. We show it can be defeated.

X is Lindelöf, so wlog, I always chooses a countable open cover.

Since II is choosing a finite subcollection, it doesn't hurt I to be restricted to increasing open covers.

In this case, it doesn't harm II to be restricted to choosing a single element of the cover.

Finally, if II chooses U , I may as well make the first member of the increasing open cover that is his response contain U .

To summarize: I chooses countable increasing open cover, each member of which contains II's previous move. II chooses a member of I's cover.

Want to show that II can defeat I's strategy.

$$\begin{array}{ccccccc}
 & & & & \{U_n\}_n & & \\
 & & & & & & \\
 \{U_{0m}\}_m & & \{U_{1m}\}_m & & \dots & & \{U_{nm}\}_m \quad \dots \\
 & & & & & & \\
 & & & & \{U_{n0k}\}_k & \{U_{n1k}\}_k & \dots \{U_{nmk}\}_k
 \end{array}$$

$$\begin{array}{ccccccc}
 & & & & \{U_n\}_n & & \\
 & & & & & & \\
 \{U_{0m}\}_m & & \{U_{1m}\}_m & & \dots & & \{U_{nm}\}_m \quad \dots \\
 & & & & & & \\
 & & & & \{U_{n0k}\}_k & \{U_{n1k}\}_k & \dots \{U_{nmk}\}_k
 \end{array}$$

Let $\{U_n\}_n$ be I's first move using the strategy.

$$\begin{array}{ccccccc}
 & & & & & & \{U_n\}_n \\
 & & & & & & \\
 \{U_{0m}\}_m & & \{U_{1m}\}_m & & \dots & & \{U_{nm}\}_m & \dots \\
 & & & & & & & \\
 & & & & & & \{U_{n0k}\}_k & \{U_{n1k}\}_k & \dots & \{U_{nmk}\}_k
 \end{array}$$

Let $\{U_n\}_n$ be I's first move using the strategy.
 If II responds with U_n , let $\{U_{nm}\}_m$ be I's next move.
 Then if II plays U_{nm} , let $\{U_{nmk}\}_k$ be I's reply.

$$\{U_n\}_n$$

$$\{U_{0m}\}_m \quad \{U_{1m}\}_m \quad \dots \quad \{U_{nm}\}_m \quad \dots$$

$$\{U_{n0k}\}_k \quad \{U_{n1k}\}_k \quad \dots \quad \{U_{nmk}\}_k$$

Let $\{U_n\}_n$ be I's first move using the strategy.

If II responds with U_n , let $\{U_{nm}\}_m$ be I's next move.

Then if II plays U_{nm} , let $\{U_{nmk}\}_k$ be I's reply.

In this way we define a "game tree" $\{U_\sigma\}_{\sigma \in \omega^{<\omega}}$. (Let $U_\emptyset = \emptyset$.)

We need to show that there is a play of the game, i.e., a branch of the game tree, for which the corresponding open sets cover.

$$\{U_n\}_n$$

$$\{U_{0m}\}_m \quad \{U_{1m}\}_m \quad \dots \quad \{U_{nm}\}_m \quad \dots$$

$$\{U_{n0k}\}_k \quad \{U_{n1k}\}_k \quad \dots \quad \{U_{nmk}\}_k$$

Let $\{U_n\}_n$ be I's first move using the strategy.

If II responds with U_n , let $\{U_{nm}\}_m$ be I's next move.

Then if II plays U_{nm} , let $\{U_{nmk}\}_k$ be I's reply.

In this way we define a "game tree" $\{U_\sigma\}_{\sigma \in \omega^{<\omega}}$. (Let $U_\emptyset = \emptyset$.)

We need to show that there is a play of the game, i.e., a branch of the game tree, for which the corresponding open sets cover.

That is, we want $f : \omega \rightarrow \omega$ such that $X = \bigcup_{n \in \omega} U_{f \upharpoonright n}$.

$$\{U_n\}_n$$

$$\{U_{0m}\}_m \quad \{U_{1m}\}_m \quad \dots \quad \{U_{nm}\}_m \quad \dots$$

$$\{U_{n0k}\}_k \quad \{U_{n1k}\}_k \quad \dots \quad \{U_{nmk}\}_k$$

Let $\{U_n\}_n$ be I's first move using the strategy.

If II responds with U_n , let $\{U_{nm}\}_m$ be I's next move.

Then if II plays U_{nm} , let $\{U_{nmk}\}_k$ be I's reply.

In this way we define a "game tree" $\{U_\sigma\}_{\sigma \in \omega^{<\omega}}$. (Let $U_\emptyset = \emptyset$.)

We need to show that there is a play of the game, i.e., a branch of the game tree, for which the corresponding open sets cover.

That is, we want $f : \omega \rightarrow \omega$ such that $X = \bigcup_{n \in \omega} U_{f \upharpoonright n}$.

A naive idea is to apply the Menger property to the countably many covers $\{U_{\sigma \frown n}\}_n$. There is a choice of one member of each that covers. But this doesn't get you a branch that covers!

Instead, we use the game tree to define covers $\{V_k^n\}_k$ as follows.

Instead, we use the game tree to define covers $\{V_k^n\}_k$ as follows.

Let $V_k^0 = U_k$; then $\{V_k^0\}_k$ is increasing open cover

Instead, we use the game tree to define covers $\{V_k^n\}_k$ as follows.

Let $V_k^0 = U_k$; then $\{V_k^0\}_k$ is increasing open cover

Next let

$$V_k^1 = U_k \cap U_{0k} \cap U_{1k} \cap \dots \cap U_{k-1,k}.$$

Instead, we use the game tree to define covers $\{V_k^n\}_k$ as follows.

Let $V_k^0 = U_k$; then $\{V_k^0\}_k$ is increasing open cover

Next let

$$V_k^1 = U_k \cap U_{0k} \cap U_{1k} \cap \dots \cap U_{k-1,k}.$$

Since we have assumed $U_{ik} \supset U_i$, note that $V_k^1 = \bigcap_{\sigma \in \omega \leq 1} U_{\sigma k}$, i.e.,

$V_k^1 =$ intersection of all k^{th} terms of all of I's plays from rounds 0 and 1

Instead, we use the game tree to define covers $\{V_k^n\}_k$ as follows.

Let $V_k^0 = U_k$; then $\{V_k^0\}_k$ is increasing open cover

Next let

$$V_k^1 = U_k \cap U_{0k} \cap U_{1k} \cap \dots \cap U_{k-1,k}.$$

Since we have assumed $U_{ik} \supset U_i$, note that $V_k^1 = \bigcap_{\sigma \in \omega \leq 1} U_{\sigma k}$, i.e.,

V_k^1 = intersection of all k^{th} terms of all of I's plays from rounds 0 and 1

Claim. $\{V_k^1\}_k$ is an increasing open cover.

Instead, we use the game tree to define covers $\{V_k^n\}_k$ as follows.

Let $V_k^0 = U_k$; then $\{V_k^0\}_k$ is increasing open cover

Next let

$$V_k^1 = U_k \cap U_{0k} \cap U_{1k} \cap \dots \cap U_{k-1,k}.$$

Since we have assumed $U_{ik} \supset U_i$, note that $V_k^1 = \bigcap_{\sigma \in \omega \leq 1} U_{\sigma k}$, i.e., V_k^1 = intersection of all k^{th} terms of all of I's plays from rounds 0 and 1

Claim. $\{V_k^1\}_k$ is an increasing open cover.

Increasing: $V_{k+1}^1 = U_{k+1} \cap \bigcap_{i \leq k} U_{i,k+1}$

Instead, we use the game tree to define covers $\{V_k^n\}_k$ as follows.

Let $V_k^0 = U_k$; then $\{V_k^0\}_k$ is increasing open cover

Next let

$$V_k^1 = U_k \cap U_{0k} \cap U_{1k} \cap \dots \cap U_{k-1,k}.$$

Since we have assumed $U_{ik} \supset U_i$, note that $V_k^1 = \bigcap_{\sigma \in \omega \leq 1} U_{\sigma k}$, i.e., $V_k^1 =$ intersection of all k^{th} terms of all of I's plays from rounds 0 and 1

Claim. $\{V_k^1\}_k$ is an increasing open cover.

Increasing: $V_{k+1}^1 = U_{k+1} \cap \bigcap_{i \leq k} U_{i,k+1} \supseteq U_k \cap (\bigcap_{i < k} U_{ik}) \cap U_{k,k+1} = U_k \cap (\bigcap_{i < k} U_{ik}) = V_k^1.$

Instead, we use the game tree to define covers $\{V_k^n\}_k$ as follows.

Let $V_k^0 = U_k$; then $\{V_k^0\}_k$ is increasing open cover

Next let

$$V_k^1 = U_k \cap U_{0k} \cap U_{1k} \cap \dots \cap U_{k-1,k}.$$

Since we have assumed $U_{ik} \supset U_i$, note that $V_k^1 = \bigcap_{\sigma \in \omega \leq 1} U_{\sigma k}$, i.e., V_k^1 = intersection of all k^{th} terms of all of I's plays from rounds 0 and 1

Claim. $\{V_k^1\}_k$ is an increasing open cover.

Increasing: $V_{k+1}^1 = U_{k+1} \cap \bigcap_{i \leq k} U_{i,k+1} \supseteq U_k \cap (\bigcap_{i < k} U_{ik}) \cap U_{k,k+1} = U_k \cap (\bigcap_{i < k} U_{ik}) = V_k^1.$

Cover: Let $x \in X$. There is k_0 with $x \in U_{k_0}$.

Instead, we use the game tree to define covers $\{V_k^n\}_k$ as follows.

Let $V_k^0 = U_k$; then $\{V_k^0\}_k$ is increasing open cover

Next let

$$V_k^1 = U_k \cap U_{0k} \cap U_{1k} \cap \dots \cap U_{k-1,k}.$$

Since we have assumed $U_{ik} \supset U_i$, note that $V_k^1 = \bigcap_{\sigma \in \omega \leq 1} U_{\sigma k}$, i.e., $V_k^1 =$ intersection of all k^{th} terms of all of I's plays from rounds 0 and 1

Claim. $\{V_k^1\}_k$ is an increasing open cover.

Increasing: $V_{k+1}^1 = U_{k+1} \cap \bigcap_{i \leq k} U_{i,k+1} \supseteq U_k \cap (\bigcap_{i < k} U_{ik}) \cap U_{k,k+1} = U_k \cap (\bigcap_{i < k} U_{ik}) = V_k^1.$

Cover: Let $x \in X$. There is k_0 with $x \in U_{k_0}$.

For each $i < k_0$, there is l_i such that $x \in U_{il_i}$.

Instead, we use the game tree to define covers $\{V_k^n\}_k$ as follows.

Let $V_k^0 = U_k$; then $\{V_k^0\}_k$ is increasing open cover

Next let

$$V_k^1 = U_k \cap U_{0k} \cap U_{1k} \cap \dots \cap U_{k-1,k}.$$

Since we have assumed $U_{ik} \supset U_i$, note that $V_k^1 = \bigcap_{\sigma \in \omega \leq 1} U_{\sigma k}$, i.e., $V_k^1 =$ intersection of all k^{th} terms of all of I's plays from rounds 0 and 1

Claim. $\{V_k^1\}_k$ is an increasing open cover.

Increasing: $V_{k+1}^1 = U_{k+1} \cap \bigcap_{i \leq k} U_{i,k+1} \supseteq U_k \cap (\bigcap_{i < k} U_{ik}) \cap U_{k,k+1} = U_k \cap (\bigcap_{i < k} U_{ik}) = V_k^1.$

Cover: Let $x \in X$. There is k_0 with $x \in U_{k_0}$.

For each $i < k_0$, there is l_i such that $x \in U_{il_i}$.

Let k be greater than k_0 and the l_i 's.

Instead, we use the game tree to define covers $\{V_k^n\}_k$ as follows.

Let $V_k^0 = U_k$; then $\{V_k^0\}_k$ is increasing open cover

Next let

$$V_k^1 = U_k \cap U_{0k} \cap U_{1k} \cap \dots \cap U_{k-1,k}.$$

Since we have assumed $U_{ik} \supset U_i$, note that $V_k^1 = \bigcap_{\sigma \in \omega \leq 1} U_{\sigma k}$, i.e., $V_k^1 =$ intersection of all k^{th} terms of all of I's plays from rounds 0 and 1

Claim. $\{V_k^1\}_k$ is an increasing open cover.

Increasing: $V_{k+1}^1 = U_{k+1} \cap \bigcap_{i \leq k} U_{i,k+1} \supseteq U_k \cap (\bigcap_{i < k} U_{ik}) \cap U_{k,k+1} = U_k \cap (\bigcap_{i < k} U_{ik}) = V_k^1.$

Cover: Let $x \in X$. There is k_0 with $x \in U_{k_0}$.

For each $i < k_0$, there is l_i such that $x \in U_{il_i}$.

Let k be greater than k_0 and the l_i 's.

Then V_k^1 is the intersection of (i) U_k ;

Instead, we use the game tree to define covers $\{V_k^n\}_k$ as follows.

Let $V_k^0 = U_k$; then $\{V_k^0\}_k$ is increasing open cover

Next let

$$V_k^1 = U_k \cap U_{0k} \cap U_{1k} \cap \dots \cap U_{k-1,k}.$$

Since we have assumed $U_{ik} \supset U_i$, note that $V_k^1 = \bigcap_{\sigma \in \omega \leq 1} U_{\sigma k}$, i.e., $V_k^1 =$ intersection of all k^{th} terms of all of I's plays from rounds 0 and 1

Claim. $\{V_k^1\}_k$ is an increasing open cover.

Increasing: $V_{k+1}^1 = U_{k+1} \cap \bigcap_{i \leq k} U_{i,k+1} \supseteq U_k \cap (\bigcap_{i < k} U_{ik}) \cap U_{k,k+1} = U_k \cap (\bigcap_{i < k} U_{ik}) = V_k^1.$

Cover: Let $x \in X$. There is k_0 with $x \in U_{k_0}$.

For each $i < k_0$, there is l_i such that $x \in U_{il_i}$.

Let k be greater than k_0 and the l_i 's.

Then V_k^1 is the intersection of (i) U_k ; (ii) $\bigcap_{i < k_0} U_{ik}$,

Instead, we use the game tree to define covers $\{V_k^n\}_k$ as follows.

Let $V_k^0 = U_k$; then $\{V_k^0\}_k$ is increasing open cover

Next let

$$V_k^1 = U_k \cap U_{0k} \cap U_{1k} \cap \dots \cap U_{k-1,k}.$$

Since we have assumed $U_{ik} \supset U_i$, note that $V_k^1 = \bigcap_{\sigma \in \omega \leq 1} U_{\sigma k}$, i.e., $V_k^1 =$ intersection of all k^{th} terms of all of I's plays from rounds 0 and 1

Claim. $\{V_k^1\}_k$ is an increasing open cover.

Increasing: $V_{k+1}^1 = U_{k+1} \cap \bigcap_{i \leq k} U_{i,k+1} \supseteq U_k \cap (\bigcap_{i < k} U_{ik}) \cap U_{k,k+1} = U_k \cap (\bigcap_{i < k} U_{ik}) = V_k^1.$

Cover: Let $x \in X$. There is k_0 with $x \in U_{k_0}$.

For each $i < k_0$, there is l_i such that $x \in U_{il_i}$.

Let k be greater than k_0 and the l_i 's.

Then V_k^1 is the intersection of (i) U_k ; (ii) $\bigcap_{i < k_0} U_{ik}$, and (iii) $\bigcap_{k_0 \leq i < k} U_{il_i}$.

Instead, we use the game tree to define covers $\{V_k^n\}_k$ as follows.

Let $V_k^0 = U_k$; then $\{V_k^0\}_k$ is increasing open cover

Next let

$$V_k^1 = U_k \cap U_{0k} \cap U_{1k} \cap \dots \cap U_{k-1,k}.$$

Since we have assumed $U_{ik} \supset U_i$, note that $V_k^1 = \bigcap_{\sigma \in \omega \leq 1} U_{\sigma k}$, i.e., V_k^1 = intersection of all k^{th} terms of all of I's plays from rounds 0 and 1

Claim. $\{V_k^1\}_k$ is an increasing open cover.

Increasing: $V_{k+1}^1 = U_{k+1} \cap \bigcap_{i \leq k} U_{i,k+1} \supseteq U_k \cap (\bigcap_{i < k} U_{ik}) \cap U_{k,k+1} = U_k \cap (\bigcap_{i < k} U_{ik}) = V_k^1.$

Cover: Let $x \in X$. There is k_0 with $x \in U_{k_0}$.

For each $i < k_0$, there is l_i such that $x \in U_{il_i}$.

Let k be greater than k_0 and the l_i 's.

Then V_k^1 is the intersection of (i) U_k ; (ii) $\bigcap_{i < k_0} U_{ik}$, and (iii) $\bigcap_{k_0 \leq i < k} U_{ik}$.

Now x is in (i) since $k \geq k_0$,

Instead, we use the game tree to define covers $\{V_k^n\}_k$ as follows.

Let $V_k^0 = U_k$; then $\{V_k^0\}_k$ is increasing open cover

Next let

$$V_k^1 = U_k \cap U_{0k} \cap U_{1k} \cap \dots \cap U_{k-1,k}.$$

Since we have assumed $U_{ik} \supset U_i$, note that $V_k^1 = \bigcap_{\sigma \in \omega \leq 1} U_{\sigma k}$, i.e., $V_k^1 =$ intersection of all k^{th} terms of all of I's plays from rounds 0 and 1

Claim. $\{V_k^1\}_k$ is an increasing open cover.

Increasing: $V_{k+1}^1 = U_{k+1} \cap \bigcap_{i \leq k} U_{i,k+1} \supseteq U_k \cap (\bigcap_{i < k} U_{ik}) \cap U_{k,k+1} = U_k \cap (\bigcap_{i < k} U_{ik}) = V_k^1.$

Cover: Let $x \in X$. There is k_0 with $x \in U_{k_0}$.

For each $i < k_0$, there is l_i such that $x \in U_{il_i}$.

Let k be greater than k_0 and the l_i 's.

Then V_k^1 is the intersection of (i) U_k ; (ii) $\bigcap_{i < k_0} U_{ik}$, and (iii) $\bigcap_{k_0 \leq i < k} U_{ik}$.

Now x is in (i) since $k \geq k_0$,

x is in (ii) since $k \geq l_i$ for all $i < k_0$, and

Instead, we use the game tree to define covers $\{V_k^n\}_k$ as follows.

Let $V_k^0 = U_k$; then $\{V_k^0\}_k$ is increasing open cover

Next let

$$V_k^1 = U_k \cap U_{0k} \cap U_{1k} \cap \dots \cap U_{k-1,k}.$$

Since we have assumed $U_{ik} \supset U_i$, note that $V_k^1 = \bigcap_{\sigma \in \omega \leq 1} U_{\sigma k}$, i.e., V_k^1 = intersection of all k^{th} terms of all of I's plays from rounds 0 and 1

Claim. $\{V_k^1\}_k$ is an increasing open cover.

Increasing: $V_{k+1}^1 = U_{k+1} \cap \bigcap_{i \leq k} U_{i,k+1} \supseteq U_k \cap (\bigcap_{i < k} U_{ik}) \cap U_{k,k+1} = U_k \cap (\bigcap_{i < k} U_{ik}) = V_k^1.$

Cover: Let $x \in X$. There is k_0 with $x \in U_{k_0}$.

For each $i < k_0$, there is l_i such that $x \in U_{il_i}$.

Let k be greater than k_0 and the l_i 's.

Then V_k^1 is the intersection of (i) U_k ; (ii) $\bigcap_{i < k_0} U_{ik}$, and (iii) $\bigcap_{k_0 \leq i < k} U_{ik}$.

Now x is in (i) since $k \geq k_0$,

x is in (ii) since $k \geq l_i$ for all $i < k_0$, and

x is in (iii) since $U_{ik} \supseteq U_{k_0}$ for all $i \geq k_0$.

Instead, we use the game tree to define covers $\{V_k^n\}_k$ as follows.

Let $V_k^0 = U_k$; then $\{V_k^0\}_k$ is increasing open cover

Next let

$$V_k^1 = U_k \cap U_{0k} \cap U_{1k} \cap \dots \cap U_{k-1,k}.$$

Since we have assumed $U_{ik} \supset U_i$, note that $V_k^1 = \bigcap_{\sigma \in \omega \leq 1} U_{\sigma k}$, i.e., $V_k^1 =$ intersection of all k^{th} terms of all of I's plays from rounds 0 and 1

Claim. $\{V_k^1\}_k$ is an increasing open cover.

Increasing: $V_{k+1}^1 = U_{k+1} \cap \bigcap_{i \leq k} U_{i,k+1} \supseteq U_k \cap (\bigcap_{i < k} U_{ik}) \cap U_{k,k+1} = U_k \cap (\bigcap_{i < k} U_{ik}) = V_k^1.$

Cover: Let $x \in X$. There is k_0 with $x \in U_{k_0}$.

For each $i < k_0$, there is l_i such that $x \in U_{il_i}$.

Let k be greater than k_0 and the l_i 's.

Then V_k^1 is the intersection of (i) U_k ; (ii) $\bigcap_{i < k_0} U_{ik}$, and (iii) $\bigcap_{k_0 \leq i < k} U_{ik}$.

Now x is in (i) since $k \geq k_0$,

x is in (ii) since $k \geq l_i$ for all $i < k_0$, and

x is in (iii) since $U_{ik} \supseteq U_{k_0}$ for all $i \geq k_0$.

So $x \in V_k^1$.

Similarly, for each n , $\{V_k^n\}_k$ is an increasing open cover, where

$$V_k^n = \bigcap_{\sigma \in \omega^{\leq n}} U_{\sigma \frown k}$$

Similarly, for each n , $\{V_k^n\}_k$ is an increasing open cover, where

$$V_k^n = \bigcap_{\sigma \in \omega^{\leq n}} U_{\sigma \frown k}$$

Since X is Menger, there is $f : \omega \rightarrow \omega$ such that $X = \bigcup_{n \in \omega} V_{f(n)}^n$.

Similarly, for each n , $\{V_k^n\}_k$ is an increasing open cover, where

$$V_k^n = \bigcap_{\sigma \in \omega^{\leq n}} U_{\sigma \frown k}$$

Since X is Menger, there is $f : \omega \rightarrow \omega$ such that $X = \bigcup_{n \in \omega} V_{f(n)}^n$.

But $V_{f(n)}^n \subset U_{(f \upharpoonright n) \frown f(n)} = U_{f \upharpoonright n+1}$.

Similarly, for each n , $\{V_k^n\}_k$ is an increasing open cover, where

$$V_k^n = \bigcap_{\sigma \in \omega^{\leq n}} U_{\sigma \frown k}$$

Since X is Menger, there is $f : \omega \rightarrow \omega$ such that $X = \bigcup_{n \in \omega} V_{f(n)}^n$.

But $V_{f(n)}^n \subset U_{(f \upharpoonright n) \frown f(n)} = U_{f \upharpoonright n+1}$.

So $X = \bigcup_{n \in \omega} U_{f \upharpoonright n+1}$, which corresponds to a play of the game in which I's strategy has been defeated. □

TUTORIAL: Topological games, lecture II

Convergence game

Let X be a space, and H a closed subset of X . Define $G(H, X)$ as follows:

Convergence game

Let X be a space, and H a closed subset of X . Define $G(H, X)$ as follows:
There are two players, O and P .

Convergence game

Let X be a space, and H a closed subset of X . Define $G(H, X)$ as follows:

There are two players, O and P .

In the n th round, O chooses an open $O_n \supset H$, and P chooses a point $p_n \in O_n$.

Convergence game

Let X be a space, and H a closed subset of X . Define $G(H, X)$ as follows:

There are two players, O and P .

In the n th round, O chooses an open $O_n \supset H$, and P chooses a point $p_n \in O_n$.

We say O *wins* the game if $p_n \rightarrow H$ in the sense that every open superset of H contains p_n for all but finitely many $n \in \omega$.

Convergence game

Let X be a space, and H a closed subset of X . Define $G(H, X)$ as follows:

There are two players, O and P .

In the n th round, O chooses an open $O_n \supset H$, and P chooses a point $p_n \in O_n$.

We say O *wins* the game if $p_n \rightarrow H$ in the sense that every open superset of H contains p_n for all but finitely many $n \in \omega$.

If $O \uparrow G(H, X)$, we call H a *W-set* in X .

Clearly X first countable implies every point of X is a W -set in X .

Clearly X first countable implies every point of X is a W -set in X . Spaces in which O has a winning strategy at every point are called W -spaces.

Clearly X first countable implies every point of X is a W -set in X . Spaces in which O has a winning strategy at every point are called W -spaces.

A W -space which is not first countable:

Clearly X first countable implies every point of X is a W -set in X . Spaces in which O has a winning strategy at every point are called W -spaces.

A W -space which is not first countable:

The one-point compactification of uncountable discrete space

Clearly X first countable implies every point of X is a W -set in X .
Spaces in which O has a winning strategy at every point are called W -spaces.

A W -space which is not first countable:

The one-point compactification of uncountable discrete space

I called spaces in which P does not have a winning strategy w -spaces.

Clearly X first countable implies every point of X is a W -set in X . Spaces in which O has a winning strategy at every point are called W -spaces.

A W -space which is not first countable:

The one-point compactification of uncountable discrete space

I called spaces in which P does not have a winning strategy w -spaces.

P . Sharma proved X is a w -space iff for each $x \in X$:

if $x \in \overline{A_n}$ for each $n \in \omega$, then there are $x_n \in A_n$ with $x_n \rightarrow x$.

Clearly X first countable implies every point of X is a W -set in X . Spaces in which O has a winning strategy at every point are called W -spaces.

A W -space which is not first countable:

The one-point compactification of uncountable discrete space

I called spaces in which P does not have a winning strategy w -spaces.

P. Sharma proved X is a w -space iff for each $x \in X$:

if $x \in \overline{A_n}$ for each $n \in \omega$, then there are $x_n \in A_n$ with $x_n \rightarrow x$.

This showed the class of w -spaces equivalent to a class introduced by Arhangel'skii (Fréchet α_2 -spaces).

Theorem

- 1 *First-countable* \Rightarrow *W-space* \Rightarrow *w-space* \Rightarrow *Fréchet*;
- 2 *W-spaces are hereditary and countably productive (closed under Σ -products, even).*

Theorem

- 1 *First-countable* \Rightarrow *W-space* \Rightarrow *w-space* \Rightarrow *Fréchet*;
- 2 *W-spaces are hereditary and countably productive (closed under Σ -products, even).*

Consider the game $G(H, X)$ played in a **compact Hausdorff space X** , where $H \subset X$ is closed.

Theorem

- 1 $\text{First-countable} \Rightarrow W\text{-space} \Rightarrow w\text{-space} \Rightarrow \text{Fréchet};$
- 2 $W\text{-spaces are hereditary and countably productive (closed under } \Sigma\text{-products, even).}$

Consider the game $G(H, X)$ played in a **compact Hausdorff space** X , where $H \subset X$ is closed.

O will have a winning strategy if H has a countable “outer base”, i.e., there is a countable collection of open supersets of H such that every open superset of H contains one. By compactness, H has a countable outer base iff H is G_δ .

Theorem

- 1 $\text{First-countable} \Rightarrow W\text{-space} \Rightarrow w\text{-space} \Rightarrow \text{Fréchet};$
- 2 $W\text{-spaces are hereditary and countably productive (closed under } \Sigma\text{-products, even).}$

Consider the game $G(H, X)$ played in a **compact Hausdorff space** X , where $H \subset X$ is closed.

O will have a winning strategy if H has a countable “outer base”, i.e., there is a countable collection of open supersets of H such that every open superset of H contains one. By compactness, H has a countable outer base iff H is G_δ .

A classical result:

Theorem (Schneider)

A compact Hausdorff space X is metrizable iff the diagonal Δ of X is G_δ in X^2 .

O can have a winning strategy in $G(H, X)$ for compact X without H being G_δ :

O can have a winning strategy in $G(H, X)$ for compact X without H being G_δ :
consider one-point compactification of uncountable discrete space.

O can have a winning strategy in $G(H, X)$ for compact X without H being G_δ :

consider one-point compactification of uncountable discrete space.

To say that O has winning strategy in $G(\Delta, X^2)$ is weaker than to say the diagonal is G_δ in X^2 .

O can have a winning strategy in $G(H, X)$ for compact X without H being G_δ :

consider one-point compactification of uncountable discrete space.

To say that O has winning strategy in $G(\Delta, X^2)$ is weaker than to say the diagonal is G_δ in X^2 .

Definition. Compact X is [Corson compact](#) iff

O can have a winning strategy in $G(H, X)$ for compact X without H being G_δ :

consider one-point compactification of uncountable discrete space.

To say that O has winning strategy in $G(\Delta, X^2)$ is weaker than to say the diagonal is G_δ in X^2 .

Definition. Compact X is **Corson compact** iff

- X is homeo to subspace of

$$\Sigma \mathbb{R}^\kappa = \{x \in \mathbb{R}^\kappa : |\{\alpha < \kappa : x(\alpha) \neq 0\}| \leq \omega\}$$

or equivalently

O can have a winning strategy in $G(H, X)$ for compact X without H being G_δ :

consider one-point compactification of uncountable discrete space.

To say that O has winning strategy in $G(\Delta, X^2)$ is weaker than to say the diagonal is G_δ in X^2 .

Definition. Compact X is **Corson compact** iff

- X is homeo to subspace of

$$\Sigma \mathbb{R}^\kappa = \{x \in \mathbb{R}^\kappa : |\{\alpha < \kappa : x(\alpha) \neq 0\}| \leq \omega\}$$

or equivalently

- X has a point-countable T_0 -separating cover by open F_σ 's.

O can have a winning strategy in $G(H, X)$ for compact X without H being G_δ :

consider one-point compactification of uncountable discrete space.

To say that O has winning strategy in $G(\Delta, X^2)$ is weaker than to say the diagonal is G_δ in X^2 .

Definition. Compact X is **Corson compact** iff

- X is homeo to subspace of

$$\Sigma \mathbb{R}^\kappa = \{x \in \mathbb{R}^\kappa : |\{\alpha < \kappa : x(\alpha) \neq 0\}| \leq \omega\}$$

or equivalently

- X has a point-countable T_0 -separating cover by open F_σ 's.

Theorem(G.G., 1984)

A compact space X is Corson compact iff $O \uparrow G(\Delta, X^2)$.

Eberlein compact: homeo to weakly compact subset of Banach space

Eberlein compact: homeo to weakly compact subset of Banach space

Eberlein \Rightarrow Corson

Eberlein compact: homeo to weakly compact subset of Banach space

Eberlein \Rightarrow Corson

Theorem(G.G., 1986)

A compact X is Eberlein compact iff O has a winning strategy in $G(\Delta, X^2)$ which depends only on P 's last move and the number of the move (i.e., a Markov strategy).

Eberlein compact: homeo to weakly compact subset of Banach space

Eberlein \Rightarrow Corson

Theorem(G.G., 1986)

A compact X is Eberlein compact iff O has a winning strategy in $G(\Delta, X^2)$ which depends only on P 's last move and the number of the move (i.e., a Markov strategy).

Strong Eberlein = scattered Eberlein

Eberlein compact: homeo to weakly compact subset of Banach space

Eberlein \Rightarrow Corson

Theorem(G.G., 1986)

A compact X is Eberlein compact iff O has a winning strategy in $G(\Delta, X^2)$ which depends only on P 's last move and the number of the move (i.e., a Markov strategy).

Strong Eberlein = scattered Eberlein

Theorem(G.G., 1984)

A compact scattered space X is strong Eberlein compact iff X is a W -space.

A recent use of the Corson result.

A recent use of the Corson result.

DEF (Tkachuk): A space X is **monotonically monolithic** if one can assign to each $F \in [X]^{<\omega}$ a countable collection $\mathcal{N}(F)$ of subsets of X such that

A recent use of the Corson result.

DEF (Tkachuk): A space X is **monotonically monolithic** if one can assign to each $F \in [X]^{<\omega}$ a countable collection $\mathcal{N}(F)$ of subsets of X such that

$$\textcircled{1} \quad F \subset F' \Rightarrow \mathcal{N}(F) \subset \mathcal{N}(F');$$

A recent use of the Corson result.

DEF (Tkachuk): A space X is **monotonically monolithic** if one can assign to each $F \in [X]^{<\omega}$ a countable collection $\mathcal{N}(F)$ of subsets of X such that

- 1 $F \subset F' \Rightarrow \mathcal{N}(F) \subset \mathcal{N}(F')$;
- 2 If U open and $x \in \overline{A} \cap U$ there is $F \in [A]^{<\omega}$ and $N \in \mathcal{N}(F)$ with $x \in N \subset U$.

A recent use of the Corson result.

DEF (Tkachuk): A space X is **monotonically monolithic** if one can assign to each $F \in [X]^{<\omega}$ a countable collection $\mathcal{N}(F)$ of subsets of X such that

- 1 $F \subset F' \Rightarrow \mathcal{N}(F) \subset \mathcal{N}(F')$;
- 2 If U open and $x \in \overline{A} \cap U$ there is $F \in [A]^{<\omega}$ and $N \in \mathcal{N}(F)$ with $x \in N \subset U$.

(That is, $\bigcup_{F \in [A]^{<\omega}} \mathcal{N}(F)$ includes a network at every point of \overline{A} .)

Tkachuk proved:

- 1 Point-countable base \Rightarrow monotonically monolithic

Tkachuk proved:

- ① Point-countable base \Rightarrow monotonically monolithic
- ② X Lindelöf $\Sigma \Rightarrow C_p(X)$ monotonically monolithic

Tkachuk proved:

- 1 Point-countable base \Rightarrow monotonically monolithic
- 2 X Lindelöf $\Sigma \Rightarrow C_p(X)$ monotonically monolithic (Lindelöf $\Sigma =$ continuous image of closed subspace of separable metric \times compact)

Tkachuk proved:

- 1 Point-countable base \Rightarrow monotonically monolithic
- 2 X Lindelöf $\Sigma \Rightarrow C_p(X)$ monotonically monolithic (Lindelöf $\Sigma =$ continuous image of closed subspace of separable metric \times compact)
- 3 Monotonically monolithic \Rightarrow hereditarily D .

D-space: $x \in N(x)^\circ \forall x \in X \Rightarrow \exists$ closed discrete D with $\{N(x) : x \in D\}$ covering X

Tkachuk proved:

- 1 Point-countable base \Rightarrow monotonically monolithic
- 2 X Lindelöf $\Sigma \Rightarrow C_p(X)$ monotonically monolithic (Lindelöf $\Sigma =$ continuous image of closed subspace of separable metric \times compact)
- 3 Monotonically monolithic \Rightarrow hereditarily D .

D-space: $x \in N(x)^\circ \forall x \in X \Rightarrow \exists$ closed discrete D with $\{N(x) : x \in D\}$ covering X

X is [Gul'ko compact](#) iff X compact and $C_p(X)$ is Lindelöf Σ . It follows that:

Tkachuk proved:

- 1 Point-countable base \Rightarrow monotonically monolithic
- 2 X Lindelöf $\Sigma \Rightarrow C_p(X)$ monotonically monolithic (Lindelöf $\Sigma =$ continuous image of closed subspace of separable metric \times compact)
- 3 Monotonically monolithic \Rightarrow hereditarily D .

D-space: $x \in N(x)^\circ \forall x \in X \Rightarrow \exists$ closed discrete D with $\{N(x) : x \in D\}$ covering X

X is **Gul'ko compact** iff X compact and $C_p(X)$ is Lindelöf Σ . It follows that:

Gul'ko compact \Rightarrow monotonically monolithic

Tkachuk proved:

- 1 Point-countable base \Rightarrow monotonically monolithic
- 2 X Lindelöf $\Sigma \Rightarrow C_p(X)$ monotonically monolithic (Lindelöf $\Sigma =$ continuous image of closed subspace of separable metric \times compact)
- 3 Monotonically monolithic \Rightarrow hereditarily D .

D-space: $x \in N(x)^\circ \forall x \in X \Rightarrow \exists$ closed discrete D with $\{N(x) : x \in D\}$ covering X

X is **Gul'ko compact** iff X compact and $C_p(X)$ is Lindelöf Σ . It follows that:

Gul'ko compact \Rightarrow monotonically monolithic

Eberlein compact \Rightarrow Gul'ko compact \Rightarrow Corson compact

Tkachuk proved:

- 1 Point-countable base \Rightarrow monotonically monolithic
- 2 X Lindelöf $\Sigma \Rightarrow C_p(X)$ monotonically monolithic (Lindelöf $\Sigma =$ continuous image of closed subspace of separable metric \times compact)
- 3 Monotonically monolithic \Rightarrow hereditarily D .

D-space: $x \in N(x)^\circ \forall x \in X \Rightarrow \exists$ closed discrete D with $\{N(x) : x \in D\}$ covering X

X is **Gul'ko compact** iff X compact and $C_p(X)$ is Lindelöf Σ . It follows that:

Gul'ko compact \Rightarrow monotonically monolithic

Eberlein compact \Rightarrow Gul'ko compact \Rightarrow Corson compact
 \Rightarrow hereditarily D

Question(Tkachuk)

- Monotonically monolithic compact \Rightarrow Corson compact?

Question(Tkachuk)

- Monotonically monolithic compact \Rightarrow Corson compact?

Yes:

Theorem (G.G., 2010)

If X is compact and monotonically monolithic, then X is Corson compact.

Lemma

Suppose X is compact and monotonically monolithic. Then O has a winning strategy in $G(H, X)$ for any closed $H \subset X$.

Lemma

Suppose X is compact and monotonically monolithic. Then O has a winning strategy in $G(H, X)$ for any closed $H \subset X$.

To prove the Theorem from the Lemma:

Lemma

Suppose X is compact and monotonically monolithic. Then O has a winning strategy in $G(H, X)$ for any closed $H \subset X$.

To prove the Theorem from the Lemma:

X compact monotonically monolithic \Rightarrow ditto for $X^2 \Rightarrow O$ has winning strategy in $G(\Delta, X^2) \Rightarrow X$ Corson compact

Proof of lemma

Proof of lemma

X compact, \mathcal{N} witnesses monotonically monolithic.

Proof of lemma

X compact, \mathcal{N} witnesses monotonically monolithic.

$H \subset X$ closed

Proof of lemma

X compact, \mathcal{N} witnesses monotonically monolithic.

$H \subset X$ closed

O plays $O_0 = X$, P plays $p_0 \in X$.

Proof of lemma

X compact, \mathcal{N} witnesses monotonically monolithic.

$H \subset X$ closed

O plays $O_0 = X$, P plays $p_0 \in X$.

O looks at $\mathcal{N}(\{p_0\}) = \{N_{00}, N_{01}, \dots\}$

Proof of lemma

X compact, \mathcal{N} witnesses monotonically monolithic.

$H \subset X$ closed

O plays $O_0 = X$, P plays $p_0 \in X$.

O looks at $\mathcal{N}(\{p_0\}) = \{N_{00}, N_{01}, \dots\}$

O chooses open $O_1 \supset H$ s.t. $\overline{O_1} \cap N_{00} = \emptyset$ if such O_1 exists; else $O_1 = X$.

Proof of lemma

X compact, \mathcal{N} witnesses monotonically monolithic.

$H \subset X$ closed

O plays $O_0 = X$, P plays $p_0 \in X$.

O looks at $\mathcal{N}(\{p_0\}) = \{N_{00}, N_{01}, \dots\}$

O chooses open $O_1 \supset H$ s.t. $\overline{O_1} \cap N_{00} = \emptyset$ if such O_1 exists; else $O_1 = X$.

P chooses $p_1 \in O_1$. Let $\mathcal{N}(\{p_0, p_1\}) = \{N_{10}, N_{11}, \dots\}$

Proof of lemma

X compact, \mathcal{N} witnesses monotonically monolithic.

$H \subset X$ closed

O plays $O_0 = X$, P plays $p_0 \in X$.

O looks at $\mathcal{N}(\{p_0\}) = \{N_{00}, N_{01}, \dots\}$

O chooses open $O_1 \supset H$ s.t. $\overline{O_1} \cap N_{00} = \emptyset$ if such O_1 exists; else $O_1 = X$.

P chooses $p_1 \in O_1$. Let $\mathcal{N}(\{p_0, p_1\}) = \{N_{10}, N_{11}, \dots\}$

O chooses $O_2 \subset O_1$ s.t., whenever possible for $i, j < 2$, $\overline{O_2} \cap N_{ij} = \emptyset$.

Proof of lemma

X compact, \mathcal{N} witnesses monotonically monolithic.

$H \subset X$ closed

O plays $O_0 = X$, P plays $p_0 \in X$.

O looks at $\mathcal{N}(\{p_0\}) = \{N_{00}, N_{01}, \dots\}$

O chooses open $O_1 \supset H$ s.t. $\overline{O_1} \cap N_{00} = \emptyset$ if such O_1 exists; else $O_1 = X$.

P chooses $p_1 \in O_1$. Let $\mathcal{N}(\{p_0, p_1\}) = \{N_{10}, N_{11}, \dots\}$

O chooses $O_2 \subset O_1$ s.t., whenever possible for $i, j < 2$, $\overline{O_2} \cap N_{ij} = \emptyset$.

Etc

Claim. This wins for O.

Claim. This wins for O .

Suppose $p_n \not\rightarrow H$. Then $\{p_n\}_n$ has limit point $q \notin H$

Claim. This wins for O .

Suppose $p_n \not\rightarrow H$. Then $\{p_n\}_n$ has limit point $q \notin H$

Let $q \in U$ open, $\overline{U} \cap H = \emptyset$.

Claim. This wins for O .

Suppose $p_n \not\rightarrow H$. Then $\{p_n\}_n$ has limit point $q \notin H$

Let $q \in U$ open, $\overline{U} \cap H = \emptyset$.

$\exists k \in \omega$ with $N \in \mathcal{N}(\{p_i\}_{i \leq k})$ and $q \in N \subset U$

Claim. This wins for O .

Suppose $p_n \not\rightarrow H$. Then $\{p_n\}_n$ has limit point $q \notin H$

Let $q \in U$ open, $\overline{U} \cap H = \emptyset$.

$\exists k \in \omega$ with $N \in \mathcal{N}(\{p_i\}_{i \leq k})$ and $q \in N \subset U$

$N = N_{kj}$ for some k, j

Claim. This wins for O.

Suppose $p_n \not\rightarrow H$. Then $\{p_n\}_n$ has limit point $q \notin H$

Let $q \in U$ open, $\bar{U} \cap H = \emptyset$.

$\exists k \in \omega$ with $N \in \mathcal{N}(\{p_i\}_{i \leq k})$ and $q \in N \subset U$

$N = N_{kj}$ for some k, j

$\bar{O}_n \cap N = \emptyset$ for $n > \max\{j, k\}$

Claim. This wins for O .

Suppose $p_n \not\rightarrow H$. Then $\{p_n\}_n$ has limit point $q \notin H$

Let $q \in U$ open, $\bar{U} \cap H = \emptyset$.

$\exists k \in \omega$ with $N \in \mathcal{N}(\{p_i\}_{i \leq k})$ and $q \in N \subset U$

$N = N_{kj}$ for some k, j

$\bar{O}_n \cap N = \emptyset$ for $n > \max\{j, k\}$

$q \notin \bar{O}_n \Rightarrow q$ not limit of $\{p_n\}_n$.

Contradiction.

How to show O wins in $G(\Delta, X^2)$ implies X Corson compact.

How to show O wins in $G(\Delta, X^2)$ implies X Corson compact.

First, easy to get from this that O wins in $G(x, X)$ for every $x \in X$; i.e., X is a W -space hence Fréchet (and so is X^2).

How to show O wins in $G(\Delta, X^2)$ implies X Corson compact.

First, easy to get from this that O wins in $G(x, X)$ for every $x \in X$; i.e., X is a W -space hence Fréchet (and so is X^2).

Use the point-countable T_0 -separating open cover characterization. Can get such if can show $X^2 \setminus \Delta$ is metalindelöf, i.e., every open cover has a point-countable open refinement.

How to show O wins in $G(\Delta, X^2)$ implies X Corson compact.

First, easy to get from this that O wins in $G(x, X)$ for every $x \in X$; i.e., X is a W -space hence Fréchet (and so is X^2).

Use the point-countable T_0 -separating open cover characterization. Can get such if can show $X^2 \setminus \Delta$ is metalindelöf, i.e., every open cover has a point-countable open refinement.

Then we have:

Theorem

Let X be compact and countably tight, and H closed. Then O has a winning strategy in $G(H, X)$ iff $X \setminus H$ is metalindelöf.

It is useful to view the game as a game in $X \setminus H$, with players K and P.

It is useful to view the game as a game in $X \setminus H$, with players K and P.

In the n^{th} round, K chooses a compact $K_n \subset X \setminus H$ (the complement of a play by O), and P responds with a point $p_n \notin K_n$.

It is useful to view the game as a game in $X \setminus H$, with players K and P.

In the n^{th} round, K chooses a compact $K_n \subset X \setminus H$ (the complement of a play by O), and P responds with a point $p_n \notin K_n$.

K *wins* if $p_n \rightarrow \infty$ (i.e., $\{p_n : n \in \omega\}$ is closed discrete in $X \setminus H$).

Replacing $X \setminus H$ with X , let us denote this game by $G_{K,P}(X)$.

It is useful to view the game as a game in $X \setminus H$, with players K and P .

In the n^{th} round, K chooses a compact $K_n \subset X \setminus H$ (the complement of a play by O), and P responds with a point $p_n \notin K_n$.

K wins if $p_n \rightarrow \infty$ (i.e., $\{p_n : n \in \omega\}$ is closed discrete in $X \setminus H$).

Replacing $X \setminus H$ with X , let us denote this game by $G_{K,P}(X)$.

Then the result becomes:

Theorem

Let X be locally compact and countably tight. Then K has a winning strategy in $G_{K,P}(X)$ iff X is metalindelöf.

(I don't know if the countable tightness assumption is necessary.)

Outline of proof, elementary submodel style

Outline of proof, elementary submodel style

Proof. If X is metalindelöf, then there is a point-countable cover \mathcal{U} of X by open sets with compact closures .

Outline of proof, elementary submodel style

Proof. If X is metalindelöf, then there is a point-countable cover \mathcal{U} of X by open sets with compact closures .

K wins by looking at the countably many members of \mathcal{U} containing P 's chosen point at each round, and choosing an increasing sequence of compact sets that eventually cover every one of these members of \mathcal{U} . It is easy to check that this wins for K .

Outline of proof, elementary submodel style

Proof. If X is metalindelöf, then there is a point-countable cover \mathcal{U} of X by open sets with compact closures .

K wins by looking at the countably many members of \mathcal{U} containing P 's chosen point at each round, and choosing an increasing sequence of compact sets that eventually cover every one of these members of \mathcal{U} . It is easy to check that this wins for K .

Now suppose K has a winning strategy σ , and let \mathcal{U} be a cover of X by open sets with compact closures. Let M be an elementary submodel (of some sufficiently large $H(\theta)$) with $X, \mathcal{U}, \sigma \in M$.

Key Claim. $\overline{M \cap X} \subset \bigcup (M \cap \mathcal{U})$.

Key Claim. $\overline{M \cap X} \subset \bigcup(M \cap \mathcal{U})$.

Proof of Key Claim. Suppose $p \in \overline{M \cap X} \setminus \bigcup(M \cap \mathcal{U})$. Let $p \in U_p \in \mathcal{U}$.

Key Claim. $\overline{M \cap X} \subset \bigcup(M \cap \mathcal{U})$.

Proof of Key Claim. Suppose $p \in \overline{M \cap X} \setminus \bigcup(M \cap \mathcal{U})$. Let $p \in U_p \in \mathcal{U}$.

Suppose $F = \{p_0, p_1, \dots, p_n\} \subset U_p \cap (M \cap X)$. Then $\sigma(F)$ is compact and in M so there exists a finite $\mathcal{U}_0 \subset \mathcal{U}$ in M covering $\sigma(F)$.

Key Claim. $\overline{M \cap X} \subset \bigcup(M \cap \mathcal{U})$.

Proof of Key Claim. Suppose $p \in \overline{M \cap X} \setminus \bigcup(M \cap \mathcal{U})$. Let $p \in U_p \in \mathcal{U}$.

Suppose $F = \{p_0, p_1, \dots, p_n\} \subset U_p \cap (M \cap X)$. Then $\sigma(F)$ is compact and in M so there exists a finite $\mathcal{U}_0 \subset \mathcal{U}$ in M covering $\sigma(F)$.

Since M also contains a finite subset of \mathcal{U} covering $\overline{\bigcup \mathcal{U}_0}$, we have $p \notin \overline{\bigcup \mathcal{U}_0}$. So there exists $p_{n+1} \in U_p \cap (M \cap X) \setminus \overline{\bigcup \mathcal{U}_0}$.

Key Claim. $\overline{M \cap X} \subset \bigcup(M \cap \mathcal{U})$.

Proof of Key Claim. Suppose $p \in \overline{M \cap X} \setminus \bigcup(M \cap \mathcal{U})$. Let $p \in U_p \in \mathcal{U}$.

Suppose $F = \{p_0, p_1, \dots, p_n\} \subset U_p \cap (M \cap X)$. Then $\sigma(F)$ is compact and in M so there exists a finite $\mathcal{U}_0 \subset \mathcal{U}$ in M covering $\sigma(F)$.

Since M also contains a finite subset of \mathcal{U} covering $\overline{\bigcup \mathcal{U}_0}$, we have $p \notin \overline{\bigcup \mathcal{U}_0}$. So there exists $p_{n+1} \in U_p \cap (M \cap X) \setminus \overline{\bigcup \mathcal{U}_0}$.

It follows that if K uses the strategy σ , P can always choose a point in $U_p \cap (M \cap X)$. But then K loses the game, a contradiction which completes the proof of Key Claim.

Since there is an M with $\mathcal{U} \subset M$, the next claim completes the proof of the theorem.

Since there is an M with $\mathcal{U} \subset M$, the next claim completes the proof of the theorem.

Claim 2. There is a point-countable open refinement \mathcal{V}_M of $M \cap \mathcal{U}$ covering $\cup(M \cap \mathcal{U})$.

Since there is an M with $\mathcal{U} \subset M$, the next claim completes the proof of the theorem.

Claim 2. There is a point-countable open refinement \mathcal{V}_M of $M \cap \mathcal{U}$ covering $\cup(M \cap \mathcal{U})$.

Proof of Claim 2. By induction on $|M| = \kappa$. Write $M = \bigcup\{M_\alpha : \alpha < \kappa\}$ and use Key Claim to put together point-countable refinements of $M_\alpha \cap \mathcal{U}$.

$G_{K,L}(X)$ is defined just like $G_{K,P}(X)$, except that P , who will be renamed L , chooses compact sets instead of points, i.e., L 's n^{th} play is a compact set L_n missing K 's previous move K_n .

$G_{K,L}(X)$ is defined just like $G_{K,P}(X)$, except that P , who will be renamed L , chooses compact sets instead of points, i.e., L 's n^{th} play is a compact set L_n missing K 's previous move K_n .

K wins iff $\{L_i\}_{i \in \omega}$ is a discrete collection.

$G_{K,L}(X)$ is defined just like $G_{K,P}(X)$, except that P , who will be renamed L , chooses compact sets instead of points, i.e., L 's n^{th} play is a compact set L_n missing K 's previous move K_n .

K wins iff $\{L_i\}_{i \in \omega}$ is a discrete collection.

$G_{K,L}^{\circ}(X)$ is the same as $G_{K,L}(X)$, except that K wins iff $\{L_i\}_{i \in \omega}$ has a discrete open expansion.

$G_{K,L}(X)$ is defined just like $G_{K,P}(X)$, except that P , who will be renamed L , chooses compact sets instead of points, i.e., L 's n^{th} play is a compact set L_n missing K 's previous move K_n .

K wins iff $\{L_i\}_{i \in \omega}$ is a discrete collection.

$G_{K,L}^{\circ}(X)$ is the same as $G_{K,L}(X)$, except that K wins iff $\{L_i\}_{i \in \omega}$ has a discrete open expansion.

It is easy to see that K has a winning strategy in any locally compact σ -compact space: K simply chooses at the n^{th} play the n^{th} set in an increasing sequence of compact sets whose interiors cover the space.

$G_{K,L}(X)$ is defined just like $G_{K,P}(X)$, except that P , who will be renamed L , chooses compact sets instead of points, i.e., L 's n^{th} play is a compact set L_n missing K 's previous move K_n .

K wins iff $\{L_i\}_{i \in \omega}$ is a discrete collection.

$G_{K,L}^{\circ}(X)$ is the same as $G_{K,L}(X)$, except that K wins iff $\{L_i\}_{i \in \omega}$ has a discrete open expansion.

It is easy to see that K has a winning strategy in any locally compact σ -compact space: K simply chooses at the n^{th} play the n^{th} set in an increasing sequence of compact sets whose interiors cover the space.

It is nearly as easy to see that K wins if X is a topological sum of locally compact σ -compact spaces, i.e., whenever X is locally compact and paracompact. The next theorem shows we have an equivalence:

Theorem

Let X be a locally compact space. Then the following are equivalent:

- 1 $K \uparrow G_{K,L}(X)$;
- 2 $K \uparrow G_{K,L}^{\circ}(X)$;
- 3 X is paracompact.

Why $G_{K,L}^o(X)$? Because it is the most natural one for attacking the following open problem:

Why $G_{K,L}^o(X)$? Because it is the most natural one for attacking the following open problem:

Question

For what (completely regular) spaces X is $C_k(X)$ a Baire space?

($C_k(X)$ is the space of continuous real-valued functions on X with the compact-open topology.)

Theorem (McCoy, Ntantu)

- 1 If $NE \uparrow BM(C_k(X))$ then $K \uparrow G_{K,L}^{\circ}(X)$;
- 2 If $C_k(X)$ is Baire, then $L \not\uparrow G_{K,L}^{\circ}(X)$;
- 3 If X is locally compact, then $NE \uparrow BM(C_k(X))$ iff $K \uparrow G_{K,L}^{\circ}(X)$.

Theorem (McCoy, Ntantu)

- 1 If $NE \uparrow BM(C_k(X))$ then $K \uparrow G_{K,L}^{\circ}(X)$;
- 2 If $C_k(X)$ is Baire, then $L \not\uparrow G_{K,L}^{\circ}(X)$;
- 3 If X is locally compact, then $NE \uparrow BM(C_k(X))$ iff $K \uparrow G_{K,L}^{\circ}(X)$.

Proof of(2) Suppose $L \uparrow G_{K,L}^{\circ}(X)$

Theorem (McCoy, Ntantu)

- 1 If $NE \uparrow BM(C_k(X))$ then $K \uparrow G_{K,L}^{\circ}(X)$;
- 2 If $C_k(X)$ is Baire, then $L \not\uparrow G_{K,L}^{\circ}(X)$;
- 3 If X is locally compact, then $NE \uparrow BM(C_k(X))$ iff $K \uparrow G_{K,L}^{\circ}(X)$.

Proof of(2) Suppose $L \uparrow G_{K,L}^{\circ}(X)$

Claim. $E \uparrow BM(C_k(X))$ (so $C_k(X)$ not Baire, contradiction).

Theorem (McCoy, Ntantu)

- 1 If $NE \uparrow BM(C_k(X))$ then $K \uparrow G_{K,L}^o(X)$;
- 2 If $C_k(X)$ is Baire, then $L \not\uparrow G_{K,L}^o(X)$;
- 3 If X is locally compact, then $NE \uparrow BM(C_k(X))$ iff $K \uparrow G_{K,L}^o(X)$.

Proof of(2) Suppose $L \uparrow G_{K,L}^o(X)$

Claim. $E \uparrow BM(C_k(X))$ (so $C_k(X)$ not Baire, contradiction).

W.l.o.g., in the n^{th} round, Non-empty chooses a basic open set of the form $B(K_n, f_n, \epsilon_n) = \{g \in C_k(X) : \forall x \in K_n (|g(x) - f_n(x)| < \epsilon_n)\}$, where K_n is compact.

Theorem (McCoy, Ntantu)

- 1 If $NE \uparrow BM(C_k(X))$ then $K \uparrow G_{K,L}^o(X)$;
- 2 If $C_k(X)$ is Baire, then $L \not\uparrow G_{K,L}^o(X)$;
- 3 If X is locally compact, then $NE \uparrow BM(C_k(X))$ iff $K \uparrow G_{K,L}^o(X)$.

Proof of(2) Suppose $L \uparrow G_{K,L}^o(X)$

Claim. $E \uparrow BM(C_k(X))$ (so $C_k(X)$ not Baire, contradiction).

W.l.o.g., in the n^{th} round, Non-empty chooses a basic open set of the form $B(K_n, f_n, \epsilon_n) = \{g \in C_k(X) : \forall x \in K_n (|g(x) - f_n(x)| < \epsilon_n)\}$, where K_n is compact.

Let L_n be L 's response to K_n in $G_{K,L}^o(X)$ using a winning strategy.

Theorem (McCoy, Ntantu)

- 1 If $NE \uparrow BM(C_k(X))$ then $K \uparrow G_{K,L}^o(X)$;
- 2 If $C_k(X)$ is Baire, then $L \not\uparrow G_{K,L}^o(X)$;
- 3 If X is locally compact, then $NE \uparrow BM(C_k(X))$ iff $K \uparrow G_{K,L}^o(X)$.

Proof of(2) Suppose $L \uparrow G_{K,L}^o(X)$

Claim. $E \uparrow BM(C_k(X))$ (so $C_k(X)$ not Baire, contradiction).

W.l.o.g., in the n^{th} round, Non-empty chooses a basic open set of the form $B(K_n, f_n, \epsilon_n) = \{g \in C_k(X) : \forall x \in K_n (|g(x) - f_n(x)| < \epsilon_n)\}$, where K_n is compact.

Let L_n be L 's response to K_n in $G_{K,L}^o(X)$ using a winning strategy.

Then *Empty* plays $B(K_n, f_n, \epsilon_n) \cap B(L_n, c_n, 1/3)$, where $c_n = \text{constant } n$

Theorem (McCoy, Ntantu)

- 1 If $\text{NE} \uparrow \text{BM}(C_k(X))$ then $K \uparrow G_{K,L}^o(X)$;
- 2 If $C_k(X)$ is Baire, then $L \not\uparrow G_{K,L}^o(X)$;
- 3 If X is locally compact, then $\text{NE} \uparrow \text{BM}(C_k(X))$ iff $K \uparrow G_{K,L}^o(X)$.

Proof of(2) Suppose $L \uparrow G_{K,L}^o(X)$

Claim. $E \uparrow \text{BM}(C_k(X))$ (so $C_k(X)$ not Baire, contradiction).

W.l.o.g., in the n^{th} round, Non-empty chooses a basic open set of the form $B(K_n, f_n, \epsilon_n) = \{g \in C_k(X) : \forall x \in K_n (|g(x) - f_n(x)| < \epsilon_n)\}$, where K_n is compact.

Let L_n be L 's response to K_n in $G_{K,L}^o(X)$ using a winning strategy.

Then *Empty* plays $B(K_n, f_n, \epsilon_n) \cap B(L_n, c_n, 1/3)$, where $c_n = \text{constant}$ n

Suppose $\phi \in \bigcap_{n \in \omega} B(L_n, c_n, 1/3)$.

Theorem (McCoy, Ntantu)

- 1 If $NE \uparrow BM(C_k(X))$ then $K \uparrow G_{K,L}^o(X)$;
- 2 If $C_k(X)$ is Baire, then $L \not\uparrow G_{K,L}^o(X)$;
- 3 If X is locally compact, then $NE \uparrow BM(C_k(X))$ iff $K \uparrow G_{K,L}^o(X)$.

Proof of(2) Suppose $L \uparrow G_{K,L}^o(X)$

Claim. $E \uparrow BM(C_k(X))$ (so $C_k(X)$ not Baire, contradiction).

W.l.o.g., in the n^{th} round, Non-empty chooses a basic open set of the form $B(K_n, f_n, \epsilon_n) = \{g \in C_k(X) : \forall x \in K_n (|g(x) - f_n(x)| < \epsilon_n)\}$, where K_n is compact.

Let L_n be L 's response to K_n in $G_{K,L}^o(X)$ using a winning strategy.

Then *Empty* plays $B(K_n, f_n, \epsilon_n) \cap B(L_n, c_n, 1/3)$, where $c_n = \text{constant } n$
Suppose $\phi \in \bigcap_{n \in \omega} B(L_n, c_n, 1/3)$. Then $\phi(L_n) \subset (n - 1/3, n + 1/3)$ for all n .

Theorem (McCoy, Ntantu)

- 1 If $NE \uparrow BM(C_k(X))$ then $K \uparrow G_{K,L}^o(X)$;
- 2 If $C_k(X)$ is Baire, then $L \not\uparrow G_{K,L}^o(X)$;
- 3 If X is locally compact, then $NE \uparrow BM(C_k(X))$ iff $K \uparrow G_{K,L}^o(X)$.

Proof of(2) Suppose $L \uparrow G_{K,L}^o(X)$

Claim. $E \uparrow BM(C_k(X))$ (so $C_k(X)$ not Baire, contradiction).

W.l.o.g., in the n^{th} round, Non-empty chooses a basic open set of the form $B(K_n, f_n, \epsilon_n) = \{g \in C_k(X) : \forall x \in K_n (|g(x) - f_n(x)| < \epsilon_n)\}$, where K_n is compact.

Let L_n be L 's response to K_n in $G_{K,L}^o(X)$ using a winning strategy.

Then *Empty* plays $B(K_n, f_n, \epsilon_n) \cap B(L_n, c_n, 1/3)$, where $c_n = \text{constant}$ n
Suppose $\phi \in \bigcap_{n \in \omega} B(L_n, c_n, 1/3)$. Then $\phi(L_n) \subset (n - 1/3, n + 1/3)$ for all n . Thus $\{L_n : n \in \omega\}$ has a discrete open expansion, contradiction.

Theorem (Ma, GG)

If X is locally compact, then $C_k(X)$ is Baire iff $L \not\preceq G_{K,L}^o(X)$.

Theorem (Ma, GG)

If X is locally compact, then $C_k(X)$ is Baire iff $L \nVdash G_{K,L}^o(X)$.

Question

Is it true that for any completely regular space X , $C_k(X)$ is Baire iff $L \nVdash G_{K,L}^o(X)$? That $NE \uparrow BM(C_k(X))$ iff $K \uparrow G_{K,L}^o(X)$?

A non-game theory characterization of “ L has no winning strategy in $G_{K,L}^o(X)$ ”

A non-game theory characterization of “ L has no winning strategy in $G_{K,L}^o(X)$ ”

Definition. A collection \mathcal{L} of non-empty compact subsets of X is said to *move off* the compact sets if for every compact subset K of X , there is some $L \in \mathcal{L}$ with $K \cap L = \emptyset$.

A non-game theory characterization of “ L has no winning strategy in $G_{K,L}^o(X)$ ”

Definition. A collection \mathcal{L} of non-empty compact subsets of X is said to *move off* the compact sets if for every compact subset K of X , there is some $L \in \mathcal{L}$ with $K \cap L = \emptyset$.

The space X is said to have the *Moving Off Property (MOP)* iff every collection \mathcal{L} which moves off the compact sets contains an infinite subcollection which has a discrete open expansion.

A non-game theory characterization of “ L has no winning strategy in $G_{K,L}^o(X)$ ”

Definition. A collection \mathcal{L} of non-empty compact subsets of X is said to *move off* the compact sets if for every compact subset K of X , there is some $L \in \mathcal{L}$ with $K \cap L = \emptyset$.

The space X is said to have the *Moving Off Property (MOP)* iff every collection \mathcal{L} which moves off the compact sets contains an infinite subcollection which has a discrete open expansion.

Theorem

TFAE:

- 1 X has the MOP;
- 2 $L \not\Uparrow G_{K,L}^o(X)$.

A non-game theory characterization of “ L has no winning strategy in $G_{K,L}^o(X)$ ”

Definition. A collection \mathcal{L} of non-empty compact subsets of X is said to *move off* the compact sets if for every compact subset K of X , there is some $L \in \mathcal{L}$ with $K \cap L = \emptyset$.

The space X is said to have the *Moving Off Property (MOP)* iff every collection \mathcal{L} which moves off the compact sets contains an infinite subcollection which has a discrete open expansion.

Theorem

TFAE:

- 1 X has the MOP;
- 2 $L \nmid G_{K,L}^o(X)$.

Question

Does X have MOP iff $C_k(X)$ Baire?

There is some subtlety in determining when X (even locally compact X) has the property “L has no winning strategy in $G_{K,L}(X)$ ”:

There is some subtlety in determining when X (even locally compact X) has the property “L has no winning strategy in $G_{K,L}(X)$ ”:
Consider spaces $T \cup A$, where T is the Cantor tree and A is a subset of the Cantor set.

There is some subtlety in determining when X (even locally compact X) has the property “ L has no winning strategy in $G_{K,L}(X)$ ”:
Consider spaces $T \cup A$, where T is the Cantor tree and A is a subset of the Cantor set.

Theorem (Ma)

The following are equivalent:

- 1 $C_k(T \cup A)$ is a Baire space;
- 2 $L \nVdash G_{K,L}(X)$
- 3 A is a γ -set.

There is some subtlety in determining when X (even locally compact X) has the property “ L has no winning strategy in $G_{K,L}(X)$ ”:

Consider spaces $T \cup A$, where T is the Cantor tree and A is a subset of the Cantor set.

Theorem (Ma)

The following are equivalent:

- 1 $C_k(T \cup A)$ is a Baire space;
- 2 $L \nVdash G_{K,L}(X)$
- 3 A is a γ -set.

$A \subset \mathbb{R}$ is a γ -set if, given any collection \mathcal{U} of open sets such that any finite subset of A is contained in some member of \mathcal{U} , there are U_0, U_1, \dots in \mathcal{U} such that $A \subset \bigcup_{n \in \omega} \bigcap_{i \geq n} U_i$.

Todorćević showed that it is consistent for there to be two γ -sets A_0 and A_1 whose topological sum is not a γ -set.

Todorćević showed that it is consistent for there to be two γ -sets A_0 and A_1 whose topological sum is not a γ -set.

Since $C_k(X_0 \oplus X_1) \cong C_k(X_0) \times C_k(X_1)$, Ma obtained the following corollary.

Todorćević showed that it is consistent for there to be two γ -sets A_0 and A_1 whose topological sum is not a γ -set.

Since $C_k(X_0 \oplus X_1) \cong C_k(X_0) \times C_k(X_1)$, Ma obtained the following corollary.

Corollary

There are, consistently, two function spaces with the compact-open topology which are Baire but whose product is not.

Todorćević showed that it is consistent for there to be two γ -sets A_0 and A_1 whose topological sum is not a γ -set.

Since $C_k(X_0 \oplus X_1) \cong C_k(X_0) \times C_k(X_1)$, Ma obtained the following corollary.

Corollary

There are, consistently, two function spaces with the compact-open topology which are Baire but whose product is not.

But we don't know about ZFC examples.

Question

Are there examples in ZFC of two Baire function spaces whose product is not Baire?

Some other games and/or applications

Tel'garsky(1975) The countably metacompact game $CM(X)$

Some other games and/or applications

Tel'garsky(1975) The countably metacompact game $CM(X)$

In the n^{th} round, I chooses closed $C_n \subset C_{n-1}$

Some other games and/or applications

Tel'garsky(1975) The countably metacompact game $CM(X)$

In the n^{th} round, I chooses closed $C_n \subset C_{n-1}$

II chooses open $U_n \supset C_n$

Some other games and/or applications

Tel'garsky(1975) The countably metacompact game $CM(X)$

In the n^{th} round, I chooses closed $C_n \subset C_{n-1}$

II chooses open $U_n \supset C_n$

I wins if $\bigcap_{n \in \omega} C_n = \emptyset$ but $\bigcap_{n \in \omega} U_n \neq \emptyset$

Some other games and/or applications

Tel'garsky(1975) The countably metacompact game $CM(X)$

In the n^{th} round, I chooses closed $C_n \subset C_{n-1}$

II chooses open $U_n \supset C_n$

I wins if $\bigcap_{n \in \omega} C_n = \emptyset$ but $\bigcap_{n \in \omega} U_n \neq \emptyset$

X not countably metacompact $\Rightarrow I \uparrow CM(X)$

Some other games and/or applications

Tel'garsky(1975) The countably metacompact game $CM(X)$

In the n^{th} round, I chooses closed $C_n \subset C_{n-1}$

II chooses open $U_n \supset C_n$

I wins if $\bigcap_{n \in \omega} C_n = \emptyset$ but $\bigcap_{n \in \omega} U_n \neq \emptyset$

X not countably metacompact $\Rightarrow I \uparrow CM(X)$

Theorem

$X \times M$ is normal for every metrizable space M iff X is normal and $II \uparrow CM(X)$.

Babinkostova (2005) The screening game $S(X)$
In the n^{th} round, I chooses open cover \mathcal{U}_n of X

Babinkostova (2005) The screening game $S(X)$

In the n^{th} round, I chooses open cover \mathcal{U}_n of X

II chooses disjoint open refinement \mathcal{V}_n of \mathcal{U}_n

Babinkostova (2005) The screening game $S(X)$

In the n^{th} round, I chooses open cover \mathcal{U}_n of X

II chooses disjoint open refinement \mathcal{V}_n of \mathcal{U}_n

II $\uparrow S(X)$ if $\bigcup_{n \in \omega} \mathcal{V}_n$ covers X

Babinkostova (2005) The screening game $S(X)$

In the n^{th} round, I chooses open cover \mathcal{U}_n of X

II chooses disjoint open refinement \mathcal{V}_n of \mathcal{U}_n

II $\uparrow S(X)$ if $\bigcup_{n \in \omega} \mathcal{V}_n$ covers X

Theorem

(Babinkostova)

- 1 II $\uparrow S(X)$ iff X is countable dimensional;
- 2 II has winning strategy in game of length $k + 1$ iff X is $\leq k$ dimensional.

A space X is *selectively separable (SS)* if, given dense sets D_0, D_1, \dots , there are finite $F_j \subset D_j$ with $\bigcup_{n \in \omega} F_n$ dense.

A space X is *selectively separable (SS)* if, given dense sets D_0, D_1, \dots , there are finite $F_j \subset D_j$ with $\bigcup_{n \in \omega} F_n$ dense.

The game **SS(X)** (Dow, Barman): In round n , I chooses dense D_n , II chooses finite $F_n \subset D_n$. II *wins* if $\bigcup_{n \in \omega} F_n$ is dense.

A space X is *selectively separable* (SS) if, given dense sets D_0, D_1, \dots , there are finite $F_j \subset D_j$ with $\bigcup_{n \in \omega} F_n$ dense.

The game $SS(X)$ (Dow, Barman): In round n , I chooses dense D_n , II chooses finite $F_n \subset D_n$. II *wins* if $\bigcup_{n \in \omega} F_n$ is dense.

X is SS^+ if $\text{II} \uparrow SS(X)$.

A space X is *selectively separable* (SS) if, given dense sets D_0, D_1, \dots , there are finite $F_j \subset D_j$ with $\bigcup_{n \in \omega} F_n$ dense.

The game $SS(X)$ (Dow, Barman): In round n , I chooses dense D_n , II chooses finite $F_n \subset D_n$. II *wins* if $\bigcup_{n \in \omega} F_n$ is dense.

X is SS^+ if $II \uparrow SS(X)$.

Countable π -base $\Rightarrow SS^+ \Rightarrow SS$

A space X is *selectively separable* (SS) if, given dense sets D_0, D_1, \dots , there are finite $F_j \subset D_j$ with $\bigcup_{n \in \omega} F_n$ dense.

The game $SS(X)$ (Dow, Barman): In round n , I chooses dense D_n , II chooses finite $F_n \subset D_n$. II *wins* if $\bigcup_{n \in \omega} F_n$ is dense.

X is SS^+ if $II \uparrow SS(X)$.

Countable π -base $\Rightarrow SS^+ \Rightarrow SS$

Separable Fréchet $\Rightarrow SS$

A space X is *selectively separable* (SS) if, given dense sets D_0, D_1, \dots , there are finite $F_j \subset D_j$ with $\bigcup_{n \in \omega} F_n$ dense.

The game $SS(X)$ (Dow, Barman): In round n , I chooses dense D_n , II chooses finite $F_n \subset D_n$. II *wins* if $\bigcup_{n \in \omega} F_n$ is dense.

X is SS^+ if $\text{II} \uparrow SS(X)$.

Countable π -base $\Rightarrow SS^+ \Rightarrow SS$

Separable Fréchet $\Rightarrow SS$

$SS \not\Rightarrow SS^+$.

Theorem

(Dow) X countable $SS^+ \Rightarrow II$ has Markov winning strategy in $SS(X)$.

So, for each dense D , for each $n \in \omega$, one can assign finite $F(D, n) \subset D$ such that, if D_0, D_1, \dots are dense, then $\bigcup_{n \in \omega} F(D_n, n)$ is dense.

Idea of proof.

Let σ be winning strategy for II.

Idea of proof.

Let σ be winning strategy for II.

X countable \Rightarrow II has only countably many possible replies

Idea of proof.

Let σ be winning strategy for II.

X countable \Rightarrow II has only countably many possible replies

For each possible first round reply F , choose $D(F)$ dense such that $\sigma(D(F)) = F$.

Idea of proof.

Let σ be winning strategy for II.

X countable \Rightarrow II has only countably many possible replies

For each possible first round reply F , choose $D(F)$ dense such that $\sigma(D(F)) = F$.

For each possible second round reply F' of II, choose dense $D(F, F')$ such that $\sigma(D(F), D(F, F')) = F'$.

Idea of proof.

Let σ be winning strategy for II.

X countable \Rightarrow II has only countably many possible replies

For each possible first round reply F , choose $D(F)$ dense such that $\sigma(D(F)) = F$.

For each possible second round reply F' of II, choose dense $D(F, F')$ such that $\sigma(D(F), D(F, F')) = F'$.

Etc.

Idea of proof.

Let σ be winning strategy for II.

X countable \Rightarrow II has only countably many possible replies

For each possible first round reply F , choose $D(F)$ dense such that $\sigma(D(F)) = F$.

For each possible second round reply F' of II, choose dense $D(F, F')$ such that $\sigma(D(F), D(F, F')) = F'$.

Etc.

This constructs a (countable) tree of finite sequences of dense sets. Let t_0, t_1, \dots be the nodes of the tree. The Markov winning strategy for II is:

Idea of proof.

Let σ be winning strategy for II.

X countable \Rightarrow II has only countably many possible replies

For each possible first round reply F , choose $D(F)$ dense such that $\sigma(D(F)) = F$.

For each possible second round reply F' of II, choose dense $D(F, F')$ such that $\sigma(D(F), D(F, F')) = F'$.

Etc.

This constructs a (countable) tree of finite sequences of dense sets. Let t_0, t_1, \dots be the nodes of the tree. The Markov winning strategy for II is:

Given dense D in round n , II plays $\sigma(t_n \widehat{\langle D \rangle})$.

Idea of proof.

Let σ be winning strategy for II.

X countable \Rightarrow II has only countably many possible replies

For each possible first round reply F , choose $D(F)$ dense such that $\sigma(D(F)) = F$.

For each possible second round reply F' of II, choose dense $D(F, F')$ such that $\sigma(D(F), D(F, F')) = F'$.

Etc.

This constructs a (countable) tree of finite sequences of dense sets. Let t_0, t_1, \dots be the nodes of the tree. The Markov winning strategy for II is:

Given dense D in round n , II plays $\sigma(t_n \widehat{\langle D \rangle})$.

[Same result for any game with II having only countably many responses, and I's legal plays unchanged during the game.]

SURVEYS

G. Gruenhage, *The story of a topological game*, Rocky Mountain Journal Math. 36 (2006), no. 6, 1885–1914.

M. Scheepers, *Topological games*, in: Encyclopedia of General Topology, K.P. Hart, J. Nagata, and J.E. Vaughan, eds., Elsevier, Amsterdam, 2004, 438-442.

R. Telgarsky, *Topological games: On the 50th anniversary of the Banach-Mazur game*, Rocky Mountain Journal Math. 17(1987), 227-276.

Y. Yajima, *Topological games and applications*, in: Topics in general topology, North-Holland, Amsterdam (1989), 523–562.

OTHER REFERENCES

- L. Babinkostova, *Selective screenability game and covering dimension*, Top. Proc. 29 (2005), 13-17.
- A. Dow and D. Barman, *Selective separability and SS^+* , Topology Proc., to appear.
- F. Galvin and R. Telgarsky, *Stationary strategies in topological games*, Topology Appl. 22(1986), 51-69.
- G. Gruenhage, *Covering properties on $X^2 \setminus \Delta$, W -sets, and compact subsets of Σ -products*, Topology Appl. 17 (1984), no. 3, 287–304.
- G. Gruenhage, *Games, covering properties, and Eberlein compacts*, Topology Appl. 23(1986), 291-297.
- W. Hurewicz, *Über eine Verallgemeinerung des Borelschen Theorems*, Math. Zeitschrift 24(1926), 401-421.
- J. C. Oxtoby, *The Banach-Mazur game and Banach category theorem, Contribution to the theory of games*, Annals of Math. Studies 39 (1957), 157-163.
- R. Telgarsky, *Spaces defined by topological games*, Fund. Math. 88(1975), 193-223.