

Non-continuous satisfaction of identities

W Taylor

BLAST, Boulder, Colorado

June 2, 2010

Four sections of the talk.

Basics

More nearly precise definitions.

An example: $\mu_1(A, \Sigma) = 0$; $\mu_2(A, \Sigma) = \text{diam}(A)$.

Some further results

Compatibility: $A \models \Sigma$

Given a *topological space* A and a set Σ of equations in operation symbols F_t , we write

$$A \models \Sigma,$$

and say that A and Σ are *compatible*, iff there exist *continuous* operations \overline{F}_t on A satisfying Σ .

Compatibility: $A \models \Sigma$

Given a *topological space* A and a set Σ of equations in operation symbols F_t , we write

$$A \models \Sigma,$$

and say that A and Σ are *compatible*, iff there exist *continuous* operations \overline{F}_t on A satisfying Σ .

Examples: Groups on S^1 , S^3 and \mathbb{R} ,

Compatibility: $A \models \Sigma$

Given a *topological space* A and a set Σ of equations in operation symbols F_t , we write

$$A \models \Sigma,$$

and say that A and Σ are *compatible*, iff there exist *continuous* operations \overline{F}_t on A satisfying Σ .

Examples: Groups on S^1 , S^3 and \mathbb{R} , various matrix groups,

Compatibility: $A \models \Sigma$

Given a *topological space* A and a set Σ of equations in operation symbols F_t , we write

$$A \models \Sigma,$$

and say that A and Σ are *compatible*, iff there exist *continuous* operations \overline{F}_t on A satisfying Σ .

Examples: Groups on S^1 , S^3 and \mathbb{R} , various matrix groups, many H-spaces,

Compatibility: $A \models \Sigma$

Given a *topological space* A and a set Σ of equations in operation symbols F_t , we write

$$A \models \Sigma,$$

and say that A and Σ are *compatible*, iff there exist *continuous* operations \overline{F}_t on A satisfying Σ .

Examples: Groups on S^1 , S^3 and \mathbb{R} , various matrix groups, many H-spaces, a lattice on $[0, 1]$,

Compatibility: $A \models \Sigma$

Given a *topological space* A and a set Σ of equations in operation symbols F_t , we write

$$A \models \Sigma,$$

and say that A and Σ are *compatible*, iff there exist *continuous* operations \overline{F}_t on A satisfying Σ .

Examples: Groups on S^1 , S^3 and \mathbb{R} , various matrix groups, many H-spaces, a lattice on $[0, 1]$, a ternary median operation on Y ,

Compatibility: $A \models \Sigma$

Given a *topological space* A and a set Σ of equations in operation symbols F_t , we write

$$A \models \Sigma,$$

and say that A and Σ are *compatible*, iff there exist *continuous* operations \overline{F}_t on A satisfying Σ .

Examples: Groups on S^1 , S^3 and \mathbb{R} , various matrix groups, many H-spaces, a lattice on $[0, 1]$, a ternary median operation on Y , simple Σ on absolute-retract A ,

Compatibility: $A \models \Sigma$

Given a *topological space* A and a set Σ of equations in operation symbols F_t , we write

$$A \models \Sigma,$$

and say that A and Σ are *compatible*, iff there exist *continuous* operations \overline{F}_t on A satisfying Σ .

Examples: Groups on S^1 , S^3 and \mathbb{R} , various matrix groups, many H-spaces, a lattice on $[0, 1]$, a ternary median operation on Y , simple Σ on absolute-retract A , Sets^n on any space A^n ,

Compatibility: $A \models \Sigma$

Given a *topological space* A and a set Σ of equations in operation symbols F_t , we write

$$A \models \Sigma,$$

and say that A and Σ are *compatible*, iff there exist *continuous* operations \overline{F}_t on A satisfying Σ .

Examples: Groups on S^1 , S^3 and \mathbb{R} , various matrix groups, many H-spaces, a lattice on $[0, 1]$, a ternary median operation on Y , simple Σ on absolute-retract A , Sets^n on any space A^n , a unital ring on $S^1 \times \mathbb{Z}$,

Compatibility: $A \models \Sigma$

Given a *topological space* A and a set Σ of equations in operation symbols F_t , we write

$$A \models \Sigma,$$

and say that A and Σ are *compatible*, iff there exist *continuous* operations \overline{F}_t on A satisfying Σ .

Examples: Groups on S^1 , S^3 and \mathbb{R} , various matrix groups, many H-spaces, a lattice on $[0, 1]$, a ternary median operation on Y , simple Σ on absolute-retract A , Sets^n on any space A^n , a unital ring on $S^1 \times \mathbb{Z}$, a Boolean algebra on $\{0, 1\}^{\aleph_0}$.

Compatibility: $A \models \Sigma$

Given a *topological space* A and a set Σ of equations in operation symbols F_t , we write

$$A \models \Sigma,$$

and say that A and Σ are *compatible*, iff there exist *continuous* operations \overline{F}_t on A satisfying Σ .

Examples: Groups on S^1 , S^3 and \mathbb{R} , various matrix groups, many H-spaces, a lattice on $[0, 1]$, a ternary median operation on Y , simple Σ on absolute-retract A , Sets ^{n} on any space A^n , a unital ring on $S^1 \times \mathbb{Z}$, a Boolean algebra on $\{0, 1\}^{\aleph_0}$. Even, for any A and Σ , the Świerczkowski free algebra $\mathbf{F}_A(\Sigma)$ (based on a set larger than A).

$A \models \Sigma$ is a mysterious, sparse relation

- ▶ Experimentally, $A \models \Sigma$ occurs only sporadically, whereas it can in many cases be proved false.

$A \models \Sigma$ is a mysterious, sparse relation

- ▶ Experientially, $A \models \Sigma$ occurs only sporadically, whereas it can in many cases be proved false.
- ▶ But $A \not\models \Sigma$ does not seem to have any uniform method of proof.

$A \models \Sigma$ is a mysterious, sparse relation

- ▶ Experientially, $A \models \Sigma$ occurs only sporadically, whereas it can in many cases be proved false.
- ▶ But $A \not\models \Sigma$ does not seem to have any uniform method of proof.
- ▶ E.g. the sphere $S^n \models \Sigma$ only for trivial Σ or for $n = 1, 3, 7$. (Hard algebraic topology to prove this.)

$A \models \Sigma$ is a mysterious, sparse relation

- ▶ Experimentally, $A \models \Sigma$ occurs only sporadically, whereas it can in many cases be proved false.
- ▶ But $A \not\models \Sigma$ does not seem to have any uniform method of proof.
- ▶ E.g. the sphere $S^n \models \Sigma$ only for trivial Σ or for $n = 1, 3, 7$. (Hard algebraic topology to prove this.)
- ▶ There is no algorithm that settles $\mathbb{R} \models \Sigma$ for finite Σ . (Uses Matiashevich solution of Hilbert's Tenth Problem.) Thus \models is not *too* sparse.

Relaxing the demands of $A \models \Sigma$.

$A \models \Sigma$ demands operations \bar{F}_t that satisfy Σ **exactly** and are **continuous**.

Relaxing the demands of $A \models \Sigma$.

$A \models \Sigma$ demands operations \bar{F}_t that satisfy Σ **exactly** and are **continuous**. We can relax those demands in two ways:

Relaxing the demands of $A \models \Sigma$.

$A \models \Sigma$ demands operations \bar{F}_t that satisfy Σ **exactly** and are **continuous**. We can relax those demands in two ways: we can consider approximate satisfaction, and

Relaxing the demands of $A \models \Sigma$.

$A \models \Sigma$ demands operations \bar{F}_t that satisfy Σ **exactly** and are **continuous**. We can relax those demands in two ways: we can consider approximate satisfaction, and we can consider approximate continuity.

Approximate replacements for $A \models \Sigma$

For (A, d) a metric space, and $\eta > 0$

$$A \models_{\eta} \Sigma,$$

will mean that there exist *continuous* operations \overline{F}_t on A satisfying Σ **within** η .

Approximate replacements for $A \models \Sigma$

For (A, d) a metric space, and $\eta > 0$

$$A \models_{\eta} \Sigma,$$

will mean that there exist *continuous* operations \overline{F}_t on A satisfying Σ **within** η . For $\varepsilon > 0$,

$$A \models^{\varepsilon} \Sigma,$$

will mean that there exist operations \overline{F}_t on A satisfying Σ and **whose discontinuities are no greater than** ε .

Approximate replacements for $A \models \Sigma$

For (A, d) a metric space, and $\eta > 0$

$$A \models_{\eta} \Sigma,$$

will mean that there exist *continuous* operations \overline{F}_t on A satisfying Σ **within** η . For $\varepsilon > 0$,

$$A \models^{\varepsilon} \Sigma,$$

will mean that there exist operations \overline{F}_t on A satisfying Σ and **whose discontinuities are no greater than** ε .

(Of course we also study $A \models_{\eta}^{\varepsilon} \Sigma!$)

The use of $A \models_{\eta} \Sigma$ and $A \models^{\varepsilon} \Sigma$.

- ▶ We said a lot about \models_{η} in “Approximate satisfaction of identities,” and have spoken on it at various times.

The use of $A \models_{\eta} \Sigma$ and $A \models^{\varepsilon} \Sigma$.

- ▶ We said a lot about \models_{η} in “Approximate satisfaction of identities,” and have spoken on it at various times.
- ▶ Today’s talk will be about \models^{ε} .

The use of $A \models_{\eta} \Sigma$ and $A \models^{\varepsilon} \Sigma$.

- ▶ We said a lot about \models_{η} in “Approximate satisfaction of identities,” and have spoken on it at various times.
- ▶ Today’s talk will be about \models^{ε} .
- ▶ Paralleling our previous work, we define

$$\mu(A, \Sigma) = \inf \{ \varepsilon : A \models^{\varepsilon} \Sigma \}.$$

The use of $A \models_{\eta} \Sigma$ and $A \models^{\varepsilon} \Sigma$.

- ▶ We said a lot about \models_{η} in “Approximate satisfaction of identities,” and have spoken on it at various times.
- ▶ Today’s talk will be about \models^{ε} .
- ▶ Paralleling our previous work, we define

$$\mu(A, \Sigma) = \inf \{ \varepsilon : A \models^{\varepsilon} \Sigma \}.$$

- ▶ One hopes that some further understanding of \models will come out of \models^{ε} and μ .

The use of $A \models_{\eta} \Sigma$ and $A \models^{\varepsilon} \Sigma$.

- ▶ We said a lot about \models_{η} in “Approximate satisfaction of identities,” and have spoken on it at various times.
- ▶ Today’s talk will be about \models^{ε} .
- ▶ Paralleling our previous work, we define

$$\mu(A, \Sigma) = \inf \{ \varepsilon : A \models^{\varepsilon} \Sigma \}.$$

- ▶ One hopes that some further understanding of \models will come out of \models^{ε} and μ .
- ▶ E.g. recursive enumerability of $A \models_{\eta} \Sigma$.

Four sections of the talk.

Basics

More nearly precise definitions.

An example: $\mu_1(A, \Sigma) = 0$; $\mu_2(A, \Sigma) = \text{diam}(A)$.

Some further results

(δ, ε) -constraints.

Let (A, d) and (B, e) be metric spaces.

(δ, ε) -constraints.

Let (A, d) and (B, e) be metric spaces.

Given $\bar{F}: B \rightarrow A$ (not necessarily continuous) and $\delta, \varepsilon > 0$, we say that \bar{F} is (δ, ε) -**constrained** if it satisfies: for all $b, b' \in B$, if $e(b, b') < \delta$, then $d(\bar{F}(b), \bar{F}(b')) < \varepsilon$.

(δ, ε) -constraints.

Let (A, d) and (B, e) be metric spaces.

Given $\bar{F}: B \rightarrow A$ (not necessarily continuous) and $\delta, \varepsilon > 0$, we say that \bar{F} is (δ, ε) -**constrained** if it satisfies: for all $b, b' \in B$, if $e(b, b') < \delta$, then $d(\bar{F}(b), \bar{F}(b')) < \varepsilon$.

We say that \bar{F} is n -constrained by (δ_0, δ_n) iff there exist $0 < \delta_0 \leq \delta_1 \leq \dots \leq \delta_n$ such that \bar{F} is (δ_0, δ_1) -constrained and (δ_1, δ_2) -constrained, and so on, up to (δ_{n-1}, δ_n) -constrained.

(δ, ε) -constraints.

Let (A, d) and (B, e) be metric spaces.

Given $\bar{F}: B \rightarrow A$ (not necessarily continuous) and $\delta, \varepsilon > 0$, we say that \bar{F} is (δ, ε) -**constrained** if it satisfies: for all $b, b' \in B$, if $e(b, b') < \delta$, then $d(\bar{F}(b), \bar{F}(b')) < \varepsilon$.

We say that \bar{F} is n -constrained by (δ_0, δ_n) iff there exist $0 < \delta_0 \leq \delta_1 \leq \dots \leq \delta_n$ such that \bar{F} is (δ_0, δ_1) -constrained and (δ_1, δ_2) -constrained, and so on, up to (δ_{n-1}, δ_n) -constrained.

(If \bar{F} is uniformly continuous, then for every $\varepsilon > 0$ there exists $\delta > 0$ so that \bar{F} is n -constrained by (δ, ε) .)

The intermediate-value theorem, revisited.

Lemma

Suppose that f maps a convex subset of \mathbb{R} into \mathbb{R} , and that f is (δ, ε) -constrained with $\delta, \varepsilon > 0$. If $a < c$ and s is between $f(a)$ and $f(c)$, then there exists b with $a \leq b \leq c$ and with $d(f(b), s) < \varepsilon/2$.

Definition of \models_n^ε and $\mu_n(A, \Sigma)$

$$A \models_n^\varepsilon \Sigma$$

Definition of \models_n^ε and $\mu_n(A, \Sigma)$

$$A \models_n^\varepsilon \Sigma$$

means that *there exists an algebra $\mathbf{A} = (A, \overline{F}_t)_{t \in T}$ modeling Σ and a real number $\delta_0 > 0$ such that each \overline{F}_t is n -constrained by (δ_0, ε) .*

Definition of \models_n^ε and $\mu_n(A, \Sigma)$

$$A \models_n^\varepsilon \Sigma$$

means that *there exists an algebra $\mathbf{A} = (A, \overline{F}_t)_{t \in T}$ modeling Σ and a real number $\delta_0 > 0$ such that each \overline{F}_t is n -constrained by (δ_0, ε) .*

We define

$$\mu_n(A, \Sigma) = \inf \{ \varepsilon : A \models_n^\varepsilon \Sigma \}.$$

Definition of \models_n^ε and $\mu_n(A, \Sigma)$

$$A \models_n^\varepsilon \Sigma$$

means that *there exists an algebra $\mathbf{A} = (A, \overline{F}_t)_{t \in T}$ modeling Σ and a real number $\delta_0 > 0$ such that each \overline{F}_t is n -constrained by (δ_0, ε) .*

We define

$$\mu_n(A, \Sigma) = \inf \{ \varepsilon : A \models_n^\varepsilon \Sigma \}.$$

It is not hard to see that

$$0 \leq \mu_1(A, \Sigma) \leq \mu_2(A, \Sigma) \leq \dots \leq \text{diam}(A).$$

Connection of \models_n^ε and $\mu_n(A, \Sigma)$ with $A \models \Sigma$

We repeat the definition:

$$A \models_n^\varepsilon \Sigma$$

means that *there exists an algebra $\mathbf{A} = (A, \overline{F}_t)_{t \in T}$ modeling Σ and a real number $\delta_0 > 0$ such that each \overline{F}_t is n -constrained by (δ_0, ε) .*

Connection of \models_n^ε and $\mu_n(A, \Sigma)$ with $A \models \Sigma$

We repeat the definition:

$$A \models_n^\varepsilon \Sigma$$

means that *there exists an algebra $\mathbf{A} = (A, \overline{F}_t)_{t \in T}$ modeling Σ and a real number $\delta_0 > 0$ such that each \overline{F}_t is n -constrained by (δ_0, ε) .*

Thus if $A \models \Sigma$, then $A \models_n^\varepsilon \Sigma$ for every n and every $\varepsilon > 0$,

Connection of \models_n^ε and $\mu_n(A, \Sigma)$ with $A \models \Sigma$

We repeat the definition:

$$A \models_n^\varepsilon \Sigma$$

means that *there exists an algebra $\mathbf{A} = (A, \overline{F}_t)_{t \in T}$ modeling Σ and a real number $\delta_0 > 0$ such that each \overline{F}_t is n -constrained by (δ_0, ε) .*

Thus if $A \models \Sigma$, then $A \models_n^\varepsilon \Sigma$ for every n and every $\varepsilon > 0$, and hence

$$\mu_n(A, \Sigma) = \inf \{ \varepsilon : A \models_n^\varepsilon \Sigma \} = 0$$

for every n .

Four sections of the talk.

Basics

More nearly precise definitions.

An example: $\mu_1(A, \Sigma) = 0$; $\mu_2(A, \Sigma) = \text{diam}(A)$.

Some further results

“ \overline{G} is a one-one binary operation on A .”

Σ will be this pair of equations:

$$F_0(G(x_0, x_1)) \approx x_0, \quad F_1(G(x_0, x_1)) \approx x_1.$$

“ \overline{G} is a one-one binary operation on A .”

Σ will be this pair of equations:

$$F_0(G(x_0, x_1)) \approx x_0, \quad F_1(G(x_0, x_1)) \approx x_1.$$

A model of Σ has

$$A^2 \xrightarrow{\overline{G}} A \xrightarrow{\overline{F}} A^2 = \text{identity},$$

“ \overline{G} is a one-one binary operation on A .”

Σ will be this pair of equations:

$$F_0(G(x_0, x_1)) \approx x_0, \quad F_1(G(x_0, x_1)) \approx x_1.$$

A model of Σ has

$$A^2 \xrightarrow{\overline{G}} A \xrightarrow{\overline{F}} A^2 = \text{identity},$$

where \overline{F} has \overline{F}_0 and \overline{F}_1 as its component functions.

“ \overline{G} is a one-one binary operation on A .”

Σ will be this pair of equations:

$$F_0(G(x_0, x_1)) \approx x_0, \quad F_1(G(x_0, x_1)) \approx x_1.$$

A model of Σ has

$$A^2 \xrightarrow{\overline{G}} A \xrightarrow{\overline{F}} A^2 = \text{identity},$$

where \overline{F} has \overline{F}_0 and \overline{F}_1 as its component functions. Thus \overline{G} must be one-one and \overline{F} must be onto.

$$\mu_1([0, 1], \Sigma) = 0$$

Repeating Σ : $A^2 \xrightarrow{\bar{G}} A \xrightarrow{\bar{F}} A^2 = \text{identity}$

$$\mu_1([0, 1], \Sigma) = 0$$

Repeating Σ : $A^2 \xrightarrow{\bar{G}} A \xrightarrow{\bar{F}} A^2 = \text{identity}$

So let \bar{F} be a Peano curve: continuous from $[0, 1]$ **onto** $[0, 1]^2$,
and let \bar{G} be any left-inverse to \bar{F} .

$$\mu_1([0, 1], \Sigma) = 0$$

Repeating Σ : $A^2 \xrightarrow{\bar{G}} A \xrightarrow{\bar{F}} A^2 = \text{identity}$

So let \bar{F} be a Peano curve: continuous from $[0, 1]$ **onto** $[0, 1]^2$,
and let \bar{G} be any left-inverse to \bar{F} . \bar{G} is perforce
discontinuous. For arbitrary $\varepsilon > 0$, define

$$G'(a_0, b_0) = \varepsilon \bar{G}(a_0, b_0); \quad F'(a) = \bar{F}(1 \wedge (a/\varepsilon)).$$

$$\mu_1([0, 1], \Sigma) = 0$$

Repeating Σ : $A^2 \xrightarrow{\bar{G}} A \xrightarrow{\bar{F}} A^2 = \text{identity}$

So let \bar{F} be a Peano curve: continuous from $[0, 1]$ **onto** $[0, 1]^2$, and let \bar{G} be any left-inverse to \bar{F} . \bar{G} is perforce discontinuous. For arbitrary $\varepsilon > 0$, define

$$G'(a_0, b_0) = \varepsilon \bar{G}(a_0, b_0); \quad F'(a) = \bar{F}(1 \wedge (a/\varepsilon)).$$

Now the discontinuities of G' are no larger than ε , and F' remains continuous, while F' and G' still satisfy Σ . Thus $A \models_1^\varepsilon$ for every $\varepsilon > 0$; hence $\mu_1([0, 1], \Sigma) = 0$. ■

$$\mu_2([0, 1], \Sigma) = 1$$

We consider $(A; \overline{G}, \overline{F}_0, \overline{F}_1)$ modeling Σ , with the operations (δ_0, δ_1) -constrained and (δ_1, δ_2) -constrained. We will show that $\delta_2 \geq 1$.

$$\mu_2([0, 1], \Sigma) = 1$$

We consider $(A; \overline{G}, \overline{F}_0, \overline{F}_1)$ modeling Σ , with the operations (δ_0, δ_1) -constrained and (δ_1, δ_2) -constrained. We will show that $\delta_2 \geq 1$.

Let $\{a_0, a_1\} = \{b_0, b_1\} = \{0, 1\}$. Compare the four values $\overline{G}(a_i, b_j)$; w.l.o.g. $\overline{G}(a_0, b_0)$ is the smallest.

$$\mu_2([0, 1], \Sigma) = 1$$

We consider $(A; \overline{G}, \overline{F}_0, \overline{F}_1)$ modeling Σ , with the operations (δ_0, δ_1) -constrained and (δ_1, δ_2) -constrained. We will show that $\delta_2 \geq 1$.

Let $\{a_0, a_1\} = \{b_0, b_1\} = \{0, 1\}$. Compare the four values $\overline{G}(a_i, b_j)$; w.l.o.g. $\overline{G}(a_0, b_0)$ is the smallest. Again w.l.o.g. we have $\overline{G}(a_1, b_0) \leq \overline{G}(a_0, b_1)$. In other words,

$$\overline{G}(a_0, b_0) \leq \overline{G}(a_1, b_0) \leq \overline{G}(a_0, b_1).$$

$$\mu_2([0, 1], \Sigma) = 1$$

We consider $(A; \overline{G}, \overline{F}_0, \overline{F}_1)$ modeling Σ , with the operations (δ_0, δ_1) -constrained and (δ_1, δ_2) -constrained. We will show that $\delta_2 \geq 1$.

Let $\{a_0, a_1\} = \{b_0, b_1\} = \{0, 1\}$. Compare the four values $\overline{G}(a_i, b_j)$; w.l.o.g. $\overline{G}(a_0, b_0)$ is the smallest. Again w.l.o.g. we have $\overline{G}(a_1, b_0) \leq \overline{G}(a_0, b_1)$. In other words,

$$\overline{G}(a_0, b_0) \leq \overline{G}(a_1, b_0) \leq \overline{G}(a_0, b_1).$$

We consider the real function $\overline{H}(x) = \overline{G}(a_0, x)$.

$$\mu_2([0, 1], \Sigma) = 1$$

We consider $(A; \overline{G}, \overline{F}_0, \overline{F}_1)$ modeling Σ , with the operations (δ_0, δ_1) -constrained and (δ_1, δ_2) -constrained. We will show that $\delta_2 \geq 1$.

Let $\{a_0, a_1\} = \{b_0, b_1\} = \{0, 1\}$. Compare the four values $\overline{G}(a_i, b_j)$; w.l.o.g. $\overline{G}(a_0, b_0)$ is the smallest. Again w.l.o.g. we have $\overline{G}(a_1, b_0) \leq \overline{G}(a_0, b_1)$. In other words,

$$\overline{G}(a_0, b_0) \leq \overline{G}(a_1, b_0) \leq \overline{G}(a_0, b_1).$$

We consider the real function $\overline{H}(x) = \overline{G}(a_0, x)$. By the Lemma (IVT), there exists $e \in [0, 1]$ with

$$d(\overline{G}(a_0, e), \overline{G}(a_1, b_0)) < \delta_1.$$

$\mu_2([0, 1], \Sigma) = 1$, concluded

Repeat: $d(\overline{G}(a_0, e), \overline{G}(a_1, b_0)) < \delta_1$.

$\mu_2([0, 1], \Sigma) = 1$, concluded

Repeat: $d(\overline{G}(a_0, e), \overline{G}(a_1, b_0)) < \delta_1$.

Because $\{a_0, a_1\} = \{0, 1\}$, because of Σ , and because the function \overline{F}_0 is (δ_1, δ_2) -constrained, we now have:

$$1 = d(a_0, a_1) = d(\overline{F}_0(\overline{G}(a_0, e)), \overline{F}_0(\overline{G}(a_1, b_0))) \leq \delta_2.$$

$\mu_2([0, 1], \Sigma) = 1$, concluded

Repeat: $d(\overline{G}(a_0, e), \overline{G}(a_1, b_0)) < \delta_1$.

Because $\{a_0, a_1\} = \{0, 1\}$, because of Σ , and because the function \overline{F}_0 is (δ_1, δ_2) -constrained, we now have:

$$1 = d(a_0, a_1) = d(\overline{F}_0(\overline{G}(a_0, e)), \overline{F}_0(\overline{G}(a_1, b_0))) \leq \delta_2.$$

Thus $\mu_2([0, 1], \Sigma)$, being the infimum of such δ_2 's, must be 1.



Four sections of the talk.

Basics

More nearly precise definitions.

An example: $\mu_1(A, \Sigma) = 0$; $\mu_2(A, \Sigma) = \text{diam}(A)$.

Some further results

Some further results . . .

- ▶ $\mu_2([0, 1]^2, \text{same } \Sigma) = 1$ — uses Borsuk-Ulam Theorem.

Some further results . . .

- ▶ $\mu_2([0, 1]^2, \text{same } \Sigma) = 1$ — uses Borsuk-Ulam Theorem.
- ▶ $\mu_2(\mathbb{R}, \text{Groups} + x^n \approx 1) \geq \text{radius}(\mathbb{R})$.

Some further results . . .

- ▶ $\mu_2([0, 1]^2, \text{same } \Sigma) = 1$ — uses Borsuk-Ulam Theorem.
- ▶ $\mu_2(\mathbb{R}, \text{Groups} + x^n \approx 1) \geq \text{radius}(\mathbb{R})$.
- ▶ $\mu_2([0, 1]^n, \text{Groups}) = \text{diameter}([0, 1]^n)$.

Some further results . . .

- ▶ $\mu_2([0, 1]^2, \text{same } \Sigma) = 1$ — uses Borsuk-Ulam Theorem.
- ▶ $\mu_2(\mathbb{R}, \text{Groups} + x^n \approx 1) \geq \text{radius}(\mathbb{R})$.
- ▶ $\mu_2([0, 1]^n, \text{Groups}) = \text{diameter}([0, 1]^n)$.
- ▶ $\mu_3(Y, \text{Lattices}) \geq 0.5$.

(Here Y stands for a Y-shaped one-dimensional space with each arm of unit length.)

... and two more

Here Σ stands for the equations defining a binary operation with left-zero and left-one, or defining a commutative idempotent binary operation, or defining a ternary majority operation.

... and two more

Here Σ stands for the equations defining a binary operation with left-zero and left-one, or defining a commutative idempotent binary operation, or defining a ternary majority operation.

- ▶ $\mu_1(S^1, \Sigma) = 2/3$.

... and two more

Here Σ stands for the equations defining a binary operation with left-zero and left-one, or defining a commutative idempotent binary operation, or defining a ternary majority operation.

- ▶ $\mu_1(S^1, \Sigma) = 2/3.$

- ▶ $\mu_1(S^2, \Sigma) \geq 2/3.$