

# Lattice valued identities

B. Šešelja and A. Tepavčević

**Blast 2010**

Boulder, June 6., 2009.

### Lattice valued function

### Lattice valued function

$X \neq \emptyset$ ;  $L$  - complete lattice with the top 1 and the bottom 0;

### Lattice valued function

$X \neq \emptyset$ ;  $L$  - complete lattice with the top 1 and the bottom 0;

$$\mu : X \rightarrow L$$

### Lattice valued function

$X \neq \emptyset$ ;  $L$  - complete lattice with the top 1 and the bottom 0;

$$\mu : X \rightarrow L$$

$\mu$  is a **lattice-valued function** or  **$L$ -valued function** on  $X$ .

### Lattice valued function

$X \neq \emptyset$ ;  $L$  - complete lattice with the top 1 and the bottom 0;

$$\mu : X \rightarrow L$$

$\mu$  is a **lattice-valued function** or  **$L$ -valued function** on  $X$ .

$\mu : X \rightarrow L$  - an  $L$ -valued function on  $X$ ;  $p \in L$ .

### Lattice valued function

$X \neq \emptyset$ ;  $L$  - complete lattice with the top 1 and the bottom 0;

$$\mu : X \rightarrow L$$

$\mu$  is a **lattice-valued function** or  **$L$ -valued function** on  $X$ .

$\mu : X \rightarrow L$  - an  $L$ -valued function on  $X$ ;  $p \in L$ .

$$\mu_p := \{x \in X \mid \mu(x) \geq p\}$$

### Lattice valued function

$X \neq \emptyset$ ;  $L$  - complete lattice with the top 1 and the bottom 0;

$$\mu : X \rightarrow L$$

$\mu$  is a **lattice-valued function** or  **$L$ -valued function** on  $X$ .

$\mu : X \rightarrow L$  - an  $L$ -valued function on  $X$ ;  $p \in L$ .

$$\mu_p := \{x \in X \mid \mu(x) \geq p\}$$

$\mu_p$  is the  **$p$ -cut**, a **cut set** or simply a **cut** of  $\mu$ .

## Lattice valued function

$X \neq \emptyset$ ;  $L$  - complete lattice with the top 1 and the bottom 0;

$$\mu : X \rightarrow L$$

$\mu$  is a **lattice-valued function** or  **$L$ -valued function** on  $X$ .

$\mu : X \rightarrow L$  - an  $L$ -valued function on  $X$ ;  $p \in L$ .

$$\mu_p := \{x \in X \mid \mu(x) \geq p\}$$

$\mu_p$  is the  **$p$ -cut**, a **cut set** or simply a **cut** of  $\mu$ .

*A  $p$ -cut of  $\mu$  is the inverse image of the principal filter in  $L$  generated by  $p$ :*

$$\mu_p = \mu^{-1}(\uparrow p).$$

# Lattice valued functions

## Properties of cuts

### Proposition

*For an  $L$ -valued function  $\mu$  on  $X$ , the following hold.*

# Lattice valued functions

## Properties of cuts

### Proposition

*For an  $L$ -valued function  $\mu$  on  $X$ , the following hold.*

- *If  $p, q \in L$  and  $p \leq q$ , then  $\mu_q \subseteq \mu_p$ .*

# Lattice valued functions

## Properties of cuts

### Proposition

For an  $L$ -valued function  $\mu$  on  $X$ , the following hold.

- If  $p, q \in L$  and  $p \leq q$ , then  $\mu_q \subseteq \mu_p$ .
- if  $L_1 \subseteq L$ , then

$$\bigcap \{ \mu_p \mid p \in L_1 \} = \mu_{\bigvee \{ p \mid p \in L_1 \}}.$$

### Proposition

For an  $L$ -valued function  $\mu$  on  $X$ , the following hold.

- If  $p, q \in L$  and  $p \leq q$ , then  $\mu_q \subseteq \mu_p$ .
- if  $L_1 \subseteq L$ , then

$$\bigcap \{\mu_p \mid p \in L_1\} = \mu_{\bigvee \{p \mid p \in L_1\}}.$$

- $\bigcup \{\mu_p \mid p \in L\} = X$ .

### Proposition

For an  $L$ -valued function  $\mu$  on  $X$ , the following hold.

- If  $p, q \in L$  and  $p \leq q$ , then  $\mu_q \subseteq \mu_p$ .
- if  $L_1 \subseteq L$ , then

$$\bigcap \{\mu_p \mid p \in L_1\} = \mu_{\bigvee \{p \mid p \in L_1\}}.$$

- $\bigcup \{\mu_p \mid p \in L\} = X$ .

$\mu_L := \{\mu_p \mid p \in L\}$  - collection of cuts of  $\mu$ .

# Lattice valued functions

## Properties of cuts

### Proposition

For an  $L$ -valued function  $\mu$  on  $X$ , the following hold.

- If  $p, q \in L$  and  $p \leq q$ , then  $\mu_q \subseteq \mu_p$ .
- if  $L_1 \subseteq L$ , then

$$\bigcap \{ \mu_p \mid p \in L_1 \} = \mu_{\bigvee \{ p \mid p \in L_1 \}}.$$

- $\bigcup \{ \mu_p \mid p \in L \} = X$ .

$\mu_L := \{ \mu_p \mid p \in L \}$  - collection of cuts of  $\mu$ .

### Proposition

For every  $x \in X$ ,

$$\bigcap \{ \mu_p \mid x \in \mu_p \} \in \mu_L.$$

# Lattice valued functions

## Properties of cuts

### Theorem

*The collection  $\mu_L$  of cuts of  $\mu : X \rightarrow L$  is a complete lattice under inclusion.*

# Lattice valued functions

## Properties of cuts

### Theorem

*The collection  $\mu_L$  of cuts of  $\mu : X \rightarrow L$  is a complete lattice under inclusion.*

## Synthesis by cuts

# Lattice valued functions

## Properties of cuts

### Theorem

*The collection  $\mu_L$  of cuts of  $\mu : X \rightarrow L$  is a complete lattice under inclusion.*

## Synthesis by cuts

### Theorem

*If  $\mu : X \rightarrow L$  is an  $L$ -valued function on  $X$ , then for every  $x \in X$*

$$\mu(x) = \bigvee \{p \in L \mid x \in \mu_p\}.$$

### Representation theorem

### Representation theorem

#### Theorem

*Let  $X$  be a nonempty set and  $F$  a family of its subsets closed under arbitrary intersections and containing  $X$  (a closure system on  $X$ ). Let also  $L$  be the lattice dual to  $(F, \subseteq)$  and  $\mu : X \rightarrow L$  an  $L$ -valued function on  $X$  defined by*

$$\mu(x) := \bigcap \{f \in F \mid x \in f\}.$$

*Then, the lattice of cut subsets of  $\mu$  is isomorphic with  $(F, \subseteq)$ , and every  $f \in F$  coincides with the corresponding cut  $\mu_f$ .*

# Lattice valued functions

Cardinal power

$$L^X := \{\mu \mid \mu : X \rightarrow L\}$$

# Lattice valued functions

Cardinal power

$$L^X := \{\mu \mid \mu : X \rightarrow L\}$$

$L^X$  is the collection of all  $L$ -valued functions on  $X$ .

# Lattice valued functions

Cardinal power

$$L^X := \{\mu \mid \mu : X \rightarrow L\}$$

$L^X$  is the collection of all  $L$ -valued functions on  $X$ .

$L^X$  is a lattice under the ordering defined by:

$$\mu \leq \nu \text{ if and only if for each } x \in X \mu(x) \leq \nu(x).$$

# Lattice valued functions

Cardinal power

$$L^X := \{\mu \mid \mu : X \rightarrow L\}$$

$L^X$  is the collection of all  $L$ -valued functions on  $X$ .

$L^X$  is a lattice under the ordering defined by:

$$\mu \leq \nu \text{ if and only if for each } x \in X \text{ } \mu(x) \leq \nu(x).$$

For  $\mu \in L^X$ , let

# Lattice valued functions

Cardinal power

$$L^X := \{\mu \mid \mu : X \rightarrow L\}$$

$L^X$  is the collection of all  $L$ -valued functions on  $X$ .

$L^X$  is a lattice under the ordering defined by:

$$\mu \leq \nu \text{ if and only if for each } x \in X \text{ } \mu(x) \leq \nu(x).$$

For  $\mu \in L^X$ , let

$$L_\mu := (\{\uparrow p \cap \mu(X) \mid p \in L\}, \subseteq).$$

# Lattice valued functions

## Cardinal power

$$L^X := \{\mu \mid \mu : X \rightarrow L\}$$

$L^X$  is the collection of all  $L$ -valued functions on  $X$ .

$L^X$  is a lattice under the ordering defined by:

$$\mu \leq \nu \text{ if and only if for each } x \in X \text{ } \mu(x) \leq \nu(x).$$

For  $\mu \in L^X$ , let

$$L_\mu := (\{\uparrow p \cap \mu(X) \mid p \in L\}, \subseteq).$$

By the definition,  $L_\mu$  consists of particular collections of images of  $\mu$  in  $L$  and is a poset under inclusion.

# Lattice valued functions

## Classification

$$L_\mu := (\{\uparrow p \cap \mu(X) \mid p \in L\}, \subseteq).$$

# Lattice valued functions

## Classification

$$L_\mu := (\{\uparrow p \cap \mu(X) \mid p \in L\}, \subseteq).$$

$\mu_L = \{\mu_p \mid p \in L\}$  - the lattice of cuts of  $\mu$ .

# Lattice valued functions

## Classification

$$L_\mu := (\{\uparrow p \cap \mu(X) \mid p \in L\}, \subseteq).$$

$\mu_L = \{\mu_p \mid p \in L\}$  - the lattice of cuts of  $\mu$ .

### Theorem

*$L_\mu$  is a lattice isomorphic with the lattice  $\mu_L$  of cuts of  $\mu$ , under*

$$f : \mu_p \mapsto \uparrow p \cap \mu(X).$$

# Lattice valued functions

## Classification

A classification of functions in the set  $L^X$  is introduced as follows.

# Lattice valued functions

## Classification

A classification of functions in the set  $L^X$  is introduced as follows.

*Let  $\sim$  be the relation on  $L^X$ , defined by:*

# Lattice valued functions

## Classification

A classification of functions in the set  $L^X$  is introduced as follows.

*Let  $\sim$  be the relation on  $L^X$ , defined by:*

*$\mu \sim \nu$  if and only if the correspondence  $f : \mu(x) \mapsto \nu(x)$ ,  $x \in X$  is a bijection from  $\mu(X)$  onto  $\nu(X)$  which has an extension to an isomorphism from the lattice  $L_\mu$  onto the lattice  $L_\nu$ , given by the mapping*

# Lattice valued functions

## Classification

A classification of functions in the set  $L^X$  is introduced as follows.

Let  $\sim$  be the relation on  $L^X$ , defined by:

$\mu \sim \nu$  if and only if the correspondence  $f : \mu(x) \mapsto \nu(x)$ ,  $x \in X$  is a bijection from  $\mu(X)$  onto  $\nu(X)$  which has an extension to an isomorphism from the lattice  $L_\mu$  onto the lattice  $L_\nu$ , given by the mapping

$$F(\uparrow p \cap \mu(X)) := \uparrow \bigwedge \{ \nu(x) \mid \mu(x) \geq p \} \cap \nu(X), p \in L.$$

# Lattice valued functions

## Classification

A classification of functions in the set  $L^X$  is introduced as follows.

*Let  $\sim$  be the relation on  $L^X$ , defined by:*

*$\mu \sim \nu$  if and only if the correspondence  $f : \mu(x) \mapsto \nu(x)$ ,  $x \in X$  is a bijection from  $\mu(X)$  onto  $\nu(X)$  which has an extension to an isomorphism from the lattice  $L_\mu$  onto the lattice  $L_\nu$ , given by the mapping*

$$F(\uparrow p \cap \mu(X)) := \uparrow \bigwedge \{ \nu(x) \mid \mu(x) \geq p \} \cap \nu(X), p \in L.$$

If  $\mu \sim \nu$ , then the  $L$ -valued functions  $\mu$  and  $\nu$  on  $X$  are said to be **equivalent**.

### Classification of functions in $L^X$

### Classification of functions in $L^X$

#### Theorem

*Let  $\mu, \nu : X \rightarrow L$ . Then  $\mu \sim \nu$  if and only if  $L$ -valued functions  $\mu$  and  $\nu$  have equal collections of cuts.*

# Lattice valued functions

Cuts applied

## Example

## Example

$$\mu = \begin{pmatrix} x & y & z \\ p & q & r \end{pmatrix} \quad \nu = \begin{pmatrix} x & y & z \\ p & q & t \end{pmatrix} \quad \pi = \begin{pmatrix} x & y & z \\ p & r & t \end{pmatrix}$$

# Lattice valued functions


Cuts applied

## Example

$$\mu = \begin{pmatrix} x & y & z \\ p & q & r \end{pmatrix}$$

$$\nu = \begin{pmatrix} x & y & z \\ p & q & t \end{pmatrix}$$




$$\pi = \begin{pmatrix} x & y & z \\ p & r & t \end{pmatrix}$$



$L$


# Lattice valued functions

Cuts applied



# Lattice valued functions


Cuts applied



$$L_\mu = (\{\uparrow p \cap \mu(X) \mid p \in L\}, \subseteq)$$

# Lattice valued functions

Cuts applied



$$\mu_p = \mu^{-1}(\uparrow p); \quad \mu_L = \{\mu_p \mid p \in L\}$$

## Representation of lattices

## Representation of lattices

### Theorem

*Let  $L$  be a lattice of finite length, and  $X$  the set of its meet irreducible elements. Then there is an  $L$ -valued function  $\mu : X \rightarrow L$  such that  $L$  is isomorphic with the dual of the lattice  $\mu_L$  of cuts of  $\mu$ , under  $p \mapsto \mu_p$ .*

# Lattice valued relations

Definition

## Lattice valued relations

# Lattice valued relations

## Definition

## Lattice valued relations

A **lattice valued relation**  $R$  on a set  $X$  is a lattice valued function on  $X^2$ :

# Lattice valued relations

## Definition

### Lattice valued relations

A **lattice valued relation**  $R$  on a set  $X$  is a lattice valued function on  $X^2$ :

$$R : X^2 \rightarrow L.$$

# Lattice valued relations

## Definition

## Lattice valued relations

A **lattice valued relation**  $R$  on a set  $X$  is a lattice valued function on  $X^2$ :

$$R : X^2 \rightarrow L.$$

A **cut-relation** of  $R$  is a  $p$ -cut of  $R$ ,  $p \in L$ :

# Lattice valued relations

## Definition

### Lattice valued relations

A **lattice valued relation**  $R$  on a set  $X$  is a lattice valued function on  $X^2$ :

$$R : X^2 \rightarrow L.$$

A **cut-relation** of  $R$  is a  $p$ -cut of  $R$ ,  $p \in L$ :

$$R_p = \{(x, y) \in X^2 \mid R(x, y) \geq p\} = R^{-1}(\uparrow p).$$

# Lattice valued relations

## Definition

## Lattice valued relations

A **lattice valued relation**  $R$  on a set  $X$  is a lattice valued function on  $X^2$ :

$$R : X^2 \rightarrow L.$$

A **cut-relation** of  $R$  is a  $p$ -cut of  $R$ ,  $p \in L$ :

$$R_p = \{(x, y) \in X^2 \mid R(x, y) \geq p\} = R^{-1}(\uparrow p).$$

An  $L$ -valued relation  $R$  on  $X$  is

## Lattice valued relations

A **lattice valued relation**  $R$  on a set  $X$  is a lattice valued function on  $X^2$ :

$$R : X^2 \rightarrow L.$$

A **cut-relation** of  $R$  is a  $p$ -cut of  $R$ ,  $p \in L$ :

$$R_p = \{(x, y) \in X^2 \mid R(x, y) \geq p\} = R^{-1}(\uparrow p).$$

An  $L$ -valued relation  $R$  on  $X$  is

**reflexive** and **symmetric** if it fulfills the analogue properties for its characteristic function; it is **transitive** if

$R(x, y) \wedge R(y, z) \leq R(x, z)$ , for all  $x, y, z \in X$ .

## Lattice valued relations

A **lattice valued relation**  $R$  on a set  $X$  is a lattice valued function on  $X^2$ :

$$R : X^2 \rightarrow L.$$

A **cut-relation** of  $R$  is a  $p$ -cut of  $R$ ,  $p \in L$ :

$$R_p = \{(x, y) \in X^2 \mid R(x, y) \geq p\} = R^{-1}(\uparrow p).$$

An  $L$ -valued relation  $R$  on  $X$  is

**reflexive** and **symmetric** if it fulfills the analogue properties for its characteristic function; it is **transitive** if

$$R(x, y) \wedge R(y, z) \leq R(x, z), \text{ for all } x, y, z \in X.$$

An  $L$ -valued relation  $R$  on  $X$  is a **lattice valued equivalence relation** on  $X$  if it is reflexive, symmetric and transitive.

### Lattice valued (sub)algebra

### Lattice valued (sub)algebra

Let  $\mathcal{A} = (A, F)$  be an algebra and  $L$  a complete lattice.

### Lattice valued (sub)algebra

Let  $\mathcal{A} = (A, F)$  be an algebra and  $L$  a complete lattice.

A **lattice valued** or  **$L$ -valued subalgebra** of  $\mathcal{A}$ , is any mapping  $\mu : A \rightarrow L$  fulfilling the following:

### Lattice valued (sub)algebra

Let  $\mathcal{A} = (A, F)$  be an algebra and  $L$  a complete lattice.

A **lattice valued** or  **$L$ -valued subalgebra** of  $\mathcal{A}$ , is any mapping

$\mu : A \rightarrow L$  fulfilling the following:

For any operation  $f$  from  $F$ ,  $f : A^n \rightarrow A$ ,  $n \in \mathbb{N}$ , and all

$x_1, \dots, x_n \in A$ ,

### Lattice valued (sub)algebra

Let  $\mathcal{A} = (A, F)$  be an algebra and  $L$  a complete lattice.

A **lattice valued** or  **$L$ -valued subalgebra** of  $\mathcal{A}$ , is any mapping

$\mu : A \rightarrow L$  fulfilling the following:

For any operation  $f$  from  $F$ ,  $f : A^n \rightarrow A$ ,  $n \in \mathbb{N}$ , and all

$x_1, \dots, x_n \in A$ ,

$$\bigwedge_{i=1}^n \mu(x_i) \leq \mu(f(x_1, \dots, x_n)).$$

### Lattice valued (sub)algebra

Let  $\mathcal{A} = (A, F)$  be an algebra and  $L$  a complete lattice.

A **lattice valued** or  **$L$ -valued subalgebra** of  $\mathcal{A}$ , is any mapping  $\mu : A \rightarrow L$  fulfilling the following:

For any operation  $f$  from  $F$ ,  $f : A^n \rightarrow A$ ,  $n \in \mathbb{N}$ , and all  $x_1, \dots, x_n \in A$ ,

$$\bigwedge_{i=1}^n \mu(x_i) \leq \mu(f(x_1, \dots, x_n)).$$

For a nullary operation (constant)  $c \in F$ ,

### Lattice valued (sub)algebra

Let  $\mathcal{A} = (A, F)$  be an algebra and  $L$  a complete lattice.

A **lattice valued** or  **$L$ -valued subalgebra** of  $\mathcal{A}$ , is any mapping  $\mu : A \rightarrow L$  fulfilling the following:

For any operation  $f$  from  $F$ ,  $f : A^n \rightarrow A$ ,  $n \in \mathbb{N}$ , and all  $x_1, \dots, x_n \in A$ ,

$$\bigwedge_{i=1}^n \mu(x_i) \leq \mu(f(x_1, \dots, x_n)).$$

For a nullary operation (constant)  $c \in F$ ,

$$\mu(c) = 1,$$

where 1 is the top element in  $L$ .

## Example

### Example

An  $L$ -valued subgroup of a group  $(G, \cdot, {}^{-1}, e)$  is a mapping  $\mu : G \rightarrow L$ , fulfilling the following:

### Example

An  $L$ -valued subgroup of a group  $(G, \cdot, {}^{-1}, e)$  is a mapping  $\mu : G \rightarrow L$ , fulfilling the following:

- $\mu(x \cdot y) \geq \mu(x) \wedge \mu(y)$ , for all  $x, y \in G$ .
- $\mu(e) = 1$ .
- $\mu(x^{-1}) \geq \mu(x)$ , for every  $x \in G$ .

### Example

An  $L$ -valued subgroup of a group  $(G, \cdot, {}^{-1}, e)$  is a mapping  $\mu : G \rightarrow L$ , fulfilling the following:

- $\mu(x \cdot y) \geq \mu(x) \wedge \mu(y)$ , for all  $x, y \in G$ .
- $\mu(e) = 1$ .
- $\mu(x^{-1}) \geq \mu(x)$ , for every  $x \in G$ .

### Theorem

*If  $\mu : A \rightarrow L$  is a lattice valued subalgebra of an algebra  $\mathcal{A}$ , then for every  $p \in L$ , the cut set  $\mu_p$  is a subalgebra of  $\mathcal{A}$ .*

### Theorem

Let  $\mathcal{A}$  be an algebra and  $\mathcal{F}$  a collection of its subuniverses closed under arbitrary intersections and containing  $A$ . Let also  $L$  be the lattice dual to  $(\mathcal{F}, \subseteq)$  and  $\mu : A \rightarrow L$  an  $L$ -valued set on  $A$  defined by

$$\mu(x) := \bigcap \{B \in \mathcal{F} \mid x \in B\}.$$

Then,  $\mu$  is an  $L$ -valued subalgebra of  $\mathcal{A}$ . In addition, the lattice of cut subalgebras of  $\mu$  is isomorphic with  $(\mathcal{F}, \subseteq)$ , and every subalgebra  $B \in \mathcal{F}$  coincides with the corresponding cut  $\mu_B$ .

### Lattice valued congruences

### Lattice valued congruences

Let  $\mathcal{A} = (A, F)$  be an algebra and  $L$  a complete lattice, and  $R : A^2 \rightarrow L$  be an  $L$ -valued relation on  $A$ .

### Lattice valued congruences

Let  $\mathcal{A} = (A, F)$  be an algebra and  $L$  a complete lattice, and  $R : A^2 \rightarrow L$  be an  $L$ -valued relation on  $A$ .

$R$  is said to be **compatible** with operations on  $\mathcal{A}$  if for any ( $n$ -ary)  $f \in F$  and all  $x_1, \dots, x_n, y_1, \dots, y_n \in A$ , we have that

$$\bigwedge_{i=1}^n R(x_i, y_i) \leq R(f(x_1, \dots, x_n), f(y_1, \dots, y_n)).$$

### Lattice valued congruences

Let  $\mathcal{A} = (A, F)$  be an algebra and  $L$  a complete lattice, and  $R : A^2 \rightarrow L$  be an  $L$ -valued relation on  $A$ .

$R$  is said to be **compatible** with operations on  $\mathcal{A}$  if for any ( $n$ -ary)  $f \in F$  and all  $x_1, \dots, x_n, y_1, \dots, y_n \in A$ , we have that

$$\bigwedge_{i=1}^n R(x_i, y_i) \leq R(f(x_1, \dots, x_n), f(y_1, \dots, y_n)).$$

An  $L$ -valued equivalence relation on  $\mathcal{A}$  which is compatible with all operations is a **lattice valued congruence** relation on  $\mathcal{A}$ .

### Lattice valued congruences

Let  $\mathcal{A} = (A, F)$  be an algebra and  $L$  a complete lattice, and  $R : A^2 \rightarrow L$  be an  $L$ -valued relation on  $A$ .

$R$  is said to be **compatible** with operations on  $\mathcal{A}$  if for any ( $n$ -ary)  $f \in F$  and all  $x_1, \dots, x_n, y_1, \dots, y_n \in A$ , we have that

$$\bigwedge_{i=1}^n R(x_i, y_i) \leq R(f(x_1, \dots, x_n), f(y_1, \dots, y_n)).$$

An  $L$ -valued equivalence relation on  $\mathcal{A}$  which is compatible with all operations is a **lattice valued congruence** relation on  $\mathcal{A}$ .

#### Theorem

*If  $R : A^2 \rightarrow L$  is a lattice valued congruence on an algebra  $\mathcal{A}$ , then for every  $p \in L$ , the cut relation  $R_p$  is a congruence on  $\mathcal{A}$ .*

# Lattice valued algebras

## Relations on $L$ -valued sets

Let  $A$  be a nonempty set,  $L$  a complete lattice and  $\mu : A \rightarrow L$  an  $L$ -valued set on  $A$ . An  $L$ -valued relation  $\rho : A^2 \rightarrow L$  on  $A$  is said to be an  **$L$ -valued relation on  $\mu$**  if for all  $x, y \in A$

$$\rho(x, y) \leq \mu(x) \wedge \mu(y). \quad (1)$$

# Lattice valued algebras

## Relations on $L$ -valued sets

Let  $A$  be a nonempty set,  $L$  a complete lattice and  $\mu : A \rightarrow L$  an  $L$ -valued set on  $A$ . An  $L$ -valued relation  $\rho : A^2 \rightarrow L$  on  $A$  is said to be an  **$L$ -valued relation on  $\mu$**  if for all  $x, y \in A$

$$\rho(x, y) \leq \mu(x) \wedge \mu(y). \quad (1)$$

Due to this boundary condition, we have the following definition.

# Lattice valued algebras

## Relations on $L$ -valued sets

Let  $A$  be a nonempty set,  $L$  a complete lattice and  $\mu : A \rightarrow L$  an  $L$ -valued set on  $A$ . An  $L$ -valued relation  $\rho : A^2 \rightarrow L$  on  $A$  is said to be an  **$L$ -valued relation on  $\mu$**  if for all  $x, y \in A$

$$\rho(x, y) \leq \mu(x) \wedge \mu(y). \quad (1)$$

Due to this boundary condition, we have the following definition. An  $L$ -valued relation  $\rho$  on an  $L$ -valued set  $\mu$  is **reflexive** if for all  $x, y \in A$ ,

$$\rho(x, x) = \mu(x). \quad (2)$$

# Lattice valued algebras

## Relations on $L$ -valued sets

Let  $A$  be a nonempty set,  $L$  a complete lattice and  $\mu : A \rightarrow L$  an  $L$ -valued set on  $A$ . An  $L$ -valued relation  $\rho : A^2 \rightarrow L$  on  $A$  is said to be an  **$L$ -valued relation on  $\mu$**  if for all  $x, y \in A$

$$\rho(x, y) \leq \mu(x) \wedge \mu(y). \quad (1)$$

Due to this boundary condition, we have the following definition. An  $L$ -valued relation  $\rho$  on an  $L$ -valued set  $\mu$  is **reflexive** if for all  $x, y \in A$ ,

$$\rho(x, x) = \mu(x). \quad (2)$$

Obviously, by (1), a reflexive relation  $\rho$  on  $\mu$  fulfils the following:

$$\text{For all } x, y \in A, \rho(x, x) \geq \rho(x, y) \text{ and } \rho(x, x) \geq \rho(y, x). \quad (3)$$

# Lattice valued algebras

## Relations on $L$ -valued sets

An  $L$ -valued relation  $\rho$  on  $\mu$  is symmetric and transitive if it fulfils the corresponding properties as an  $L$ -valued relation on  $A$ .

# Lattice valued algebras

## Relations on $L$ -valued sets

An  $L$ -valued relation  $\rho$  on  $\mu$  is symmetric and transitive if it fulfils the corresponding properties as an  $L$ -valued relation on  $A$ .

A reflexive, symmetric and transitive relation  $\rho$  on  $\mu$  is an  **$L$ -valued equivalence** on this  $L$ -valued set.

# Lattice valued algebras

## Relations on $L$ -valued sets

An  $L$ -valued relation  $\rho$  on  $\mu$  is symmetric and transitive if it fulfils the corresponding properties as an  $L$ -valued relation on  $A$ .

A reflexive, symmetric and transitive relation  $\rho$  on  $\mu$  is an  **$L$ -valued equivalence** on this  $L$ -valued set.

An  $L$ -valued equivalence relation  $\rho$  on  $\mu$ , fulfilling

$$\text{For all } x, y \in A, \rho(x, x) > \rho(x, y) \text{ and } \rho(x, x) > \rho(y, x), \quad (4)$$

is called an  **$L$ -valued equality** relation on an  $L$ -valued set  $\mu$ .

# Lattice valued algebras

## Relations on $L$ -valued sets

An  $L$ -valued relation  $\rho$  on  $\mu$  is symmetric and transitive if it fulfils the corresponding properties as an  $L$ -valued relation on  $A$ .

A reflexive, symmetric and transitive relation  $\rho$  on  $\mu$  is an  **$L$ -valued equivalence** on this  $L$ -valued set.

An  $L$ -valued equivalence relation  $\rho$  on  $\mu$ , fulfilling

$$\text{For all } x, y \in A, \rho(x, x) > \rho(x, y) \text{ and } \rho(x, x) > \rho(y, x), \quad (4)$$

is called an  **$L$ -valued equality** relation on an  $L$ -valued set  $\mu$ .

In addition, an  $L$ -valued relation  $\rho$  on  $\mu$  is **compatible** with the operations on this  $L$ -valued subalgebra if for any ( $n$ -ary)  $f \in F$  and all  $x_1, \dots, x_n, y_1, \dots, y_n \in A$ , we have that

$$\bigwedge_{i=1}^n R(x_i, y_i) \leq R(f(x_1, \dots, x_n), f(y_1, \dots, y_n)).$$

# Lattice valued algebras

## Relations on $L$ -valued sets

A compatible  $L$ -valued equivalence on  $\mu$  is an  **$L$ -valued congruence** on this  $L$ -valued subalgebra.

# Lattice valued algebras

## Relations on $L$ -valued sets

A compatible  $L$ -valued equivalence on  $\mu$  is an  **$L$ -valued congruence** on this  $L$ -valued subalgebra.

A compatible lattice valued equality is also called an  **$L$ -valued equality** on  $\mu$ .

# Lattice valued algebras

## Relations on $L$ -valued sets

A compatible  $L$ -valued equivalence on  $\mu$  is an  **$L$ -valued congruence** on this  $L$ -valued subalgebra.

A compatible lattice valued equality is also called an  **$L$ -valued equality** on  $\mu$ .

Denote by  $LCon \mu$  and  $LEq \mu$  the collections of all  $L$ -valued congruences and all compatible  $L$ -valued equalities (respectively) on an  $L$ -valued subalgebra  $\mu$  of an algebra  $\mathcal{A}$ . These can be naturally ordered by componentwise order  $\leq$ , inherited from  $L$ .

# Lattice valued algebras

## Relations on $L$ -valued sets

A compatible  $L$ -valued equivalence on  $\mu$  is an  **$L$ -valued congruence** on this  $L$ -valued subalgebra.

A compatible lattice valued equality is also called an  **$L$ -valued equality** on  $\mu$ .

Denote by  $L\text{Con } \mu$  and  $L\text{Eq } \mu$  the collections of all  $L$ -valued congruences and all compatible  $L$ -valued equalities (respectively) on an  $L$ -valued subalgebra  $\mu$  of an algebra  $\mathcal{A}$ . These can be naturally ordered by componentwise order  $\leq$ , inherited from  $L$ .

### Theorem

*The poset  $(L\text{Con } \mu, \leq)$  is a complete lattice, and the poset  $(L\text{Eq } \mu, \leq)$  is a meet-semilattice, a semi-ideal in the former.*

# Lattice valued algebras

Connection to the underlying algebra

We have an algebra  $\mathcal{A} = (A, F)$  and  $L$ -valued relations on it, i.e., mappings from  $A^2$  to a complete lattice  $L$ .

# Lattice valued algebras

## Connection to the underlying algebra

We have an algebra  $\mathcal{A} = (A, F)$  and  $L$ -valued relations on it, i.e., mappings from  $A^2$  to a complete lattice  $L$ .

We say that  $\rho : A^2 \rightarrow L$  is an  **$L$ -valued weakly reflexive** relation on  $\mathcal{A}$ , if

$$\rho(c, c) = 1 \text{ for every constant } c \in F. \quad (5)$$

# Lattice valued algebras

## Connection to the underlying algebra

We have an algebra  $\mathcal{A} = (A, F)$  and  $L$ -valued relations on it, i.e., mappings from  $A^2$  to a complete lattice  $L$ .

We say that  $\rho : A^2 \rightarrow L$  is an  **$L$ -valued weakly reflexive** relation on  $\mathcal{A}$ , if

$$\rho(c, c) = 1 \text{ for every constant } c \in F. \quad (5)$$

An  $L$ -valued relation  $\rho : A^2 \rightarrow L$  on an algebra  $\mathcal{A}$  which is weakly reflexive, symmetric and transitive, is called a **weak  $L$ -valued equivalence** on  $\mathcal{A}$ .

# Lattice valued algebras

## Connection to the underlying algebra

We have an algebra  $\mathcal{A} = (A, F)$  and  $L$ -valued relations on it, i.e., mappings from  $A^2$  to a complete lattice  $L$ .

We say that  $\rho : A^2 \rightarrow L$  is an  **$L$ -valued weakly reflexive** relation on  $\mathcal{A}$ , if

$$\rho(c, c) = 1 \text{ for every constant } c \in F. \quad (5)$$

An  $L$ -valued relation  $\rho : A^2 \rightarrow L$  on an algebra  $\mathcal{A}$  which is weakly reflexive, symmetric and transitive, is called a **weak  $L$ -valued equivalence** on  $\mathcal{A}$ .

If, in addition,  $\rho$  fulfills also the condition

$$\text{For all } x, y \in A, \rho(x, x) > \rho(x, y) \text{ and } \rho(x, x) > \rho(y, x),$$

then  $\rho$  is a **weak  $L$ -valued equality** on  $\mathcal{A}$ .

# Lattice valued algebras

Connection to the underlying algebra

For compatible weak  $L$ -valued equivalences we use the name **weak  $L$ -valued congruences** on  $\mathcal{A}$ . A subclass of weak  $L$ -valued congruences are **compatible weak  $L$ -valued equalities**.

# Lattice valued algebras

## Connection to the underlying algebra

For compatible weak  $L$ -valued equivalences we use the name **weak  $L$ -valued congruences** on  $\mathcal{A}$ . A subclass of weak  $L$ -valued congruences are **compatible weak  $L$ -valued equalities**.

### Theorem

If  $\rho : A^2 \rightarrow L$  is a weak  $L$ -valued congruence on an algebra  $\mathcal{A}$ , then the mapping  $\mu_\rho : A \rightarrow L$ , defined by

$$\mu_\rho(x) := \rho(x, x) \quad (6)$$

is an  $L$ -valued subalgebra of  $\mathcal{A}$ .

# Lattice valued algebras

Connection to the underlying algebra

The previous theorem gives a link between  $L$ -valued congruences on  $L$ -valued subalgebras and weak  $L$ -valued congruences on the whole algebra.

# Lattice valued algebras

## Connection to the underlying algebra

The previous theorem gives a link between  $L$ -valued congruences on  $L$ -valued subalgebras and weak  $L$ -valued congruences on the whole algebra.

### Theorem

*A weak  $L$ -valued congruence  $\rho : A^2 \rightarrow L$  on an algebra  $\mathcal{A}$  is an  $L$ -valued congruence on the  $L$ -valued subalgebra  $\mu_\rho$  of  $\mathcal{A}$ .  
Conversely, an  $L$ -valued congruence  $\rho$  on an  $L$ -valued subalgebra  $\mu$  of  $\mathcal{A}$  is a weak  $L$ -valued congruence on the whole algebra  $\mathcal{A}$ .*

# Lattice valued algebras

Lattice of all weak  $L$ -valued congruences

## Theorem

*The collection of all weak  $L$ -valued congruences on an algebra  $\mathcal{A}$  is a complete lattice. Its sublattice of diagonal relations is isomorphic to the lattice of all  $L$ -valued subalgebras of  $\mathcal{A}$ . The lattice of  $L$ -valued congruences on each  $L$ -valued subalgebra of  $\mathcal{A}$  is an interval sublattice.*

# Lattice valued algebras

## Identities

Let  $\mathcal{A} = (A, F)$  be an algebra and  $L$  a complete lattice.

# Lattice valued algebras

## Identities

Let  $\mathcal{A} = (A, F)$  be an algebra and  $L$  a complete lattice.

Let  $\mu$  be an  $L$ -valued subalgebra of  $\mathcal{A}$  and  $E : A^2 \rightarrow L$  an  $L$ -valued equality on  $\mu$ .

# Lattice valued algebras

## Identities

Let  $\mathcal{A} = (A, F)$  be an algebra and  $L$  a complete lattice.

Let  $\mu$  be an  $L$ -valued subalgebra of  $\mathcal{A}$  and  $E : A^2 \rightarrow L$  an  $L$ -valued equality on  $\mu$ .

If  $t_1, t_2$  are terms in the language of  $\mathcal{A}$ , we consider the expression  $E(t_1, t_2)$  as an  **$L$ -valued identity with respect to  $E$** , or (briefly)  **$L$ -valued identity**, if  $E$  is fixed.

# Lattice valued algebras

## Identities

Let  $\mathcal{A} = (A, F)$  be an algebra and  $L$  a complete lattice.

Let  $\mu$  be an  $L$ -valued subalgebra of  $\mathcal{A}$  and  $E : A^2 \rightarrow L$  an  $L$ -valued equality on  $\mu$ .

If  $t_1, t_2$  are terms in the language of  $\mathcal{A}$ , we consider the expression  $E(t_1, t_2)$  as an  **$L$ -valued identity with respect to  $E$** , or (briefly)  **$L$ -valued identity**, if  $E$  is fixed.

Suppose that  $x_1, \dots, x_n$  are variables appearing in terms  $t_1, t_2$ . We say that an  $L$ -valued subalgebra  $\mu$  of  $\mathcal{A}$  **satisfies** the  $L$ -valued identity  $E(t_1, t_2)$  (or that this  $L$ -valued identity is valid on  $L$ -valued subalgebra  $\mu$ ) if for all  $x_1, \dots, x_n \in A$

$$\bigwedge_{i=1}^n \mu(x_i) \leq E(t_1, t_2). \quad (7)$$

### Proposition

*Let  $\mu : A \rightarrow L$  be an  $L$ -valued subalgebra of an algebra  $\mathcal{A}$  and  $E : A^2 \rightarrow L$  an  $L$ -valued equality on  $\mu$ . If  $\mu$  satisfies an  $L$ -valued identity  $E(f, g)$ , then also  $\mu$  satisfies the identity  $E_1(f, g)$ , for every  $L$ -valued equality  $E_1$  on  $\mu$ , such that  $E \leq E_1$ .*

### Proposition

Let  $\mu : A \rightarrow L$  be an  $L$ -valued subalgebra of an algebra  $\mathcal{A}$  and  $E : A^2 \rightarrow L$  an  $L$ -valued equality on  $\mu$ . If  $\mu$  satisfies an  $L$ -valued identity  $E(f, g)$ , then also  $\mu$  satisfies the identity  $E_1(f, g)$ , for every  $L$ -valued equality  $E_1$  on  $\mu$ , such that  $E \leq E_1$ .

### Lemma

Let  $\mu : A \rightarrow L$  be an  $L$ -valued subalgebra of an algebra  $\mathcal{A}$ ,  $\{E_i : A^2 \rightarrow L, i \in I\}$  a family of  $L$ -valued equalities on  $\mu$ , and  $f, g$  terms in the language of  $\mathcal{A}$ . Now, if  $\mu$  satisfies the identity  $E_i(f, g)$  for every  $i \in I$ , then  $\mu$  also satisfies the identity  $E(f, g)$ , where  $E = \bigwedge_{i \in I} E_i$ .

### Corollary

*If an  $L$ -valued subalgebra  $\mu$  of  $\mathcal{A}$  satisfies the identity  $E(f, g)$  for an  $L$ -valued equality  $E$ , then there is the least  $L$ -valued equality on  $\mu$ , denoted by  $E_{\mu(f,g)}$ , such that  $\mu$  satisfies  $E_{\mu(f,g)}(f, g)$ .*

### Corollary

*If an  $L$ -valued subalgebra  $\mu$  of  $\mathcal{A}$  satisfies the identity  $E(f, g)$  for an  $L$ -valued equality  $E$ , then there is the least  $L$ -valued equality on  $\mu$ , denoted by  $E_{\mu(f,g)}$ , such that  $\mu$  satisfies  $E_{\mu(f,g)}(f, g)$ .*

### Corollary

*Let  $\mu$  be an  $L$ -valued subalgebra of an algebra  $\mathcal{A}$ . Then  $\mathcal{A}$  satisfies the identity  $f = g$  if and only if  $\mu$  satisfies the identity  $E(f, g)$  for every  $E \in \text{fEq } \mu$ .*

### Corollary

*Let  $\mu$  be an  $L$ -valued subalgebra of an algebra  $\mathcal{A}$  and  $L$  a complete lattice. Let also  $f, g$  be terms in the language of  $\mathcal{A}$ . Then the following hold.*

*The cut subalgebra  $\mu_p$  of  $\mu$  for  $p \in L$  satisfies the identity  $f = g$  if and only if the cut-relation  $(E_{\mu(f,g)})_p$  of the least equality  $E_{\mu(f,g)}$  is the ordinary equality on  $\mu_p$ .*

## References

## References

- A. Di Nola, G. Gerla, *Lattice valued algebras*, Stochastica 11 (1987) 137150.

## References

- A. Di Nola, G. Gerla, *Lattice valued algebras*, Stochastica 11 (1987) 137-150.
- M. Demirci, *Vague Groups*, J. Math. Anal. Appl. 230,(1999) 142-156.

## References

- A. Di Nola, G. Gerla, *Lattice valued algebras*, Stochastica 11 (1987) 137-150.
- M. Demirci, *Vague Groups*, J. Math. Anal. Appl. 230,(1999) 142-156.
- R. Bělohlávek, *Fuzzy Relational Systems: Foundations and Principles*, Kluwer Academic/Plenum Publishers, New York, 2002.

- B. Šešelja, A. Tepavčević, *Equivalent fuzzy sets*, Kybernetika 41 (2005), No.2, 115-128.

# Lattice valued identities

## References

- B. Šešelja, A. Tepavčević, *Equivalent fuzzy sets*, Kybernetika 41 (2005), No.2, 115-128.
- B. Šešelja, *Lattice-valued Covering Relation and Ordering: An Abstract Approach*, Computational Intelligence, Theory and Applications, Bernd Reusch (Ed.), Springer, 2006, 295-300.

- B. Šešelja, A. Tepavčević, *Equivalent fuzzy sets*, Kybernetika 41 (2005), No.2, 115-128.
- B. Šešelja, *Lattice-valued Covering Relation and Ordering: An Abstract Approach*, Computational Intelligence, Theory and Applications, Bernd Reusch (Ed.), Springer, 2006, 295-300.
- V. Janis, B. Šešelja, A. Tepavčević, *Non-standard cut classification of lattice-valued sets*, Information Sciences 177 (2007) 161-169.

- B. Šešelja, A. Tepavčević, *Equivalent fuzzy sets*, Kybernetika 41 (2005), No.2, 115-128.
- B. Šešelja, *Lattice-valued Covering Relation and Ordering: An Abstract Approach*, Computational Intelligence, Theory and Applications, Bernd Reusch (Ed.), Springer, 2006, 295-300.
- V. Janis, B. Šešelja, A. Tepavčević, *Non-standard cut classification of lattice-valued sets*, Information Sciences 177 (2007) 161-169.
- B. Borchardt, A. Maletti, B. Šešelja, A. Tepavčević, H. Vogler, *Cut sets as recognizable tree languages*, Fuzzy Sets and Systems 157 (2006) 1560-1571.

**Thank you for your attention!**