

Lifters, free
sets, ladders

The Conc
functor

k -ladders

Critical points

From lifting
objects to
lifting
diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating
Kuratowski
indexes of
finite posets

Large free sets versus k -ladders: combinatorial issues raised by “From lifting objects to lifting diagrams”

Friedrich Wehrung

Université de Caen

LMNO, UMR 6139

Département de Mathématiques

14032 Caen cedex

E-mail: wehrung@math.unicaen.fr

URL: <http://www.math.unicaen.fr/~wehrung>

Most of the results discussed here obtained with Pierre Gillibert.

June 2-6, 2010

An example with the Con_c functor

- Denote by $\text{Con}_c A$ the $(\vee, 0)$ -semilattice of all **compact** (i.e., **finitely generated**) congruences of an (universal) algebra A .

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

An example with the Con_c functor

- Denote by $\text{Con}_c A$ the $(\vee, 0)$ -semilattice of all **compact** (i.e., **finitely generated**) congruences of an (universal) algebra A .
- A $(\vee, 0)$ -semilattice S is **distributive** if for all $a, b, c \in S$, if $c \leq a \vee b$ then there are $x \leq a$ and $y \leq b$ such that $c = x \vee y$.

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

An example with the Con_c functor

- Denote by $\text{Con}_c A$ the $(\vee, 0)$ -semilattice of all **compact** (i.e., **finitely generated**) congruences of an (universal) algebra A .
- A $(\vee, 0)$ -semilattice S is **distributive** if for all $a, b, c \in S$, if $c \leq a \vee b$ then there are $x \leq a$ and $y \leq b$ such that $c = x \vee y$.
- Equivalently, $\text{Id } S$ is a distributive lattice.

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

An example with the Con_c functor

- Denote by $\text{Con}_c A$ the $(\vee, 0)$ -semilattice of all **compact** (i.e., **finitely generated**) congruences of an (universal) algebra A .
- A $(\vee, 0)$ -semilattice S is **distributive** if for all $a, b, c \in S$, if $c \leq a \vee b$ then there are $x \leq a$ and $y \leq b$ such that $c = x \vee y$.
- Equivalently, $\text{Id } S$ is a distributive lattice.
- In particular, every **Boolean** semilattice is distributive.

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

An example with the Con_c functor

- Denote by $\text{Con}_c A$ the $(\vee, 0)$ -semilattice of all **compact** (i.e., **finitely generated**) congruences of an (universal) algebra A .
- A $(\vee, 0)$ -semilattice S is **distributive** if for all $a, b, c \in S$, if $c \leq a \vee b$ then there are $x \leq a$ and $y \leq b$ such that $c = x \vee y$.
- Equivalently, $\text{Id } S$ is a distributive lattice.
- In particular, every **Boolean** semilattice is distributive.
- Let \mathcal{V} be a **variety** of algebras.

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

An example with the Con_c functor

- Denote by $\text{Con}_c A$ the $(\vee, 0)$ -semilattice of all **compact** (i.e., **finitely generated**) congruences of an (universal) algebra A .
- A $(\vee, 0)$ -semilattice S is **distributive** if for all $a, b, c \in S$, if $c \leq a \vee b$ then there are $x \leq a$ and $y \leq b$ such that $c = x \vee y$.
- Equivalently, $\text{Id } S$ is a distributive lattice.
- In particular, every **Boolean** semilattice is distributive.
- Let \mathcal{V} be a **variety** of algebras.
- Assume that every **finite Boolean** semilattice S is isomorphic to $\text{Con}_c A$ for some $A \in \mathcal{V}$ (we say that S can be **lifted** in \mathcal{V} , or that it is **representable in \mathcal{V}**)

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

An example with the Con_c functor

- Denote by $\text{Con}_c A$ the $(\vee, 0)$ -semilattice of all **compact** (i.e., **finitely generated**) congruences of an (universal) algebra A .
- A $(\vee, 0)$ -semilattice S is **distributive** if for all $a, b, c \in S$, if $c \leq a \vee b$ then there are $x \leq a$ and $y \leq b$ such that $c = x \vee y$.
- Equivalently, $\text{Id } S$ is a distributive lattice.
- In particular, every **Boolean** semilattice is distributive.
- Let \mathcal{V} be a **variety** of algebras.
- Assume that every **finite Boolean** semilattice S is isomorphic to $\text{Con}_c A$ for some $A \in \mathcal{V}$ (we say that S can be **lifted** in \mathcal{V} , or that it is **representable in \mathcal{V}**) (**more assumptions coming**).

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

An example with the Con_c functor

- Denote by $\text{Con}_c A$ the $(\vee, 0)$ -semilattice of all **compact** (i.e., **finitely generated**) congruences of an (universal) algebra A .
- A $(\vee, 0)$ -semilattice S is **distributive** if for all $a, b, c \in S$, if $c \leq a \vee b$ then there are $x \leq a$ and $y \leq b$ such that $c = x \vee y$.
- Equivalently, $\text{Id } S$ is a distributive lattice.
- In particular, every **Boolean** semilattice is distributive.
- Let \mathcal{V} be a **variety** of algebras.
- Assume that every **finite Boolean** semilattice S is isomorphic to $\text{Con}_c A$ for some $A \in \mathcal{V}$ (we say that S can be **lifted** in \mathcal{V} , or that it is **representable in \mathcal{V}**) (**more assumptions coming**).
- We would like to prove that every **countable** distributive $(\vee, 0)$ -semilattice can be lifted in \mathcal{V} .

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Con_c example, continued (1)

- Without additional assumptions, this can't be done (e.g., let \mathcal{V} be the variety of all distributive lattices).

Lifters, free sets, ladders

The Con_c functor

k-ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, \langle \lambda \rangle) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Con_c example, continued (1)

- Without additional assumptions, this can't be done (e.g., let \mathcal{V} be the variety of all distributive lattices).
- Now suppose that for each $A \in \mathcal{V}$, each finite Boolean semilattice S , every $(\vee, 0)$ -homomorphism $\varphi: \text{Con}_c A \rightarrow S$ is (up to iso) of the form $\text{Con}_c f: \text{Con}_c A \rightarrow \text{Con}_c B$, for some $B \in \mathcal{V}$ and some homomorphism $f: A \rightarrow B$.

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, <\lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Con_c example, continued (1)

- Without additional assumptions, this can't be done (e.g., let \mathcal{V} be the variety of all distributive lattices).
- Now suppose that for each $A \in \mathcal{V}$, each finite Boolean semilattice S , every $(\vee, 0)$ -homomorphism $\varphi: \text{Con}_c A \rightarrow S$ is (up to iso) of the form $\text{Con}_c f: \text{Con}_c A \rightarrow \text{Con}_c B$, for some $B \in \mathcal{V}$ and some homomorphism $f: A \rightarrow B$.
Important observation: Con_c is a (quite nice) functor.

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, <\lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Con_c example, continued (1)

- Without additional assumptions, this can't be done (e.g., let \mathcal{V} be the variety of all distributive lattices).
- Now suppose that for each $A \in \mathcal{V}$, each finite Boolean semilattice S , every $(\vee, 0)$ -homomorphism $\varphi: \text{Con}_c A \rightarrow S$ is (up to iso) of the form $\text{Con}_c f: \text{Con}_c A \rightarrow \text{Con}_c B$, for some $B \in \mathcal{V}$ and some homomorphism $f: A \rightarrow B$.
Important observation: Con_c is a (quite nice) functor.
(one-dimensional amalgamation property for Con_c).

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, <\lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Con_c example, continued (1)

- Without additional assumptions, this can't be done (e.g., let \mathcal{V} be the variety of all distributive lattices).
- Now suppose that for each $A \in \mathcal{V}$, each finite Boolean semilattice S , every $(\vee, 0)$ -homomorphism $\varphi: \text{Con}_c A \rightarrow S$ is (up to iso) of the form $\text{Con}_c f: \text{Con}_c A \rightarrow \text{Con}_c B$, for some $B \in \mathcal{V}$ and some homomorphism $f: A \rightarrow B$.
Important observation: Con_c is a (quite nice) functor. (one-dimensional amalgamation property for Con_c).
- Now write $S = \varinjlim_{n < \omega} S_n$, with all S_n Boolean (Bulman-Fleming and McDowell, 1978).

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Con_c example, continued (1)

- Without additional assumptions, this can't be done (e.g., let \mathcal{V} be the variety of all distributive lattices).
- Now suppose that for each $A \in \mathcal{V}$, each finite Boolean semilattice S , every $(\vee, 0)$ -homomorphism $\varphi: \text{Con}_c A \rightarrow S$ is (up to iso) of the form $\text{Con}_c f: \text{Con}_c A \rightarrow \text{Con}_c B$, for some $B \in \mathcal{V}$ and some homomorphism $f: A \rightarrow B$.
Important observation: Con_c is a (quite nice) functor. (one-dimensional amalgamation property for Con_c).
- Now write $S = \varinjlim_{n < \omega} S_n$, with all S_n Boolean (Bulman-Fleming and McDowell, 1978).
- Suppose $S_n \cong \text{Con}_c A_n$. By the assumption above, represent the transition map $S_n \rightarrow S_{n+1}$ as $\text{Con}_c f_n$, for some $f_n: A_n \rightarrow A_{n+1}$.

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Con_c example, continued (1)

Lifters, free sets, ladders

The Con_c functor

k-ladders

Critical points

From lifting objects to lifting diagrams

λ-lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

- Without additional assumptions, this can't be done (e.g., let \mathcal{V} be the variety of all distributive lattices).
- Now suppose that for each $A \in \mathcal{V}$, each finite Boolean semilattice S , every $(\vee, 0)$ -homomorphism $\varphi: \text{Con}_c A \rightarrow S$ is (up to iso) of the form $\text{Con}_c f: \text{Con}_c A \rightarrow \text{Con}_c B$, for some $B \in \mathcal{V}$ and some homomorphism $f: A \rightarrow B$.
Important observation: Con_c is a (quite nice) functor. (one-dimensional amalgamation property for Con_c).
- Now write $S = \varinjlim_{n < \omega} S_n$, with all S_n Boolean (Bulman-Fleming and McDowell, 1978).
- Suppose $S_n \cong \text{Con}_c A_n$. By the assumption above, represent the transition map $S_n \rightarrow S_{n+1}$ as $\text{Con}_c f_n$, for some $f_n: A_n \rightarrow A_{n+1}$.
- Let $A = \varinjlim_{n < \omega} A_n$. Then $\text{Con}_c A \cong S$.

Con_c example, continued (2)

Lifters, free
sets, ladders

The Con_c
functor

k-ladders

Critical points

From lifting
objects to
lifting
diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating
Kuratowski
indexes of
finite posets

- How to extend this to larger cardinalities?

Con_c example, continued (2)

Lifters, free sets, ladders

The Con_c functor

k-ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

- How to extend this to larger cardinalities?
- Example for \aleph_1 instructive.

Con_c example, continued (2)

Lifters, free sets, ladders

The Con_c functor

k-ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

- How to extend this to larger cardinalities?
- Example for \aleph_1 instructive.
- 1-dimensional amalgamation no longer sufficient.

Con_c example, continued (2)

Lifters, free sets, ladders

The Con_c functor

k-ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

- How to extend this to larger cardinalities?
- Example for \aleph_1 instructive.
- 1-dimensional amalgamation no longer sufficient.
- **2-dimensional amalgamation property** necessary.

Con_c example, continued (2)

Lifters, free sets, ladders

The Con_c functor

k-ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

- How to extend this to larger cardinalities?
- Example for \aleph_1 instructive.
- 1-dimensional amalgamation no longer sufficient.
- 2-dimensional amalgamation property necessary.
- ... Remaining question: what plays the role of $\omega = \{0, 1, 2, \dots\}$ (index poset of a direct system)?

k -ladders

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

- A poset P is **lower finite** if $P \downarrow p := \{x \in P \mid x \leq p\}$ is finite $\forall p \in P$.

k -ladders

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, <\lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

- A poset P is **lower finite** if $P \downarrow p := \{x \in P \mid x \leq p\}$ is finite $\forall p \in P$.
- For a positive integer k , a **k -ladder** is a lower finite lattice where every element has $\leq k$ lower covers.

k -ladders

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, <\lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

- A poset P is **lower finite** if $P \downarrow p := \{x \in P \mid x \leq p\}$ is finite $\forall p \in P$.
- For a positive integer k , a **k -ladder** is a lower finite lattice where every element has $\leq k$ lower covers.

Theorem (Ditor 1984)

k -ladders

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

- A poset P is **lower finite** if $P \downarrow p := \{x \in P \mid x \leq p\}$ is finite $\forall p \in P$.
- For a positive integer k , a **k -ladder** is a lower finite lattice where every element has $\leq k$ lower covers.

Theorem (Ditor 1984)

- (i) Every k -ladder has cardinality $\leq \aleph_{k-1}$.

k -ladders

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

- A poset P is **lower finite** if $P \downarrow p := \{x \in P \mid x \leq p\}$ is finite $\forall p \in P$.
- For a positive integer k , a **k -ladder** is a lower finite lattice where every element has $\leq k$ lower covers.

Theorem (Ditor 1984)

- (i) Every k -ladder has cardinality $\leq \aleph_{k-1}$.
- (ii) There exists a 2-ladder of cardinality \aleph_1 .

k -ladders

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

- A poset P is **lower finite** if $P \downarrow p := \{x \in P \mid x \leq p\}$ is finite $\forall p \in P$.
- For a positive integer k , a **k -ladder** is a lower finite lattice where every element has $\leq k$ lower covers.

Theorem (Ditor 1984)

- Every k -ladder has cardinality $\leq \aleph_{k-1}$.
- There exists a 2-ladder of cardinality \aleph_1 .

Observe that the 1-ladders are exactly the finite chains together with $\omega = \{0, 1, 2, \dots\}$.

Construction of a 2-ladder of cardinality \aleph_1

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Idea of proof of (ii): $F = \bigcup_{\alpha < \omega_1} F_\alpha$. Start with $F_0 := \{0\}$.
Suppose F_α constructed.

Construction of a 2-ladder of cardinality \aleph_1

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points


From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Idea of proof of (ii): $F = \bigcup_{\alpha < \omega_1} F_\alpha$. Start with $F_0 := \{0\}$. Suppose F_α constructed.



Construction of a 2-ladder of cardinality \aleph_1

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points


From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Idea of proof of (ii): $F = \bigcup_{\alpha < \omega_1} F_\alpha$. Start with $F_0 := \{0\}$. Suppose F_α constructed.



Construction of a 2-ladder of cardinality \aleph_1

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points


From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Idea of proof of (ii): $F = \bigcup_{\alpha < \omega_1} F_\alpha$. Start with $F_0 := \{0\}$. Suppose F_α constructed.



Construction of a 2-ladder of cardinality \aleph_1

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points


From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Idea of proof of (ii): $F = \bigcup_{\alpha < \omega_1} F_\alpha$. Start with $F_0 := \{0\}$. Suppose F_α constructed.



A few applications of 2-ladders

Lifters, free
sets, ladders

The Conc
functor

k -ladders

Critical points

From lifting
objects to
lifting
diagrams

λ -lifters

$(\kappa, <\lambda) \rightsquigarrow P$

Calculating
Kuratowski
indexes of
finite posets

Theorem

A few applications of 2-ladders

Lifters, free
sets, ladders

The Conc
functor

k -ladders

Critical points

From lifting
objects to
lifting
diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating
Kuratowski
indexes of
finite posets

Theorem

Every distributive $(\vee, 0)$ -semilattice of cardinality \aleph_1 is isomorphic to

A few applications of 2-ladders

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Theorem

Every distributive $(\vee, 0)$ -semilattice of cardinality \aleph_1 is isomorphic to

- $\text{Con}_c L$ for some lattice L (Huhn 1989),

A few applications of 2-ladders

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Theorem

Every distributive $(\vee, 0)$ -semilattice of cardinality \aleph_1 is isomorphic to

- $\text{Con}_c L$ for some lattice L (Huhn 1989),
- the normal subgroup lattice of some group, or the submodule lattice of some module (Růžička, Tůma, and W., 2007).

A few applications of 2-ladders

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Theorem

Every distributive $(\vee, 0)$ -semilattice of cardinality \aleph_1 is isomorphic to

- $\text{Con}_c L$ for some lattice L (Huhn 1989),
- the normal subgroup lattice of some group, or the submodule lattice of some module (Růžička, Tůma, and W., 2007).

In both results above, the \aleph_1 bound is optimal (**different methods**).

... Looking ahead: representation results in cardinality \aleph_2 ?

Problem (Ditor 1984)

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

... Looking ahead: representation results in cardinality \aleph_2 ?

Problem (Ditor 1984)

Does there exist a 3-ladder of cardinality \aleph_2 ?

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

... Looking ahead: representation results in cardinality \aleph_2 ?

Problem (Ditor 1984)

Does there exist a 3-ladder of cardinality \aleph_2 ?

Theorem (W. 2008)

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

... Looking ahead: representation results in cardinality \aleph_2 ?

Problem (Ditor 1984)

Does there exist a 3-ladder of cardinality \aleph_2 ?

Theorem (W. 2008)

Suppose that either $\text{MA}(\aleph_1)$ holds or there exists a gap-1 morass. Then there exists a 3-ladder of cardinality \aleph_2 .

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

... Looking ahead: representation results in cardinality \aleph_2 ?

Problem (Ditor 1984)

Does there exist a 3-ladder of cardinality \aleph_2 ?

Theorem (W. 2008)

Suppose that either $\text{MA}(\aleph_1)$ holds or there exists a gap-1 morass. Then there exists a 3-ladder of cardinality \aleph_2 .

Gap-1 morasses exist in $\mathbf{L}[A]$ for any $A \subseteq \omega_1$; hence, **if there is no gap-1 morass, then ω_2 is inaccessible in \mathbf{L} .**

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

... Looking ahead: representation results in cardinality \aleph_2 ?

Problem (Ditor 1984)

Does there exist a 3-ladder of cardinality \aleph_2 ?

Theorem (W. 2008)

Suppose that either $\text{MA}(\aleph_1)$ holds or there exists a gap-1 morass. Then there exists a 3-ladder of cardinality \aleph_2 .

Gap-1 morasses exist in $\mathbf{L}[A]$ for any $A \subseteq \omega_1$; hence, **if there is no gap-1 morass, then ω_2 is inaccessible in \mathbf{L} .**

Corollary

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

... Looking ahead: representation results in cardinality \aleph_2 ?

Problem (Ditor 1984)

Does there exist a 3-ladder of cardinality \aleph_2 ?

Theorem (W. 2008)

Suppose that either $\text{MA}(\aleph_1)$ holds or there exists a gap-1 morass. Then there exists a 3-ladder of cardinality \aleph_2 .

Gap-1 morasses exist in $\mathbf{L}[A]$ for any $A \subseteq \omega_1$; hence, **if there is no gap-1 morass, then ω_2 is inaccessible in \mathbf{L} .**

Corollary

If there is no 3-ladder of cardinality \aleph_2 , then ω_2 is inaccessible in \mathbf{L} .

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

... Looking ahead: representation results in cardinality \aleph_2 ?

Problem (Ditor 1984)

Does there exist a 3-ladder of cardinality \aleph_2 ?

Theorem (W. 2008)

Suppose that either $\text{MA}(\aleph_1)$ holds or there exists a gap-1 morass. Then there exists a 3-ladder of cardinality \aleph_2 .

Gap-1 morasses exist in $\mathbf{L}[A]$ for any $A \subseteq \omega_1$; hence, **if there is no gap-1 morass, then ω_2 is inaccessible in \mathbf{L} .**

Corollary

If there is no 3-ladder of cardinality \aleph_2 , then ω_2 is inaccessible in \mathbf{L} .

On the other hand, **3-dimensional amalgamation fails in any variety with a nontrivial member.**

Critical points

- Set $\text{Con}_c \mathcal{V} := \{S \mid (\exists A \in \mathcal{V})(S \cong \text{Con}_c A)\}$, for any class \mathcal{V} of algebras.

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Critical points

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

- Set $\text{Con}_c \mathcal{V} := \{S \mid (\exists A \in \mathcal{V})(S \cong \text{Con}_c A)\}$, for any class \mathcal{V} of algebras.
- **Critical point crit** $(\mathcal{A}; \mathcal{B})$, for classes \mathcal{A} and \mathcal{B} of algebras: least possible cardinality of a member of $(\text{Con}_c \mathcal{A}) \setminus (\text{Con}_c \mathcal{B})$ (and ∞ if $\text{Con}_c \mathcal{A} \subseteq \text{Con}_c \mathcal{B}$).

Critical points

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

- Set $\text{Con}_c \mathcal{V} := \{S \mid (\exists A \in \mathcal{V})(S \cong \text{Con}_c A)\}$, for any class \mathcal{V} of algebras.
- **Critical point** $\text{crit}(\mathcal{A}; \mathcal{B})$, for classes \mathcal{A} and \mathcal{B} of algebras: least possible cardinality of a member of $(\text{Con}_c \mathcal{A}) \setminus (\text{Con}_c \mathcal{B})$ (and ∞ if $\text{Con}_c \mathcal{A} \subseteq \text{Con}_c \mathcal{B}$).

Theorem (Gillibert 2007)

Critical points

- Set $\text{Con}_c \mathcal{V} := \{S \mid (\exists A \in \mathcal{V})(S \cong \text{Con}_c A)\}$, for any class \mathcal{V} of algebras.
- **Critical point** $\mathbf{crit}(\mathcal{A}; \mathcal{B})$, for classes \mathcal{A} and \mathcal{B} of algebras: least possible cardinality of a member of $(\text{Con}_c \mathcal{A}) \setminus (\text{Con}_c \mathcal{B})$ (and ∞ if $\text{Con}_c \mathcal{A} \subseteq \text{Con}_c \mathcal{B}$).

Theorem (Gillibert 2007)

Let \mathcal{A} and \mathcal{B} be varieties of **algebras**, with \mathcal{A} locally finite and \mathcal{B} finitely generated congruence-distributive. If $\text{Con}_c \mathcal{A} \not\subseteq \text{Con}_c \mathcal{B}$, then $\mathbf{crit}(\mathcal{A}; \mathcal{B}) < \aleph_\omega$.

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Critical points

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

- Set $\text{Con}_c \mathcal{V} := \{S \mid (\exists A \in \mathcal{V})(S \cong \text{Con}_c A)\}$, for any class \mathcal{V} of algebras.
- **Critical point** $\mathbf{crit}(\mathcal{A}; \mathcal{B})$, for classes \mathcal{A} and \mathcal{B} of algebras: least possible cardinality of a member of $(\text{Con}_c \mathcal{A}) \setminus (\text{Con}_c \mathcal{B})$ (and ∞ if $\text{Con}_c \mathcal{A} \subseteq \text{Con}_c \mathcal{B}$).

Theorem (Gillibert 2007)

Let \mathcal{A} and \mathcal{B} be varieties of **algebras**, with \mathcal{A} locally finite and \mathcal{B} finitely generated congruence-distributive. If $\text{Con}_c \mathcal{A} \not\subseteq \text{Con}_c \mathcal{B}$, then $\mathbf{crit}(\mathcal{A}; \mathcal{B}) < \aleph_\omega$.

Theorem (Gillibert 2009)

Critical points

- Set $\text{Con}_c \mathcal{V} := \{S \mid (\exists A \in \mathcal{V})(S \cong \text{Con}_c A)\}$, for any class \mathcal{V} of algebras.
- **Critical point** $\mathbf{crit}(\mathcal{A}; \mathcal{B})$, for classes \mathcal{A} and \mathcal{B} of algebras: least possible cardinality of a member of $(\text{Con}_c \mathcal{A}) \setminus (\text{Con}_c \mathcal{B})$ (and ∞ if $\text{Con}_c \mathcal{A} \subseteq \text{Con}_c \mathcal{B}$).

Theorem (Gillibert 2007)

Let \mathcal{A} and \mathcal{B} be varieties of **algebras**, with \mathcal{A} locally finite and \mathcal{B} finitely generated congruence-distributive. If $\text{Con}_c \mathcal{A} \not\subseteq \text{Con}_c \mathcal{B}$, then $\mathbf{crit}(\mathcal{A}; \mathcal{B}) < \aleph_\omega$.

Theorem (Gillibert 2009)

Let \mathcal{A} and \mathcal{B} be varieties of **lattices** such that every simple member of \mathcal{B} has a prime interval. If $\mathcal{A} \not\subseteq \mathcal{B}$ and $\mathcal{A} \not\subseteq \mathcal{B}^{\text{dual}}$, then $\text{Con}_c \mathcal{A} \not\subseteq \text{Con}_c \mathcal{B}$ and $\mathbf{crit}(\mathcal{A}; \mathcal{B}) \leq \aleph_2$. The bound \aleph_2 is optimal.

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

A question about critical points

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, <\lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Question

A question about critical points

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, <\lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Question

Are there varieties \mathcal{A} and \mathcal{B} , on finite similarity types, such that $\mathbf{crit}(\mathcal{A}; \mathcal{B}) = \aleph_3$?

A question about critical points

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Question

Are there varieties \mathcal{A} and \mathcal{B} , on finite similarity types, such that $\mathbf{crit}(\mathcal{A}; \mathcal{B}) = \aleph_3$?

Formally related to the existence problem of 3-ladders of cardinality $\aleph_2 \dots$

A question about critical points

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Question

Are there varieties \mathcal{A} and \mathcal{B} , on finite similarity types, such that $\mathbf{crit}(\mathcal{A}; \mathcal{B}) = \aleph_3$?

Formally related to the existence problem of 3-ladders of cardinality \aleph_2 . . . On the other hand, 3-dimensional amalgamation fails!

What to think?

What makes critical points small?

Lifters, free
sets, ladders

The Conc
functor

k -ladders

Critical points

From lifting
objects to
lifting
diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating
Kuratowski
indexes of
finite posets

- Using the chain ω makes critical points $\geq \aleph_1$.

What makes critical points small?

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

- Using the chain ω makes critical points $\geq \aleph_1$.
- Using 2-ladders of cardinality \aleph_1 makes critical points $\geq \aleph_2$.

What makes critical points small?

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

- Using the chain ω makes critical points $\geq \aleph_1$.
- Using 2-ladders of cardinality \aleph_1 makes critical points $\geq \aleph_2$.
- Maybe some day, 3-ladders of cardinality \aleph_2 will help finding a critical point $\aleph_3 \dots$

What makes critical points small?

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

- Using the chain ω makes critical points $\geq \aleph_1$.
- Using 2-ladders of cardinality \aleph_1 makes critical points $\geq \aleph_2$.
- Maybe some day, 3-ladders of cardinality \aleph_2 will help finding a critical point \aleph_3 ...
- ... But what makes critical points small?

What makes critical points small?

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

- Using the chain ω makes critical points $\geq \aleph_1$.
- Using 2-ladders of cardinality \aleph_1 makes critical points $\geq \aleph_2$.
- Maybe some day, 3-ladders of cardinality \aleph_2 will help finding a critical point \aleph_3 ...
- ... But what makes critical points small?
- Goto category theory...

General categorical settings

Lifters, free
sets, ladders

The Con_c
functor

k -ladders

Critical points

From lifting
objects to
lifting
diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating
Kuratowski
indexes of
finite posets

We are given **categories** \mathcal{A} , \mathcal{B} , \mathcal{S} together with **functors**
 $\Phi: \mathcal{A} \rightarrow \mathcal{S}$ and $\Psi: \mathcal{B} \rightarrow \mathcal{S}$.

General categorical settings

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters


$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

We are given **categories** \mathcal{A} , \mathcal{B} , \mathcal{S} together with **functors** $\Phi: \mathcal{A} \rightarrow \mathcal{S}$ and $\Psi: \mathcal{B} \rightarrow \mathcal{S}$. We are trying to find a **functor** $\Gamma: \mathcal{A} \rightarrow \mathcal{B}$ such that $\Phi(A) \cong \Psi\Gamma(A)$, **naturally in \mathcal{A}** , for “many” (ideally, all) $A \in \mathcal{A}$.

General categorical settings

We are given **categories** \mathcal{A} , \mathcal{B} , \mathcal{S} together with **functors** $\Phi: \mathcal{A} \rightarrow \mathcal{S}$ and $\Psi: \mathcal{B} \rightarrow \mathcal{S}$. We are trying to find a **functor** $\Gamma: \mathcal{A} \rightarrow \mathcal{B}$ such that $\Phi(A) \cong \Psi\Gamma(A)$, **naturally in \mathcal{A}** , for “many” (ideally, all) $A \in \mathcal{A}$.



Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams



λ -lifters

$(\kappa, <\lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

General categorical settings

We are given **categories** \mathcal{A} , \mathcal{B} , \mathcal{S} together with **functors** $\Phi: \mathcal{A} \rightarrow \mathcal{S}$ and $\Psi: \mathcal{B} \rightarrow \mathcal{S}$. We are trying to find a **functor** $\Gamma: \mathcal{A} \rightarrow \mathcal{B}$ such that $\Phi(A) \cong \Psi\Gamma(A)$, **naturally in \mathcal{A}** , for “many” (ideally, all) $A \in \mathcal{A}$.



Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams



λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

General categorical settings

We are given **categories** \mathcal{A} , \mathcal{B} , \mathcal{S} together with **functors** $\Phi: \mathcal{A} \rightarrow \mathcal{S}$ and $\Psi: \mathcal{B} \rightarrow \mathcal{S}$. We are trying to find a **functor** $\Gamma: \mathcal{A} \rightarrow \mathcal{B}$ such that $\Phi(A) \cong \Psi\Gamma(A)$, **naturally in \mathcal{A}** , for “many” (ideally, all) $A \in \mathcal{A}$.



Hence we need an assumption of the form “for many $A \in \mathcal{A}$, there exists $B \in \mathcal{B}$ such that $\Phi(A) \cong \Psi(B)$ ”.

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams



λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

General categorical settings

We are given **categories** \mathcal{A} , \mathcal{B} , \mathcal{S} together with **functors** $\Phi: \mathcal{A} \rightarrow \mathcal{S}$ and $\Psi: \mathcal{B} \rightarrow \mathcal{S}$. We are trying to find a **functor** $\Gamma: \mathcal{A} \rightarrow \mathcal{B}$ such that $\Phi(A) \cong \Psi\Gamma(A)$, **naturally in \mathcal{A}** , for “many” (ideally, all) $A \in \mathcal{A}$.



Hence we need an assumption of the form “for many $A \in \mathcal{A}$, there exists $B \in \mathcal{B}$ such that $\Phi(A) \cong \Psi(B)$ ”. Ask for $\Gamma: \mathcal{A} \rightarrow \mathcal{B}$ to be a **functor** (at least on a large enough subcategory of \mathcal{A}).

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

From lifting objects to lifting diagrams

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points


From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

CLL Theorem (Gillibert and W., 2008)



From lifting objects to lifting diagrams

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

CLL Theorem (Gillibert and W., 2008)

In the context above, suppose that

$$(\text{For many } A \in \mathcal{A})(\exists B \in \mathcal{B})(\Phi(A) \cong \Psi(B)).$$

From lifting objects to lifting diagrams

CLL Theorem (Gillibert and W., 2008)

In the context above, suppose that

$$(\text{For many } A \in \mathcal{A})(\exists B \in \mathcal{B})(\Phi(A) \cong \Psi(B)).$$

Suppose that the categorical settings are **nice** (ladders...)

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

From lifting objects to lifting diagrams

CLL Theorem (Gillibert and W., 2008)

In the context above, suppose that

$$(\text{For many } A \in \mathcal{A})(\exists B \in \mathcal{B})(\Phi(A) \cong \Psi(B)).$$

Suppose that the categorical settings are **nice** (**ladders...**) and let P be a **nice** poset. Then

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

From lifting objects to lifting diagrams

CLL Theorem (Gillibert and W., 2008)

In the context above, suppose that

$$(\text{For many } A \in \mathcal{A})(\exists B \in \mathcal{B})(\Phi(A) \cong \Psi(B)).$$

Suppose that the categorical settings are **nice** (**ladders...**) and let P be a **nice** poset. Then

$$(\text{For many } A \in \mathcal{A}^P)(\exists B \in \mathcal{B}^P)(\Phi(A) \cong \Psi(B)).$$

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

From lifting objects to lifting diagrams

CLL Theorem (Gillibert and W., 2008)

In the context above, suppose that

$$(\text{For many } A \in \mathcal{A})(\exists B \in \mathcal{B})(\Phi(A) \cong \Psi(B)).$$

Suppose that the categorical settings are **nice** (**ladders**...) and let P be a **nice** poset. Then

$$(\text{For many } A \in \mathcal{A}^P)(\exists B \in \mathcal{B}^P)(\Phi(A) \cong \Psi(B)).$$

That is, if Ψ lifts many **objects**, then it lifts many (P -indexed) **diagrams**.

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

From lifting objects to lifting diagrams

CLL Theorem (Gillibert and W., 2008)

In the context above, suppose that

$$(\text{For many } A \in \mathcal{A})(\exists B \in \mathcal{B})(\Phi(A) \cong \Psi(B)).$$

Suppose that the categorical settings are **nice** (**ladders...**) and let P be a **nice** poset. Then

$$(\text{For many } A \in \mathcal{A}^P)(\exists B \in \mathcal{B}^P)(\Phi(A) \cong \Psi(B)).$$

That is, if Ψ lifts many **objects**, then it lifts many (P -indexed) **diagrams**.

CLL turns **diagram counterexamples** to **object counterexamples**.

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

From lifting objects to lifting diagrams

CLL Theorem (Gillibert and W., 2008)

In the context above, suppose that

$$(\text{For many } A \in \mathcal{A})(\exists B \in \mathcal{B})(\Phi(A) \cong \Psi(B)).$$

Suppose that the categorical settings are **nice** (**ladders...**) and let P be a **nice** poset. Then

$$(\text{For many } A \in \mathcal{A}^P)(\exists B \in \mathcal{B}^P)(\Phi(A) \cong \Psi(B)).$$

That is, if Ψ lifts many **objects**, then it lifts many (P -indexed) **diagrams**.

CLL turns **diagram counterexamples** to **object counterexamples**.
For critical points, Φ and Ψ are both Con_c .

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

What is special about the poset P ?

- For a subset X in a poset P , denote by ∇X the set of all minimal elements of $P \uparrow X := \{p \in P \mid X \leq p\}$.

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

What is special about the poset P ?

- For a subset X in a poset P , denote by ∇X the set of all minimal elements of $P \uparrow X := \{p \in P \mid X \leq p\}$.
- Say that X is ∇ -closed if $\nabla Y \subseteq X$, $\forall Y \subseteq X$ finite.

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

What is special about the poset P ?

- For a subset X in a poset P , denote by ∇X the set of all minimal elements of $P \uparrow X := \{p \in P \mid X \leq p\}$.
- Say that X is **∇ -closed** if $\nabla Y \subseteq X, \forall Y \subseteq X$ finite.

Definition (Gillibert and W., 2008)

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

What is special about the poset P ?

- For a subset X in a poset P , denote by ∇X the set of all minimal elements of $P \uparrow X := \{p \in P \mid X \leq p\}$.
- Say that X is **∇ -closed** if $\nabla Y \subseteq X, \forall Y \subseteq X$ finite.

Definition (Gillibert and W., 2008)

A poset P is

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

What is special about the poset P ?

- For a subset X in a poset P , denote by ∇X the set of all minimal elements of $P \uparrow X := \{p \in P \mid X \leq p\}$.
- Say that X is **∇ -closed** if $\nabla Y \subseteq X, \forall Y \subseteq X$ finite.

Definition (Gillibert and W., 2008)

A poset P is

- a **pseudo join-semilattice** if $P \uparrow X$ is a finitely generated upper subset, $\forall X \subseteq P$ finite;

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

What is special about the poset P ?

- For a subset X in a poset P , denote by ∇X the set of all minimal elements of $P \uparrow X := \{p \in P \mid X \leq p\}$.
- Say that X is **∇ -closed** if $\nabla Y \subseteq X$, $\forall Y \subseteq X$ finite.

Definition (Gillibert and W., 2008)

A poset P is

- a **pseudo join-semilattice** if $P \uparrow X$ is a finitely generated upper subset, $\forall X \subseteq P$ finite;
- **supported** if it is a pseudo join-semilattice and the ∇ -closure of any finite subset of P is finite (sometimes such a P is called **mub-complete**);

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

What is special about the poset P ?

- For a subset X in a poset P , denote by ∇X the set of all minimal elements of $P \uparrow X := \{p \in P \mid X \leq p\}$.
- Say that X is **∇ -closed** if $\nabla Y \subseteq X$, $\forall Y \subseteq X$ finite.

Definition (Gillibert and W., 2008)

A poset P is

- a **pseudo join-semilattice** if $P \uparrow X$ is a finitely generated upper subset, $\forall X \subseteq P$ finite;
- **supported** if it is a pseudo join-semilattice and the ∇ -closure of any finite subset of P is finite (sometimes such a P is called **mub-complete**);
- an **almost join-semilattice** if it is a pseudo join-semilattice and $P \downarrow a$ is a join-semilattice $\forall a \in P$.

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Examples (pseudo join-semilattice, etc.)

Lifters, free sets, ladders

- almost join-semilattice \Rightarrow supported \Rightarrow pseudo join-semilattice.

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, \langle \lambda \rangle) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Examples (pseudo join-semilattice, etc.)

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

- almost join-semilattice \Rightarrow supported \Rightarrow pseudo join-semilattice.
- The following posets separate the three classes (and the leftmost one is not a pseudo join-semilattice):

Examples (pseudo join-semilattice, etc.)

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points


From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

- almost join-semilattice \Rightarrow supported \Rightarrow pseudo join-semilattice.
- The following posets separate the three classes (and the leftmost one is not a pseudo join-semilattice):



Norm-coverings, sharp ideals

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, \langle \lambda \rangle) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

- A **norm-covering** of a poset P is a pseudo join-semilattice X , together with an isotone map $\partial: X \rightarrow P$.

Norm-coverings, sharp ideals

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, <\lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

- A **norm-covering** of a poset P is a pseudo join-semilattice X , together with an isotone map $\partial: X \rightarrow P$.
- An ideal (i.e., a nonempty, upward directed, lower subset) \mathbf{x} of X is **sharp** if $\{\partial x \mid x \in \mathbf{x}\}$ has a largest element, then denoted by $\partial \mathbf{x}$.

Norm-coverings, sharp ideals

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

- A **norm-covering** of a poset P is a pseudo join-semilattice X , together with an isotone map $\partial: X \rightarrow P$.
- An ideal (i.e., a nonempty, upward directed, lower subset) \mathbf{x} of X is **sharp** if $\{\partial x \mid x \in \mathbf{x}\}$ has a largest element, then denoted by $\partial \mathbf{x}$.
- Then set $\mathbf{X}^= := \{\mathbf{x} \in \mathbf{X} \mid \partial \mathbf{x} \text{ not maximal}\}$, for every set \mathbf{X} of sharp ideals.

λ -lifters

The posets P involved in the statement of CLL are those for which there exists a λ -lifter. Here, λ may be thought of as an upper bound for the sizes of the **diagram data**, while the cardinality of X may be thought as an upper bound for the sizes of the **object data**.

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, <\lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

λ -lifters

The posets P involved in the statement of CLL are those for which there exists a λ -lifter. Here, λ may be thought of as an upper bound for the sizes of the **diagram data**, while the cardinality of X may be thought as an upper bound for the sizes of the **object data**.

Definition (Gillibert and W. 2008)

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

λ -lifters

The posets P involved in the statement of CLL are those for which there exists a λ -lifter. Here, λ may be thought of as an upper bound for the sizes of the **diagram data**, while the cardinality of X may be thought as an upper bound for the sizes of the **object data**.

Definition (Gillibert and W. 2008)

For an infinite cardinal λ , a λ -lifter of a poset P is a pair (X, \mathbf{X}) , where X is a norm-covering of P , \mathbf{X} is a set of sharp ideals of X , and

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

λ -lifters

The posets P involved in the statement of CLL are those for which there exists a λ -lifter. Here, λ may be thought of as an upper bound for the sizes of the **diagram data**, while the cardinality of X may be thought as an upper bound for the sizes of the **object data**.

Definition (Gillibert and W. 2008)

For an infinite cardinal λ , a λ -lifter of a poset P is a pair (X, \mathbf{X}) , where X is a norm-covering of P , \mathbf{X} is a set of sharp ideals of X , and

- (i) Every principal ideal of $\mathbf{X}^=$ has cardinality $< \text{cf}(\lambda)$.

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

λ -lifters

The posets P involved in the statement of CLL are those for which there exists a λ -lifter. Here, λ may be thought of as an upper bound for the sizes of the **diagram data**, while the cardinality of X may be thought as an upper bound for the sizes of the **object data**.

Definition (Gillibert and W. 2008)

For an infinite cardinal λ , a λ -lifter of a poset P is a pair (X, \mathbf{X}) , where X is a norm-covering of P , \mathbf{X} is a set of sharp ideals of X , and

- (i) Every principal ideal of $\mathbf{X}^=$ has cardinality $< \text{cf}(\lambda)$.
- (ii) For every map $S: \mathbf{X}^= \rightarrow [X]^{<\lambda}$ there exists an isotone section σ of ∂ such that $S\sigma(a) \cap \sigma(b) \subseteq \sigma(a)$ for all $a < b$ in P .

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, <\lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

λ -lifters

The posets P involved in the statement of CLL are those for which there exists a λ -lifter. Here, λ may be thought of as an upper bound for the sizes of the **diagram data**, while the cardinality of X may be thought as an upper bound for the sizes of the **object data**.

Definition (Gillibert and W. 2008)

For an infinite cardinal λ , a λ -lifter of a poset P is a pair (X, \mathbf{X}) , where X is a norm-covering of P , \mathbf{X} is a set of sharp ideals of X , and

- (i) Every principal ideal of $\mathbf{X}^=$ has cardinality $< \text{cf}(\lambda)$.
- (ii) For every map $S: \mathbf{X}^= \rightarrow [X]^{<\lambda}$ there exists an isotone section σ of ∂ such that $S\sigma(a) \cap \sigma(b) \subseteq \sigma(a)$ for all $a < b$ in P .
- (iii) If $\lambda = \aleph_0$, then X is supported.

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, <\lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

The shape of liftable posets

A severe restriction about the shape of P is that **If P has a λ -lifter, then it is an almost join-semilattice.** More precisely,

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, \langle \lambda \rangle) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

The shape of liftable posets

A severe restriction about the shape of P is that **If P has a λ -lifter, then it is an almost join-semilattice.** More precisely,

Theorem (Gillibert and W., 2008)

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, <\lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

The shape of liftable posets

A severe restriction about the shape of P is that **If P has a λ -lifter, then it is an almost join-semilattice.** More precisely,

Theorem (Gillibert and W., 2008)

Let λ be an infinite cardinal and let (X, \mathbf{X}) be a λ -lifter of a poset P . Put $\kappa := \text{card } \mathbf{X}$. Then the following statements hold:

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

The shape of liftable posets

A severe restriction about the shape of P is that **If P has a λ -lifter, then it is an almost join-semilattice.** More precisely,

Theorem (Gillibert and W., 2008)

Let λ be an infinite cardinal and let (X, \mathbf{X}) be a λ -lifter of a poset P . Put $\kappa := \text{card } \mathbf{X}$. Then the following statements hold:

- (i) P is a disjoint union of finitely many almost join-semilattices with zero.

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

The shape of liftable posets

A severe restriction about the shape of P is that **If P has a λ -lifter, then it is an almost join-semilattice.** More precisely,

Theorem (Gillibert and W., 2008)

Let λ be an infinite cardinal and let (X, \mathbf{X}) be a λ -lifter of a poset P . Put $\kappa := \text{card } \mathbf{X}$. Then the following statements hold:

- (i) P is a disjoint union of finitely many almost join-semilattices with zero.
- (ii) For every **isotone** map $F: [\kappa]^{<\text{cf}(\lambda)} \rightarrow [\kappa]^{<\lambda}$, there exists a **one-to-one** map $\sigma: P \rightarrow \kappa$ such that

$$(\forall a < b \text{ in } P)(F\sigma(P \downarrow a) \cap \sigma(P \downarrow b) \subseteq \sigma(P \downarrow a)).$$

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, <\lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

These posets are too heavy...

Lifters, free
sets, ladders

The Conc
functor

k -ladders

Critical points

From lifting
objects to
lifting
diagrams

λ -lifters

$(\kappa, <\lambda) \rightsquigarrow P$

Calculating
Kuratowski
indexes of
finite posets

In particular, none of the following posets has a lifter:

These posets are too heavy...

Lifters, free sets, ladders

The Conc functor

k -ladders


Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets



Well-foundedness and liftability

Lifters, free sets, ladders

Theorem (Gillibert + W. 2010)

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Well-foundedness and liftability

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Theorem (Gillibert + W. 2010)

Let λ be an infinite cardinal such that

$$(\forall \mu < \text{cf}(\lambda)) (\mu^{\aleph_0} < \lambda).$$

Well-foundedness and liftability

Theorem (Gillibert + W. 2010)

Let λ be an infinite cardinal such that

$$(\forall \mu < \text{cf}(\lambda)) (\mu^{\aleph_0} < \lambda).$$

If a poset has a λ -lifter, then it is **well-founded**.

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Well-foundedness and liftability

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Theorem (Gillibert + W. 2010)

Let λ be an infinite cardinal such that

$$(\forall \mu < \text{cf}(\lambda)) (\mu^{\aleph_0} < \lambda).$$

If a poset has a λ -lifter, then it is **well-founded**.

The poset $(\omega + 1)^{\text{dual}}$:

$$0 > 1 > 2 > \dots > \omega.$$

Well-foundedness and liftability

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, <\lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Theorem (Gillibert + W. 2010)

Let λ be an infinite cardinal such that

$$(\forall \mu < \text{cf}(\lambda)) (\mu^{\aleph_0} < \lambda).$$

If a poset has a λ -lifter, then it is **well-founded**.

The poset $(\omega + 1)^{\text{dual}}$:

$$0 > 1 > 2 > \dots > \omega.$$

Corollary

Well-foundedness and liftability

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Theorem (Gillibert + W. 2010)

Let λ be an infinite cardinal such that

$$(\forall \mu < \text{cf}(\lambda)) (\mu^{\aleph_0} < \lambda).$$

If a poset has a λ -lifter, then it is **well-founded**.

The poset $(\omega + 1)^{\text{dual}}$:

$$0 > 1 > 2 > \dots > \omega.$$

Corollary

$(\omega + 1)^{\text{dual}}$ has no $(2^{\aleph_0})^+$ -lifter.

A question about liftability of infinite posets

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, <\lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Question

A question about liftability of infinite posets

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Question

Let λ be an infinite cardinal. Does $(\omega + 1)^{\text{dual}}$ have a λ -lifter?

A question about liftability of infinite posets

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Question

Let λ be an infinite cardinal. Does $(\omega + 1)^{\text{dual}}$ have a λ -lifter?

For $\lambda := \aleph_1$, this is related to the following question:

A question about liftability of infinite posets

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Question

Let λ be an infinite cardinal. Does $(\omega + 1)^{\text{dual}}$ have a λ -lifter?

For $\lambda := \aleph_1$, this is related to the following question:

Question

A question about liftability of infinite posets

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Question

Let λ be an infinite cardinal. Does $(\omega + 1)^{\text{dual}}$ have a λ -lifter?

For $\lambda := \aleph_1$, this is related to the following question:

Question

Does there exist a cardinal κ such that for every **isotone** $f: [\kappa]^{\leq \aleph_0} \rightarrow [\kappa]^{\leq \aleph_0}$, there exists a sequence $(\xi_n \mid n < \omega)$ of elements of κ such that $\xi_n \notin f(\{\xi_k \mid k > n\})$ for each $n < \omega$?

A question about liftability of infinite posets

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Question

Let λ be an infinite cardinal. Does $(\omega + 1)^{\text{dual}}$ have a λ -lifter?

For $\lambda := \aleph_1$, this is related to the following question:

Question

Does there exist a cardinal κ such that for every **isotone** $f: [\kappa]^{\leq \aleph_0} \rightarrow [\kappa]^{\leq \aleph_0}$, there exists a sequence $(\xi_n \mid n < \omega)$ of elements of κ such that $\xi_n \notin f(\{\xi_k \mid k > n\})$ for each $n < \omega$?

If “isotone” is removed from the assumptions above, then the answer to the corresponding question is easily seen to be negative.

A diagram counterexample with no associated object counterexample so far

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

The following uses 1998 results from Kearnes and Szendrei on commutator theory.

Theorem (Tůma and W., 2006)

A diagram counterexample with no associated object counterexample so far

The following uses 1998 results from Kearnes and Szendrei on commutator theory.

Theorem (Tůma and W., 2006)

There exists a **diagram**, indexed by a **finite poset** P , of **finite Boolean semilattices** and **$(\vee, 0)$ -embeddings**, which cannot be lifted, with respect to the Con_c functor, in any **variety** satisfying a nontrivial congruence identity.

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, <\lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

A diagram counterexample with no associated object counterexample so far

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, <\lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

The following uses 1998 results from Kearnes and Szendrei on commutator theory.

Theorem (Tůma and W., 2006)

There exists a **diagram**, indexed by a **finite poset** P , of **finite Boolean semilattices** and **$(\vee, 0)$ -embeddings**, which cannot be lifted, with respect to the Con_c functor, in any **variety** satisfying a nontrivial congruence identity.

- In particular, this diagram **cannot** be lifted by **lattices**, **majority algebras**, **groups**, **modules**, etc.

A diagram counterexample with no associated object counterexample so far

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, <\lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets


The following uses 1998 results from Kearnes and Szendrei on commutator theory.

Theorem (Tůma and W., 2006)

There exists a **diagram**, indexed by a **finite poset** P , of **finite Boolean semilattices** and **$(\vee, 0)$ -embeddings**, which cannot be lifted, with respect to the Con_c functor, in any **variety** satisfying a nontrivial congruence identity.

- In particular, this diagram **cannot** be lifted by **lattices**, **majority algebras**, **groups**, **modules**, etc.
- Nevertheless, it **can** be lifted by **groupoids**.

This diagram has the following form:



Lifters, free
sets, ladders

The Conc
functor

k -ladders

Critical points

From lifting
objects to
lifting
diagrams

λ -lifters

$(\kappa, <\lambda) \rightsquigarrow P$

Calculating
Kuratowski
indexes of
finite posets

- Its underlying poset does not have any lifter (because it is not an almost join-semilattice).

- Its underlying poset does not have any lifter (because it is **not an almost join-semilattice**).
- Thus we cannot apply CLL (or its known extensions) to it.

- Its underlying poset does not have any lifter (because it is **not an almost join-semilattice**).
- Thus we cannot apply CLL (or its known extensions) to it.
- **In particular, it is still unknown whether every distributive $(\vee, 0)$ -semilattice is isomorphic to $\text{Con}_c M$ for some majority algebra M .**

The $(\kappa, <\lambda) \rightsquigarrow P$ notation

The following is a more user-friendly variant of the definition of a lifter.

Definition (Gillibert and W., 2008)

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, <\lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

The $(\kappa, <\lambda) \rightsquigarrow P$ notation

The following is a more user-friendly variant of the definition of a lifter.

Definition (Gillibert and W., 2008)

For infinite cardinals κ, λ and a poset P , let $(\kappa, <\lambda) \rightsquigarrow P$ hold if for every mapping $F: \mathfrak{P}(\kappa) \rightarrow [\kappa]^{<\lambda}$ there exists a one-to-one map $\sigma: P \rightarrow \kappa$ such that

$$(\forall x < y \text{ in } P)(F\sigma(P \downarrow x) \cap \sigma(P \downarrow y) \subseteq \sigma(P \downarrow x)).$$

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, <\lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

The $(\kappa, <\lambda) \rightsquigarrow P$ notation

The following is a more user-friendly variant of the definition of a lifter.

Definition (Gillibert and W., 2008)

For infinite cardinals κ, λ and a poset P , let $(\kappa, <\lambda) \rightsquigarrow P$ hold if for every mapping $F: \mathfrak{P}(\kappa) \rightarrow [\kappa]^{<\lambda}$ there exists a one-to-one map $\sigma: P \rightarrow \kappa$ such that

$$(\forall x < y \text{ in } P)(F\sigma(P \downarrow x) \cap \sigma(P \downarrow y) \subseteq \sigma(P \downarrow x)).$$

In case P is **lower finite**, it is sufficient to replace $P \downarrow z$ by $J(P) \downarrow z$ in the statement above (**convenient for verifications on specific posets**).

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, <\lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

The liftable finite posets

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, \langle \lambda \rangle) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Theorem (Gillibert and W., 2008)

The liftable finite posets

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Theorem (Gillibert and W., 2008)

Let P be a finite poset and let λ be an infinite cardinal. Then P has a λ -lifter iff P is a disjoint union of almost join-semilattices with zero.

The liftable finite posets

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Theorem (Gillibert and W., 2008)

Let P be a finite poset and let λ be an infinite cardinal. Then P has a λ -lifter iff P is a disjoint union of almost join-semilattices with zero.

- The smallest possible cardinality of a λ -lifter of a finite poset P can be estimated precisely, via the $(\kappa, < \lambda) \rightsquigarrow P$ notation.

The liftable finite posets

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Theorem (Gillibert and W., 2008)

Let P be a finite poset and let λ be an infinite cardinal. Then P has a λ -lifter iff P is a disjoint union of almost join-semilattices with zero.

- The smallest possible cardinality of a λ -lifter of a finite poset P can be estimated precisely, *via* the $(\kappa, < \lambda) \rightsquigarrow P$ notation.
- A convenient way to do this is to introduce the **Kuratowski index**.

The Kuratowski index of a finite poset

Definition (Gillibert and W., 2008)

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, <\lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

The Kuratowski index of a finite poset

Definition (Gillibert and W., 2008)

The **Kuratowski index** of a finite poset P , denoted by $\text{kur}(P)$, is defined as 0 if P is an antichain, and, otherwise, as the least $n > 0$ such that $(\lambda^{+(n-1)}, < \lambda) \rightsquigarrow P$ for each infinite cardinal λ .

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

The Kuratowski index of a finite poset

Definition (Gillibert and W., 2008)

The **Kuratowski index** of a finite poset P , denoted by $\text{kur}(P)$, is defined as 0 if P is an antichain, and, otherwise, as the least $n > 0$ such that $(\lambda^{+(n-1)}, < \lambda) \rightsquigarrow P$ for each infinite cardinal λ .

Theorem (Gillibert + W., 2008)

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

The Kuratowski index of a finite poset

Definition (Gillibert and W., 2008)

The **Kuratowski index** of a finite poset P , denoted by $\text{kur}(P)$, is defined as 0 if P is an antichain, and, otherwise, as the least $n > 0$ such that $(\lambda^{+(n-1)}, < \lambda) \rightsquigarrow P$ for each infinite cardinal λ .

Theorem (Gillibert + W., 2008)

The Kuratowski index of P always exists, and $\text{kur}(P) \leq \dim(P) \leq \text{width } J(P) \leq \text{card } J(P)$. Furthermore, if P is a join-semilattice, then $\text{breadth}(P) \leq \text{kur}(P)$.

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

The Kuratowski index of a finite poset

Definition (Gillibert and W., 2008)

The **Kuratowski index** of a finite poset P , denoted by $\text{kur}(P)$, is defined as 0 if P is an antichain, and, otherwise, as the least $n > 0$ such that $(\lambda^{+(n-1)}, < \lambda) \rightsquigarrow P$ for each infinite cardinal λ .

Theorem (Gillibert + W., 2008)

The Kuratowski index of P always exists, and $\text{kur}(P) \leq \dim(P) \leq \text{width } J(P) \leq \text{card } J(P)$. Furthermore, if P is a join-semilattice, then $\text{breadth}(P) \leq \text{kur}(P)$.

Theorem (Gillibert + W., 2008)

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

The Kuratowski index of a finite poset

Definition (Gillibert and W., 2008)

The **Kuratowski index** of a finite poset P , denoted by $\text{kur}(P)$, is defined as 0 if P is an antichain, and, otherwise, as the least $n > 0$ such that $(\lambda^{+(n-1)}, < \lambda) \rightsquigarrow P$ for each infinite cardinal λ .

Theorem (Gillibert + W., 2008)

The Kuratowski index of P always exists, and $\text{kur}(P) \leq \dim(P) \leq \text{width } J(P) \leq \text{card } J(P)$. Furthermore, if P is a join-semilattice, then $\text{breadth}(P) \leq \text{kur}(P)$.

Theorem (Gillibert + W., 2008)

For every infinite cardinal λ , every nontrivial finite almost join-semilattice P with zero has a λ -lifter (X, \mathbf{X}) with $\text{card } X = \text{card } \mathbf{X} = \lambda^{+(\text{kur}(P)-1)}$.

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Sometimes, $\text{kur}(P)$ is easy to calculate:

- $\text{kur}(T) = 1$ whenever T is a nontrivial finite **tree**.

Lifters, free
sets, ladders

The Conc
functor

k -ladders

Critical points

From lifting
objects to
lifting
diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating
Kuratowski
indexes of
finite posets

Sometimes, $\text{kur}(P)$ is easy to calculate:

- $\text{kur}(T) = 1$ whenever T is a nontrivial finite **tree**.
- $P \subseteq Q$ implies that $\text{kur}(P) \leq \text{kur}(Q)$.

Lifters, free
sets, ladders

The Conc
functor

k -ladders

Critical points

From lifting
objects to
lifting
diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating
Kuratowski
indexes of
finite posets

Sometimes, $\text{kur}(P)$ is easy to calculate:

- $\text{kur}(T) = 1$ whenever T is a nontrivial finite **tree**.
- $P \subseteq Q$ implies that $\text{kur}(P) \leq \text{kur}(Q)$.
- $\text{kur}(P \times Q) \leq \text{kur}(P) + \text{kur}(Q)$, for all finite posets P and Q with zero.

Lifters, free
sets, ladders

The Conc
functor

k -ladders

Critical points

From lifting
objects to
lifting
diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating
Kuratowski
indexes of
finite posets

Sometimes, $\text{kur}(P)$ is easy to calculate:

- $\text{kur}(T) = 1$ whenever T is a nontrivial finite tree.
- $P \subseteq Q$ implies that $\text{kur}(P) \leq \text{kur}(Q)$.
- $\text{kur}(P \times Q) \leq \text{kur}(P) + \text{kur}(Q)$, for all finite posets P and Q with zero.
- Thus $\text{kur}(P) = n$ in case P is a product of n nontrivial finite trees.

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Sometimes, $\text{kur}(P)$ is easy to calculate:

- $\text{kur}(T) = 1$ whenever T is a nontrivial finite tree.
- $P \subseteq Q$ implies that $\text{kur}(P) \leq \text{kur}(Q)$.
- $\text{kur}(P \times Q) \leq \text{kur}(P) + \text{kur}(Q)$, for all finite posets P and Q with zero.
- Thus $\text{kur}(P) = n$ in case P is a product of n nontrivial finite trees.
- In particular, $\text{kur}(2^n) = n$ (also follows from Kuratowski's Free Set Theorem).

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Sometimes, $\text{kur}(P)$ is easy to calculate:

- $\text{kur}(T) = 1$ whenever T is a nontrivial finite **tree**.
- $P \subseteq Q$ implies that $\text{kur}(P) \leq \text{kur}(Q)$.
- $\text{kur}(P \times Q) \leq \text{kur}(P) + \text{kur}(Q)$, for all finite posets P and Q with zero.
- Thus $\text{kur}(P) = n$ in case P is a product of n nontrivial finite trees.
- In particular, $\text{kur}(\mathbf{2}^n) = n$ (also follows from Kuratowski's Free Set Theorem).
- **Warning:** the $P \mapsto \text{kur}(P)$ function is **not absolute** (in the set-theoretical sense).

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams


λ -lifters

$(\kappa, <\lambda) \rightsquigarrow P$


Calculating Kuratowski indexes of finite posets

... sometimes, not so easy:

- Consider the finite lattices P and Q represented below.



P



Q

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams



λ -lifters

$(\kappa, <\lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

... sometimes, not so easy:

- Consider the finite lattices P and Q represented below.



- $\text{breadth}(P) = \text{breadth}(Q) = 2$ while $\text{dim}(P) = \text{dim}(Q) = 3$.

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams


λ -lifters

$(\kappa, \langle \lambda \rangle) \rightsquigarrow P$


Calculating Kuratowski indexes of finite posets

... sometimes, not so easy:

- Consider the finite lattices P and Q represented below.



P



Q

- $\text{breadth}(P) = \text{breadth}(Q) = 2$ while $\text{dim}(P) = \text{dim}(Q) = 3$.
- Thus $2 \leq \text{kur}(P) \leq \text{kur}(Q) \leq 3$.

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, <\lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Lifters, free
sets, ladders

The Con_c
functor

k -ladders

Critical points

From lifting
objects to
lifting
diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating
Kuratowski
indexes of
finite posets

We don't know more. For example, $\text{kur}(P) = 2$ would mean
the following:

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, <\lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

We don't know more. For example, $\text{kur}(P) = 2$ would mean the following:

For every infinite cardinal λ and every $F: [\lambda^+]^{<\omega} \rightarrow [\lambda^+]^{<\lambda}$, there are distinct $\xi_0, \xi_1, \xi_2, \eta_0, \eta_1, \eta_2 < \lambda$ such that $\xi_i \notin F(\{\xi_j, \eta_j\})$, $\eta_i \notin F(\{\xi_j, \eta_j\})$, and $\eta_i \notin F(\{\xi_0, \xi_1, \xi_2\})$ for all $i \neq j$ in $\{0, 1, 2\}$.

Truncated m -cubes

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, \langle \lambda \rangle) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

For $1 \leq r \leq m$, we define the **truncated m -dimensional cube**

$$B_m(\leq r) := \{X \in \mathfrak{P}(m) \mid \text{either } \text{card } X \leq r \text{ or } X = m\},$$

endowed with containment.

Truncated m -cubes

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

For $1 \leq r \leq m$, we define the **truncated m -dimensional cube**

$$B_m(\leq r) := \{X \in \mathfrak{P}(m) \mid \text{either } \text{card } X \leq r \text{ or } X = m\},$$

endowed with containment.

Proposition (Gillibert + W., 2008)

Truncated m -cubes

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

For $1 \leq r \leq m$, we define the **truncated m -dimensional cube**

$$B_m(\leq r) := \{X \in \mathfrak{P}(m) \mid \text{either } \text{card } X \leq r \text{ or } X = m\},$$

endowed with containment.

Proposition (Gillibert + W., 2008)

Let r and m be integers with $1 \leq r < m$ and let κ and λ be infinite cardinals. Then $(\kappa, < \lambda) \rightsquigarrow B_m(\leq r)$ iff $(\kappa, r, \lambda) \rightarrow m$, that is, for each $F: [\kappa]^r \rightarrow [\kappa]^{< \lambda}$, there exists $H \in [\kappa]^m$ such that $F(X) \cap H \subseteq X$ for each $X \in [H]^r$ (**we say that H is free with respect to F**).

Kuratowski indexes of near-bottom truncated cubes

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Theorem

Kuratowski indexes of near-bottom truncated cubes

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Theorem

- (i) If GCH holds, then $(\lambda^{+r}, r, \lambda) \rightarrow \lambda^+$ for every infinite cardinal λ and every integer $r \geq 1$ (Erdős *et al.*, 1984).

Kuratowski indexes of near-bottom truncated cubes

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Theorem

- (i) If GCH holds, then $(\lambda^{+r}, r, \lambda) \rightarrow \lambda^+$ for every infinite cardinal λ and every integer $r \geq 1$ (Erdős *et al.*, 1984).
- (ii) $(\lambda^{+2}, 2, \lambda) \rightarrow m$ for every integer $m > 0$ (Hajnal and Máté, 1975). Hence $\text{kur } B_m(\leq 2) = 3$ for all $m \geq 3$.

Kuratowski indexes of near-bottom truncated cubes

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Theorem

- (i) If GCH holds, then $(\lambda^{+r}, r, \lambda) \rightarrow \lambda^+$ for every infinite cardinal λ and every integer $r \geq 1$ (Erdős *et al.*, 1984).
- (ii) $(\lambda^{+2}, 2, \lambda) \rightarrow m$ for every integer $m > 0$ (Hajnal and Máté, 1975). Hence $\text{kur } B_m(\leq 2) = 3$ for all $m \geq 3$.
- (iii) $(\lambda^{+3}, 3, \lambda) \rightarrow m$ for every integer $m > 0$ (Hajnal, 1984). Hence $\text{kur } B_m(\leq 3) = 4$ for all $m \geq 4$.

Kuratowski indexes of near-bottom truncated cubes

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Theorem

- (i) If GCH holds, then $(\lambda^{+r}, r, \lambda) \rightarrow \lambda^+$ for every infinite cardinal λ and every integer $r \geq 1$ (Erdős *et al.*, 1984).
- (ii) $(\lambda^{+2}, 2, \lambda) \rightarrow m$ for every integer $m > 0$ (Hajnal and Máté, 1975). Hence $\text{kur } B_m(\leq 2) = 3$ for all $m \geq 3$.
- (iii) $(\lambda^{+3}, 3, \lambda) \rightarrow m$ for every integer $m > 0$ (Hajnal, 1984). Hence $\text{kur } B_m(\leq 3) = 4$ for all $m \geq 4$.

Item (ii) above has been used by Ploščica and Gillibert to evaluate various critical points between finitely generated modular lattice varieties.

Higher truncated cubes

Set $t_0 := 5$, $t_1 := 7$, and, for each $n > 0$, let $t_{n+1} \rightarrow (t_n, 7)^5$
(they exist, due to Ramsey's Theorem).

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Higher truncated cubes

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Set $t_0 := 5$, $t_1 := 7$, and, for each $n > 0$, let $t_{n+1} \rightarrow (t_n, 7)^5$ (they exist, due to Ramsey's Theorem).

Theorem (Komjáth + Shelah, 2000)

Higher truncated cubes

Set $t_0 := 5$, $t_1 := 7$, and, for each $n > 0$, let $t_{n+1} \rightarrow (t_n, 7)^5$
(they exist, due to Ramsey's Theorem).

Theorem (Komjáth + Shelah, 2000)

For every $n > 0$, there exists a generic extension of the universe
in which $(\aleph_n, 4, \aleph_0) \not\rightarrow t_n$.

Lifters, free
sets, ladders

The Con_c
functor

k -ladders

Critical points

From lifting
objects to
lifting
diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating
Kuratowski
indexes of
finite posets

Higher truncated cubes

Set $t_0 := 5$, $t_1 := 7$, and, for each $n > 0$, let $t_{n+1} \rightarrow (t_n, 7)^5$ (they exist, due to Ramsey's Theorem).

Theorem (Komjáth + Shelah, 2000)

For every $n > 0$, there exists a generic extension of the universe in which $(\aleph_n, 4, \aleph_0) \not\rightarrow t_n$.

In particular,

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Higher truncated cubes

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Set $t_0 := 5$, $t_1 := 7$, and, for each $n > 0$, let $t_{n+1} \rightarrow (t_n, 7)^5$ (they exist, due to Ramsey's Theorem).

Theorem (Komjáth + Shelah, 2000)

For every $n > 0$, there exists a generic extension of the universe in which $(\aleph_n, 4, \aleph_0) \not\rightarrow t_n$.

In particular,

- There exists a generic extension of the universe in which $(\aleph_4, 4, \aleph_0) \not\rightarrow t_4$.

Higher truncated cubes

Set $t_0 := 5$, $t_1 := 7$, and, for each $n > 0$, let $t_{n+1} \rightarrow (t_n, 7)^5$ (they exist, due to Ramsey's Theorem).

Theorem (Komjáth + Shelah, 2000)

For every $n > 0$, there exists a generic extension of the universe in which $(\aleph_n, 4, \aleph_0) \not\rightarrow t_n$.

In particular,

- There exists a generic extension of the universe in which $(\aleph_4, 4, \aleph_0) \not\rightarrow t_4$.
- Hence $\text{kur } \mathbb{B}_{t_4}(\leq 4) = 5$ in any universe with GCH, while $\text{kur } \mathbb{B}_{t_4}(\leq 4) \geq 6$ in some generic extension (hence $P \mapsto \text{kur}(P)$ is not absolute).

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

A new lower bound (size of a free set)

It is not hard to verify that

breadth $B_{n+2}(\leq n) = \dim B_{n+2}(\leq n) = n + 1$, for all $n > 0$.

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

A new lower bound (size of a free set)

It is not hard to verify that

breadth $B_{n+2}(\leq n) = \dim B_{n+2}(\leq n) = n + 1$, for all $n > 0$.

Hence $\text{kur } B_{n+2}(\leq n) = n + 1$, and so

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

A new lower bound (size of a free set)

It is not hard to verify that

breadth $B_{n+2}(\leq n) = \dim B_{n+2}(\leq n) = n + 1$, for all $n > 0$.

Hence $\text{kur } B_{n+2}(\leq n) = n + 1$, and so

Theorem (Gillibert 2008)

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

A new lower bound (size of a free set)

It is not hard to verify that

breadth $B_{n+2}(\leq n) = \dim B_{n+2}(\leq n) = n + 1$, for all $n > 0$.

Hence $\text{kur } B_{n+2}(\leq n) = n + 1$, and so

Theorem (Gillibert 2008)

The relation $(\lambda^{+n}, n, \lambda) \rightarrow n + 2$ holds for each infinite cardinal λ and each positive integer n .

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

A new lower bound (size of a free set)

It is not hard to verify that

breadth $B_{n+2}(\leq n) = \dim B_{n+2}(\leq n) = n + 1$, for all $n > 0$.

Hence $\text{kur } B_{n+2}(\leq n) = n + 1$, and so

Theorem (Gillibert 2008)

The relation $(\aleph^{+n}, n, \aleph) \rightarrow n + 2$ holds for each infinite cardinal \aleph and each positive integer n .

- In particular, $(\aleph_4, 4, \aleph_0) \rightarrow 6$.

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

A new lower bound (size of a free set)

It is not hard to verify that
breadth $B_{n+2}(\leq n) = \dim B_{n+2}(\leq n) = n + 1$, for all $n > 0$.
Hence $\text{kur } B_{n+2}(\leq n) = n + 1$, and so

Theorem (Gillibert 2008)

The relation $(\aleph^{+n}, n, \lambda) \rightarrow n + 2$ holds for each infinite cardinal λ and each positive integer n .

- In particular, $(\aleph_4, 4, \aleph_0) \rightarrow 6$.
- **Previously known bound:** $(\aleph_4, 4, \aleph_0) \rightarrow 5$ (due to Kuratowski's Free Set Theorem).

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

A new lower bound (size of a free set)

It is not hard to verify that

breadth $B_{n+2}(\leq n) = \dim B_{n+2}(\leq n) = n + 1$, for all $n > 0$.

Hence $\text{kur } B_{n+2}(\leq n) = n + 1$, and so

Theorem (Gillibert 2008)

The relation $(\aleph^{+n}, n, \lambda) \rightarrow n + 2$ holds for each infinite cardinal λ and each positive integer n .

- In particular, $(\aleph_4, 4, \aleph_0) \rightarrow 6$.
- **Previously known bound:** $(\aleph_4, 4, \aleph_0) \rightarrow 5$ (due to Kuratowski's Free Set Theorem).
- **Next open question:** $(\aleph_4, 4, \aleph_0) \rightarrow 7$. That is, does $\text{kur } B_7(\leq 4) = 5$?

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

A new lower bound (size of a free set)

It is not hard to verify that

breadth $B_{n+2}(\leq n) = \dim B_{n+2}(\leq n) = n + 1$, for all $n > 0$.

Hence $\text{kur } B_{n+2}(\leq n) = n + 1$, and so

Theorem (Gillibert 2008)

The relation $(\lambda^{+n}, n, \lambda) \rightarrow n + 2$ holds for each infinite cardinal λ and each positive integer n .

- In particular, $(\aleph_4, 4, \aleph_0) \rightarrow 6$.
- **Previously known bound:** $(\aleph_4, 4, \aleph_0) \rightarrow 5$ (due to Kuratowski's Free Set Theorem).
- **Next open question:** $(\aleph_4, 4, \aleph_0) \rightarrow 7$. That is, does $\text{kur } B_7(\leq 4) = 5$?
- **The unreachable upper limit:** remember that $(\aleph_4, 4, \aleph_0) \not\rightarrow t_4$ in some generic extension.

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Using further order-dimension estimates

- Remember the estimate $\text{kur}(P) \leq \dim(P)$.

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Using further order-dimension estimates

- Remember the estimate $\text{kur}(P) \leq \dim(P)$.
- Estimating the order-dimension $\text{kur } B_m(\leq r)$ has been an object of intensive study (Dushnik, Füredi + Kahn, Kierstead, Hajnal, Spencer, and others).

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, \leq \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Using further order-dimension estimates

- Remember the estimate $\text{kur}(P) \leq \dim(P)$.
- Estimating the order-dimension $\text{kur } B_m(\leq r)$ has been an object of intensive study (Dushnik, Füredi + Kahn, Kierstead, Hajnal, Spencer, and others).
- Using the various estimates available, we obtain, for example:

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Using further order-dimension estimates

- Remember the estimate $\text{kur}(P) \leq \dim(P)$.
- Estimating the order-dimension $\text{kur} B_m(\leq r)$ has been an object of intensive study (Dushnik, Füredi + Kahn, Kierstead, Hajnal, Spencer, and others).
- Using the various estimates available, we obtain, for example:

Theorem (Gillibert + W., 2008)

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Using further order-dimension estimates

- Remember the estimate $\text{kur}(P) \leq \dim(P)$.
- Estimating the order-dimension $\text{kur} B_m(\leq r)$ has been an object of intensive study (Dushnik, Füredi + Kahn, Kierstead, Hajnal, Spencer, and others).
- Using the various estimates available, we obtain, for example:

Theorem (Gillibert + W., 2008)

- $(\aleph_7, 4, \aleph_0) \rightarrow 10$ (using Dushnik's estimate).

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Using further order-dimension estimates

- Remember the estimate $\text{kur}(P) \leq \dim(P)$.
- Estimating the order-dimension $\text{kur} B_m(\leq r)$ has been an object of intensive study (Dushnik, Füredi + Kahn, Kierstead, Hajnal, Spencer, and others).
- Using the various estimates available, we obtain, for example:

Theorem (Gillibert + W., 2008)

- $(\aleph_7, 4, \aleph_0) \rightarrow 10$ (using Dushnik's estimate).
- $(\aleph_9, 5, \aleph_0) \rightarrow 12$ (using Dushnik's estimate).

Lifters, free sets, ladders

The Conc functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Using further order-dimension estimates

- Remember the estimate $\text{kur}(P) \leq \dim(P)$.
- Estimating the order-dimension $\text{kur } B_m(\leq r)$ has been an object of intensive study (Dushnik, Füredi + Kahn, Kierstead, Hajnal, Spencer, and others).
- Using the various estimates available, we obtain, for example:

Theorem (Gillibert + W., 2008)

- $(\aleph_7, 4, \aleph_0) \rightarrow 10$ (using Dushnik's estimate).
- $(\aleph_9, 5, \aleph_0) \rightarrow 12$ (using Dushnik's estimate).
- $(\aleph_{109}, 4, \aleph_0) \rightarrow 257$ (using Füredi and Kahn's estimate).

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

Using further order-dimension estimates

- Remember the estimate $\text{kur}(P) \leq \dim(P)$.
- Estimating the order-dimension $\text{kur } B_m(\leq r)$ has been an object of intensive study (Dushnik, Füredi + Kahn, Kierstead, Hajnal, Spencer, and others).
- Using the various estimates available, we obtain, for example:

Theorem (Gillibert + W., 2008)

- $(\aleph_7, 4, \aleph_0) \rightarrow 10$ (using Dushnik's estimate).
- $(\aleph_9, 5, \aleph_0) \rightarrow 12$ (using Dushnik's estimate).
- $(\aleph_{109}, 4, \aleph_0) \rightarrow 257$ (using Füredi and Kahn's estimate).
- $(\aleph_{210}, 4, \aleph_0) \rightarrow 32,768$ (using Hajnal and Spencer's estimate).

Lifters, free sets, ladders

The Con_c functor

k -ladders

Critical points

From lifting objects to lifting diagrams

λ -lifters

$(\kappa, < \lambda) \rightsquigarrow P$

Calculating Kuratowski indexes of finite posets

For general n and r , $(\aleph_n, r, \aleph_0) \rightarrow E(n, r)$ with

$$\lg \lg E(n, r) \sim \frac{n}{r 2^r \log 2} \text{ as } n \gg r \gg 0$$

(again by using Hajnal and Spencer's estimate).