

Coproducts and colimits of κ -quantales

R. N. Ball and A. Pultr

BLAST

6 June 2010

Motivation

Motivation

Frame quotients are relatively easy to compute in many cases for two reasons.

Motivation

Frame quotients are relatively easy to compute in many cases for two reasons.

- Each equivalence class (fiber of the quotient map) has a greatest element.

Frame quotients are relatively easy to compute in many cases for two reasons.

- Each equivalence class (fiber of the quotient map) has a greatest element.
- The frame law asserts that the meet operation preserves all joins, and therefore has an adjoint map, which is the Heyting arrow.

Frame quotients are relatively easy to compute in many cases for two reasons.

- Each equivalence class (fiber of the quotient map) has a greatest element.
- The frame law asserts that the meet operation preserves all joins, and therefore has an adjoint map, which is the Heyting arrow. Although this operation is not part of the formal frame apparatus, it provides a simple technique for extending a generating relation to a frame congruence.

Frame quotients are relatively easy to compute in many cases for two reasons.

- Each equivalence class (fiber of the quotient map) has a greatest element.
- The frame law asserts that the meet operation preserves all joins, and therefore has an adjoint map, which is the Heyting arrow. Although this operation is not part of the formal frame apparatus, it provides a simple technique for extending a generating relation to a frame congruence.

We will extend the second point to κ -quantales, substituting the monoid multiplication for the meet operation.

Frame quotients are relatively easy to compute in many cases for two reasons.

- Each equivalence class (fiber of the quotient map) has a greatest element.
- The frame law asserts that the meet operation preserves all joins, and therefore has an adjoint map, which is the Heyting arrow. Although this operation is not part of the formal frame apparatus, it provides a simple technique for extending a generating relation to a frame congruence.

We will extend the second point to κ -quantales, substituting the monoid multiplication for the meet operation. And we will use the first point on suitable extensions to give information about quotients of κ -quantales.

The symbols λ and κ will represent infinite regular cardinals,

The symbols λ and κ will represent infinite regular cardinals, or the symbol 0 ,

The symbols λ and κ will represent infinite regular cardinals, or the symbol 0 , or the symbol ∞ .

The symbols λ and κ will represent infinite regular cardinals, or the symbol 0 , or the symbol ∞ . The order is

$$0 \leq \lambda \leq \kappa \leq \infty$$

The symbols λ and κ will represent infinite regular cardinals, or the symbol 0 , or the symbol ∞ . The order is

$$0 \leq \lambda \leq \kappa \leq \infty$$

Definition

A κ -set is a set of cardinality strictly less than κ .

The symbols λ and κ will represent infinite regular cardinals, or the symbol 0 , or the symbol ∞ . The order is

$$0 \leq \lambda \leq \kappa \leq \infty$$

Definition

A κ -set is a set of cardinality strictly less than κ . There are no 0 -sets,

The symbols λ and κ will represent infinite regular cardinals, or the symbol 0 , or the symbol ∞ . The order is

$$0 \leq \lambda \leq \kappa \leq \infty$$

Definition

A κ -set is a set of cardinality strictly less than κ . There are no 0 -sets, an ω -set is a finite set,

The symbols λ and κ will represent infinite regular cardinals, or the symbol 0 , or the symbol ∞ . The order is

$$0 \leq \lambda \leq \kappa \leq \infty$$

Definition

A κ -set is a set of cardinality strictly less than κ . There are no 0 -sets, an ω -set is a finite set, an ω_1 -set is a countable set,

The symbols λ and κ will represent infinite regular cardinals, or the symbol 0 , or the symbol ∞ . The order is

$$0 \leq \lambda \leq \kappa \leq \infty$$

Definition

A κ -set is a set of cardinality strictly less than κ . There are no 0 -sets, an ω -set is a finite set, an ω_1 -set is a countable set, and an ∞ -set is just a set.

The symbols λ and κ will represent infinite regular cardinals, or the symbol 0 , or the symbol ∞ . The order is

$$0 \leq \lambda \leq \kappa \leq \infty$$

Definition

A κ -set is a set of cardinality strictly less than κ . There are no 0 -sets, an ω -set is a finite set, an ω_1 -set is a countable set, and an ∞ -set is just a set. A κ -subset is a subset $A \subseteq B$ which is a κ -set; we write $A \subseteq_{\kappa} B$.

The symbols λ and κ will represent infinite regular cardinals, or the symbol 0 , or the symbol ∞ . The order is

$$0 \leq \lambda \leq \kappa \leq \infty$$

Definition

A κ -set is a set of cardinality strictly less than κ . There are no 0 -sets, an ω -set is a finite set, an ω_1 -set is a countable set, and an ∞ -set is just a set. A κ -subset is a subset $A \subseteq B$ which is a κ -set; we write $A \subseteq_{\kappa} B$. Likewise for κ -union, κ -join, etc.

For the purposes of this talk,

Definition

A κ -quantale is a commutative monoid and a κ -complete lattice

For the purposes of this talk,

Definition

A κ -quantale is a commutative monoid and a κ -complete lattice such that the identity element of the monoid is the top element of the lattice,

For the purposes of this talk,

Definition

A κ -quantale is a commutative monoid and a κ -complete lattice such that the identity element of the monoid is the top element of the lattice, and the monoid multiplication distributes across κ -joins.

For the purposes of this talk,

Definition

A κ -quantale is a commutative monoid and a κ -complete lattice such that the identity element of the monoid is the top element of the lattice, and the monoid multiplication distributes across κ -joins. κ -morphisms preserve all relevant structure. The category is $\kappa\mathbf{Qnt}$.

For the purposes of this talk,

Definition

A κ -quantale is a commutative monoid and a κ -complete lattice such that the identity element of the monoid is the top element of the lattice, and the monoid multiplication distributes across κ -joins. κ -morphisms preserve all relevant structure. The category is $\kappa\mathbf{Qnt}$. 0-quantales have no order, and are therefore simply commutative monoids. We write $\mathbf{0Qnt}$ as \mathbf{CMon} .

For the purposes of this talk,

Definition

A κ -quantale is a commutative monoid and a κ -complete lattice such that the identity element of the monoid is the top element of the lattice, and the monoid multiplication distributes across κ -joins. κ -morphisms preserve all relevant structure. The category is $\kappa\mathbf{Qnt}$. 0-quantales have no order, and are therefore simply commutative monoids. We write $\mathbf{0Qnt}$ as \mathbf{CMon} . We refer to ∞ -quantales as simply quantales, and write $\infty\mathbf{Qnt}$ as \mathbf{Qnt} .

For the purposes of this talk,

Definition

A κ -quantale is a commutative monoid and a κ -complete lattice such that the identity element of the monoid is the top element of the lattice, and the monoid multiplication distributes across κ -joins. κ -morphisms preserve all relevant structure. The category is $\kappa\mathbf{Qnt}$. 0-quantales have no order, and are therefore simply commutative monoids. We write $\mathbf{0Qnt}$ as \mathbf{CMon} . We refer to ∞ -quantales as simply quantales, and write $\infty\mathbf{Qnt}$ as \mathbf{Qnt} .

- By distributivity, $x \cdot y$ is monotone in both variables.

For the purposes of this talk,

Definition

A κ -quantale is a commutative monoid and a κ -complete lattice such that the identity element of the monoid is the top element of the lattice, and the monoid multiplication distributes across κ -joins. κ -morphisms preserve all relevant structure. The category is $\kappa\mathbf{Qnt}$. 0-quantales have no order, and are therefore simply commutative monoids. We write $\mathbf{0Qnt}$ as \mathbf{CMon} . We refer to ∞ -quantales as simply quantales, and write $\infty\mathbf{Qnt}$ as \mathbf{Qnt} .

- By distributivity, $x \cdot y$ is monotone in both variables.
- $xy \leq x, y$, since 1 is the top, and $x \cdot 0 = 0$ since $x \cdot 0 \leq 1 \cdot 0 = 0$.

For the purposes of this talk,

Definition

A κ -quantale is a commutative monoid and a κ -complete lattice such that the identity element of the monoid is the top element of the lattice, and the monoid multiplication distributes across κ -joins. κ -morphisms preserve all relevant structure. The category is $\kappa\mathbf{Qnt}$. 0-quantales have no order, and are therefore simply commutative monoids. We write $\mathbf{0Qnt}$ as \mathbf{CMon} . We refer to ∞ -quantales as simply quantales, and write $\infty\mathbf{Qnt}$ as \mathbf{Qnt} .

- By distributivity, $x \cdot y$ is monotone in both variables.
- $xy \leq x, y$, since 1 is the top, and $x \cdot 0 = 0$ since $x \cdot 0 \leq 1 \cdot 0 = 0$.
- $xy = x \wedge y$ iff the monoid is idempotent, since in that case $z \leq x, y$ implies $z = z \cdot z \leq x \cdot y$.

For the purposes of this talk,

Definition

A κ -quantale is a commutative monoid and a κ -complete lattice such that the identity element of the monoid is the top element of the lattice, and the monoid multiplication distributes across κ -joins. κ -morphisms preserve all relevant structure. The category is $\kappa\mathbf{Qnt}$. 0-quantales have no order, and are therefore simply commutative monoids. We write $\mathbf{0Qnt}$ as \mathbf{CMon} . We refer to ∞ -quantales as simply quantales, and write $\infty\mathbf{Qnt}$ as \mathbf{Qnt} .

- By distributivity, $x \cdot y$ is monotone in both variables.
- $xy \leq x, y$, since 1 is the top, and $x \cdot 0 = 0$ since $x \cdot 0 \leq 1 \cdot 0 = 0$.
- $xy = x \wedge y$ iff the monoid is idempotent, since in that case $z \leq x, y$ implies $z = z \cdot z \leq x \cdot y$.
- The ideals of a commutative ring with unit form a quantale.

The first tool: quantale quotients

The first tool: quantale quotients

Let R be a binary relation on a quantale L .

The first tool: quantale quotients

Let R be a binary relation on a quantale L .

Definition

An element $s \in S$ is said to be **R -saturated**, or simply **saturated**, if

$$\forall a, b, c \quad aRb \implies (ac \leq s \text{ iff } bc \leq s).$$

The set of all saturated elements will be denoted by L/R .

The first tool: quantale quotients

Let R be a binary relation on a quantale L .

Definition

An element $s \in S$ is said to be **R -saturated**, or simply **saturated**, if

$$\forall a, b, c \quad aRb \implies (ac \leq s \text{ iff } bc \leq s).$$

The set of all saturated elements will be denoted by L/R .

The distributivity $a \cdot \bigvee b_i = \bigvee (a \cdot b_i)$ in L can be interpreted as saying that the mappings $(x \mapsto a \cdot x) : L \rightarrow L$ preserve all suprema, and hence they are left Galois adjoints.

The first tool: quantale quotients

Let R be a binary relation on a quantale L .

Definition

An element $s \in S$ is said to be **R -saturated**, or simply **saturated**, if

$$\forall a, b, c \quad aRb \implies (ac \leq s \text{ iff } bc \leq s).$$

The set of all saturated elements will be denoted by L/R .

The distributivity $a \cdot \bigvee b_i = \bigvee (a \cdot b_i)$ in L can be interpreted as saying that the mappings $(x \mapsto a \cdot x) : L \rightarrow L$ preserve all suprema, and hence they are left Galois adjoints. This gives rise to an operation \rightarrow on L such that

$$ab \leq c \quad \text{iff} \quad a \leq b \rightarrow c.$$

R -saturated elements

Definition

An element $s \in S$ is said to be R -saturated if

$$\forall a, b, c \quad aRb \implies (ac \leq s \text{ iff } bc \leq s).$$

Observation

- *The meet of an arbitrary set of saturated elements is saturated.*

R -saturated elements

Definition

An element $s \in S$ is said to be R -saturated if

$$\forall a, b, c \quad aRb \implies (ac \leq s \text{ iff } bc \leq s).$$

Observation

- *The meet of an arbitrary set of saturated elements is saturated.*
- *If s is saturated then so is $x \rightarrow s$ for any x .*

R -saturated elements

Definition

An element $s \in S$ is said to be R -saturated if

$$\forall a, b, c \quad aRb \implies (ac \leq s \text{ iff } bc \leq s).$$

Observation

- *The meet of an arbitrary set of saturated elements is saturated.*
- *If s is saturated then so is $x \rightarrow s$ for any x .*

Proof.

Suppose aRb . Then

$$ac \leq x \rightarrow s \iff acx \leq s \iff bcx \leq s \iff bc \leq x \rightarrow s$$



The quotient map

Definition

Define a mapping $\mu_R \equiv (x \mapsto \bigwedge_{x \leq s \in L/R} s) : L \longrightarrow L/R$.

The quotient map

Definition

Define a mapping $\mu_R \equiv (x \mapsto \bigwedge_{x \leq s \in L/R} s) : L \longrightarrow L/R$.

Lemma

- 1 $x \leq \mu(x)$, μ is monotone, and $\mu\mu(x) = \mu(x)$,
- 2 $\mu(xy) = \mu(\mu(x)\mu(y))$.

The quotient map

Definition

Define a mapping $\mu_R \equiv (x \mapsto \bigwedge_{x \leq s \in L/R} s) : L \longrightarrow L/R$.

Lemma

- 1 $x \leq \mu(x)$, μ is monotone, and $\mu\mu(x) = \mu(x)$,
- 2 $\mu(xy) = \mu(\mu(x)\mu(y))$.

Theorem

*L/R is a complete lattice, and if it is endowed with the multiplication $x * y = \mu(xy)$ it becomes a quantale and μ_R becomes a quantale morphism $L \rightarrow L/R$.*

The quotient map

Definition

Define a mapping $\mu_R \equiv (x \mapsto \bigwedge_{x \leq s \in L/R} s) : L \longrightarrow L/R$.

Lemma

- 1 $x \leq \mu(x)$, μ is monotone, and $\mu\mu(x) = \mu(x)$,
- 2 $\mu(xy) = \mu(\mu(x)\mu(y))$.

Theorem

*L/R is a complete lattice, and if it is endowed with the multiplication $x * y = \mu(xy)$ it becomes a quantale and μ_R becomes a quantale morphism $L \rightarrow L/R$.*

If aRb then $\mu_R(a) = \mu_R(b)$, and for every κ -morphism $h : L \rightarrow M$ such that $aRb \Rightarrow h(a) = h(b)$ there is a unique quantale morphism $\bar{h} : L/R \rightarrow M$ such that $\bar{h}\mu_R = h$. Moreover, $\bar{h}(a) = h(a)$ for all $a \in L/R$.

The second tool: the free κ -quantale over a λ -quantale

The second tool: the free κ -quantale over a λ -quantale

The forgetful functor $\kappa\mathbf{Qnt} \rightarrow \lambda\mathbf{Qnt}$ has a left adjoint, which we label $\mathfrak{F}_\lambda^\kappa$.

The second tool: the free κ -quantale over a λ -quantale

The forgetful functor $\kappa\mathbf{Qnt} \rightarrow \lambda\mathbf{Qnt}$ has a left adjoint, which we label $\mathfrak{F}_\lambda^\kappa$. We begin by describing $\mathfrak{F}_\kappa^0 S$, the free κ -quantale over a commutative monoid S .

The second tool: the free κ -quantale over a λ -quantale

The forgetful functor $\kappa\mathbf{Qnt} \rightarrow \lambda\mathbf{Qnt}$ has a left adjoint, which we label $\mathfrak{F}_\lambda^\kappa$. We begin by describing $\mathfrak{F}_\kappa^0 S$, the free κ -quantale over a commutative monoid S . Fix $\kappa > 0$.

The second tool: the free κ -quantale over a λ -quantale

The forgetful functor $\kappa\mathbf{Qnt} \rightarrow \lambda\mathbf{Qnt}$ has a left adjoint, which we label $\mathfrak{F}_\lambda^\kappa$. We begin by describing $\mathfrak{F}_\kappa^0 S$, the free κ -quantale over a commutative monoid S . Fix $\kappa > 0$.

Definition

A **pre-ideal** in a commutative monoid S is a subset $U \subseteq S$ such that

$$u \in U \ \& \ s \in S \quad \implies \quad us \in U.$$

The second tool: the free κ -quantale over a λ -quantale

The forgetful functor $\kappa\mathbf{Qnt} \rightarrow \lambda\mathbf{Qnt}$ has a left adjoint, which we label $\mathfrak{F}_\lambda^\kappa$. We begin by describing $\mathfrak{F}_\kappa^0 S$, the free κ -quantale over a commutative monoid S . Fix $\kappa > 0$.

Definition

A **pre-ideal** in a commutative monoid S is a subset $U \subseteq S$ such that

$$u \in U \ \& \ s \in S \quad \implies \quad us \in U.$$

The pre-ideal generated by an arbitrary subset $A \subseteq S$ is

$$[A] \equiv \{as : a \in A, s \in S\} = \bigcup_A [a].$$

The second tool: the free κ -quantale over a λ -quantale

The forgetful functor $\kappa\mathbf{Qnt} \rightarrow \lambda\mathbf{Qnt}$ has a left adjoint, which we label $\mathfrak{F}_\lambda^\kappa$. We begin by describing $\mathfrak{F}_\kappa^0 S$, the free κ -quantale over a commutative monoid S . Fix $\kappa > 0$.

Definition

A **pre-ideal** in a commutative monoid S is a subset $U \subseteq S$ such that

$$u \in U \ \& \ s \in S \quad \implies \quad us \in U.$$

The pre-ideal generated by an arbitrary subset $A \subseteq S$ is

$$[A] \equiv \{as : a \in A, s \in S\} = \bigcup_A [a].$$

The smallest pre-ideal containing an element $a \in S$ is the principal pre-ideal

$$[a] = \{as : s \in S\}.$$

Properties of pre-ideals

Though a pre-ideal need not be a downset, a downset is a pre-ideal in any quantale,

Properties of pre-ideals

Though a pre-ideal need not be a downset, a downset is a pre-ideal in any quantale, and the pre-ideals of a meet-semilattice are exactly the downsets. In that case $[a] = \downarrow a$.

Properties of pre-ideals

Though a pre-ideal need not be a downset, a downset is a pre-ideal in any quantale, and the pre-ideals of a meet-semilattice are exactly the downsets. In that case $[a] = \downarrow a$.

Lemma

Let S be a commutative monoid.

- 1 If U_i , $i \in I$, are pre-ideals then so is $\bigcup_i U_i$.
- 2 If U and V are pre-ideals then $U \cdot V = \{uv : u \in U, v \in V\}$ is a pre-ideal. This operation is associative and commutative. If the monoid is idempotent, i.e., a meet semilattice, then $U \cdot U = U$.
- 3 $U \cdot S = U$.
- 4 $U \cdot (\bigcup_i V_i) = \bigcup_i (U \cdot V_i)$.
- 5 $[a] \cdot [b] = [ab]$, and $[1] = S$.

Properties of pre-ideals

Though a pre-ideal need not be a downset, a downset is a pre-ideal in any quantale, and the pre-ideals of a meet-semilattice are exactly the downsets. In that case $[a] = \downarrow a$.

Lemma

Let S be a commutative monoid.

- 1 If U_i , $i \in I$, are pre-ideals then so is $\bigcup_i U_i$.
- 2 If U and V are pre-ideals then $U \cdot V = \{uv : u \in U, v \in V\}$ is a pre-ideal. This operation is associative and commutative. If the monoid is idempotent, i.e., a meet semilattice, then $U \cdot U = U$.
- 3 $U \cdot S = U$.
- 4 $U \cdot (\bigcup_i V_i) = \bigcup_i (U \cdot V_i)$.
- 5 $[a] \cdot [b] = [ab]$, and $[1] = S$.

The content of this lemma is that the family of pre-ideals on S forms a quantale in the inclusion order.

Properties of pre-ideals

Though a pre-ideal need not be a downset, a downset is a pre-ideal in any quantale, and the pre-ideals of a meet-semilattice are exactly the downsets. In that case $[a] = \downarrow a$.

Lemma

Let S be a commutative monoid.

- 1 If U_i , $i \in I$, are pre-ideals then so is $\bigcup_i U_i$.
- 2 If U and V are pre-ideals then $U \cdot V = \{uv : u \in U, v \in V\}$ is a pre-ideal. This operation is associative and commutative. If the monoid is idempotent, i.e., a meet semilattice, then $U \cdot U = U$.
- 3 $U \cdot S = U$.
- 4 $U \cdot (\bigcup_i V_i) = \bigcup_i (U \cdot V_i)$.
- 5 $[a] \cdot [b] = [ab]$, and $[1] = S$.

The content of this lemma is that the family of pre-ideals on S forms a quantale in the inclusion order. Note: order from no order.

The free κ -quantale over a commutative monoid

The free κ -quantale over a commutative monoid

Definition

For a commutative monoid S let

$$\mathfrak{F}_\kappa^0 S \equiv \{[A] : A \subseteq_\kappa S\} \cdot,$$

The free κ -quantale over a commutative monoid

Definition

For a commutative monoid S let

$$\mathfrak{F}_\kappa^0 S \equiv \{[A] : A \subseteq_\kappa S\} .,$$

Define the mapping

$$\rho_{\kappa S}^0 : S \rightarrow \mathfrak{F}_\kappa^0 S \equiv (a \mapsto [a], a \in S).$$

Abbreviate $\rho_{\infty S}^0$ to ρ_S^0 .

The free κ -quantale over a commutative monoid

Theorem

$\rho_{\kappa S}^0 : S \rightarrow \mathfrak{F}_{\kappa}^0 S$ is the free κ -quantale over the commutative monoid S . That is, for every κ -quantale L and monoid homomorphism $h : S \rightarrow L$ there is precisely one κ -morphism $f : \mathfrak{F}_{\kappa}^0 S \rightarrow L$ such that the diagram

$$\begin{array}{ccc} \mathfrak{F}_{\kappa}^0 S & \xrightarrow{f} & L \\ \rho_{\kappa S}^0 \uparrow & \nearrow h & \\ S & & \end{array}$$

commutes.

The free κ -quantale over a λ -quantale, $\lambda > 0$

The free κ -quantale over a λ -quantale, $\lambda > 0$

We can expect to construct this object by

The free κ -quantale over a λ -quantale, $\lambda > 0$

We can expect to construct this object by

- first forgetting the order on L , i.e., applying the forgetful functor to **CMon**,

The free κ -quantale over a λ -quantale, $\lambda > 0$

We can expect to construct this object by

- first forgetting the order on L , i.e., applying the forgetful functor to **CMon**,
- then passing to $\mathfrak{F}_\kappa^0 L$, the free κ -quantale over the monoid L ,

The free κ -quantale over a λ -quantale, $\lambda > 0$

We can expect to construct this object by

- first forgetting the order on L , i.e., applying the forgetful functor to **CMon**,
- then passing to $\mathfrak{F}_\kappa^0 L$, the free κ -quantale over the monoid L ,
- and then factoring $\mathfrak{F}_\kappa^0 L$ by the relation R which forces the insertion of L into $\mathfrak{F}_\kappa^0 L$ to respect κ -joins.

The free κ -quantale over a λ -quantale, $\lambda > 0$

We can expect to construct this object by

- first forgetting the order on L , i.e., applying the forgetful functor to **CMon**,
- then passing to $\mathfrak{F}_\kappa^0 L$, the free κ -quantale over the monoid L ,
- and then factoring $\mathfrak{F}_\kappa^0 L$ by the relation R which forces the insertion of L into $\mathfrak{F}_\kappa^0 L$ to respect κ -joins.

Definition

A **λ -ideal** in a λ -quantale L is a downset $U \subseteq L$ such that $\forall A \in U$ for all $A \subseteq_\lambda U$.

The free κ -quantale over a λ -quantale, $\lambda > 0$

We can expect to construct this object by

- first forgetting the order on L , i.e., applying the forgetful functor to **CMon**,
- then passing to $\mathfrak{F}_\kappa^0 L$, the free κ -quantale over the monoid L ,
- and then factoring $\mathfrak{F}_\kappa^0 L$ by the relation R which forces the insertion of L into $\mathfrak{F}_\kappa^0 L$ to respect κ -joins.

Definition

A λ -ideal in a λ -quantale L is a downset $U \subseteq L$ such that $\bigvee A \in U$ for all $A \subseteq_\lambda U$. Denote the smallest λ -ideal containing $A \subseteq L$ by

$$\langle A \rangle_\lambda \equiv \downarrow \left\{ \bigvee B : B \subseteq_\lambda A \right\}$$

Let

$$\mathfrak{F}_\kappa^\lambda L = \{ \langle A \rangle_\lambda : A \subseteq_k L \},$$

The free κ -quantale over a λ -quantale, $\lambda > 0$

We can expect to construct this object by

- first forgetting the order on L , i.e., applying the forgetful functor to **CMon**,
- then passing to $\mathfrak{F}_\kappa^0 L$, the free κ -quantale over the monoid L ,
- and then factoring $\mathfrak{F}_\kappa^0 L$ by the relation R which forces the insertion of L into $\mathfrak{F}_\kappa^0 L$ to respect κ -joins.

Definition

A λ -ideal in a λ -quantale L is a downset $U \subseteq L$ such that $\bigvee A \in U$ for all $A \subseteq_\lambda U$. Denote the smallest λ -ideal containing $A \subseteq L$ by

$$\langle A \rangle_\lambda \equiv \downarrow \left\{ \bigvee B : B \subseteq_\lambda A \right\}$$

Let

$$\mathfrak{F}_\kappa^\lambda L = \{ \langle A \rangle_\lambda : A \subseteq_k L \},$$

and let $\rho_{\kappa L}^\lambda(a) : L \rightarrow \mathfrak{F}_\kappa^\lambda L \equiv (a \mapsto \downarrow a, a \in L)$.

The free κ -quantale over a λ -quantale, $\lambda > 0$

We can expect to construct this object by

- first forgetting the order on L , i.e., applying the forgetful functor to **CMon**,
- then passing to $\mathfrak{F}_\kappa^0 L$, the free κ -quantale over the monoid L ,
- and then factoring $\mathfrak{F}_\kappa^0 L$ by the relation R which forces the insertion of L into $\mathfrak{F}_\kappa^0 L$ to respect κ -joins.

Definition

A λ -ideal in a λ -quantale L is a downset $U \subseteq L$ such that $\bigvee A \in U$ for all $A \subseteq_\lambda U$. Denote the smallest λ -ideal containing $A \subseteq L$ by

$$\langle A \rangle_\lambda \equiv \downarrow \left\{ \bigvee B : B \subseteq_\lambda A \right\}$$

Let

$$\mathfrak{F}_\kappa^\lambda L = \{ \langle A \rangle_\lambda : A \subseteq_k L \},$$

and let $\rho_{\kappa L}^\lambda(a) : L \rightarrow \mathfrak{F}_\kappa^\lambda L \equiv (a \mapsto \downarrow a, a \in L)$. Abbreviate $\rho_{\infty L}^\lambda$ to ρ_L^λ .

The κ -free quantale over a λ -quantale, $\lambda > 0$

Theorem

Let L be a λ -quantale. Then $\mathfrak{F}_\kappa^\lambda L$ is a quantale with respect to the operations

$$U \cdot V = \downarrow \{uv : u \in U, v \in V\},$$

$$\bigvee_I V_i = \downarrow \left\{ \bigvee A : A \subseteq_\lambda \bigcup_I V_i \right\}$$

In fact, $\rho_{\kappa L}^\lambda(a) : L \rightarrow \mathfrak{F}_L^\lambda$ is the free κ -quantale over L .

λ -coherent κ -quantales

λ -coherent κ -quantales

It is a surprising fact that those κ -quantales L which are free over some λ -quantale M can be intrinsically characterized.

λ -coherent κ -quantales

It is a surprising fact that those κ -quantales L which are free over some λ -quantale M can be intrinsically characterized. Even more surprisingly, M can be recovered from L .

It is a surprising fact that those κ -quantales L which are free over some λ -quantale M can be intrinsically characterized. Even more surprisingly, M can be recovered from L .

Definition

Let L be a κ -quantale. An element $a \in L$ is called a **λ -element** if for all $A \subseteq_{\kappa} L$ such that $\bigvee A \geq a$ there is some $A_0 \subseteq_{\lambda} A$ such that $\bigvee A_0 \geq a$.

It is a surprising fact that those κ -quantales L which are free over some λ -quantale M can be intrinsically characterized. Even more surprisingly, M can be recovered from L .

Definition

Let L be a κ -quantale. An element $a \in L$ is called a **λ -element** if for all $A \subseteq_{\kappa} L$ such that $\bigvee A \geq a$ there is some $A_0 \subseteq_{\lambda} A$ such that $\bigvee A_0 \geq a$. The set of λ -elements of L is designated $\mathfrak{E}_{\lambda}^{\kappa} L$.

It is a surprising fact that those κ -quantales L which are free over some λ -quantale M can be intrinsically characterized. Even more surprisingly, M can be recovered from L .

Definition

Let L be a κ -quantale. An element $a \in L$ is called a **λ -element** if for all $A \subseteq_{\kappa} L$ such that $\bigvee A \geq a$ there is some $A_0 \subseteq_{\lambda} A$ such that $\bigvee A_0 \geq a$. The set of λ -elements of L is designated $\mathfrak{E}_{\lambda}^{\kappa} L$. This set is evidently closed under λ -joins,

It is a surprising fact that those κ -quantales L which are free over some λ -quantale M can be intrinsically characterized. Even more surprisingly, M can be recovered from L .

Definition

Let L be a κ -quantale. An element $a \in L$ is called a **λ -element** if for all $A \subseteq_{\kappa} L$ such that $\bigvee A \geq a$ there is some $A_0 \subseteq_{\lambda} A$ such that $\bigvee A_0 \geq a$. The set of λ -elements of L is designated $\mathfrak{E}_{\lambda}^{\kappa} L$. This set is evidently closed under λ -joins, and we call L **λ -coherent** if $\mathfrak{E}_{\lambda}^{\kappa} L$ forms a generating sub- λ -frame of L .

It is a surprising fact that those κ -quantales L which are free over some λ -quantale M can be intrinsically characterized. Even more surprisingly, M can be recovered from L .

Definition

Let L be a κ -quantale. An element $a \in L$ is called a **λ -element** if for all $A \subseteq_{\kappa} L$ such that $\bigvee A \geq a$ there is some $A_0 \subseteq_{\lambda} A$ such that $\bigvee A_0 \geq a$. The set of λ -elements of L is designated $\mathfrak{E}_{\lambda}^{\kappa} L$. This set is evidently closed under λ -joins, and we call L **λ -coherent** if $\mathfrak{E}_{\lambda}^{\kappa} L$ forms a generating sub- λ -frame of L . More explicitly, L is λ -coherent if

- every element of L is a supremum of a κ -set of λ -elements,

It is a surprising fact that those κ -quantales L which are free over some λ -quantale M can be intrinsically characterized. Even more surprisingly, M can be recovered from L .

Definition

Let L be a κ -quantale. An element $a \in L$ is called a **λ -element** if for all $A \subseteq_{\kappa} L$ such that $\bigvee A \geq a$ there is some $A_0 \subseteq_{\lambda} A$ such that $\bigvee A_0 \geq a$. The set of λ -elements of L is designated $\mathfrak{E}_{\lambda}^{\kappa} L$. This set is evidently closed under λ -joins, and we call L **λ -coherent** if $\mathfrak{E}_{\lambda}^{\kappa} L$ forms a generating sub- λ -frame of L . More explicitly, L is λ -coherent if

- every element of L is a supremum of a κ -set of λ -elements,
- the product of finitely many λ -elements is a λ -element,

It is a surprising fact that those κ -quantales L which are free over some λ -quantale M can be intrinsically characterized. Even more surprisingly, M can be recovered from L .

Definition

Let L be a κ -quantale. An element $a \in L$ is called a **λ -element** if for all $A \subseteq_{\kappa} L$ such that $\bigvee A \geq a$ there is some $A_0 \subseteq_{\lambda} A$ such that $\bigvee A_0 \geq a$. The set of λ -elements of L is designated $\mathfrak{E}_{\lambda}^{\kappa} L$. This set is evidently closed under λ -joins, and we call L **λ -coherent** if $\mathfrak{E}_{\lambda}^{\kappa} L$ forms a generating sub- λ -frame of L . More explicitly, L is λ -coherent if

- every element of L is a supremum of a κ -set of λ -elements,
- the product of finitely many λ -elements is a λ -element,
- and 1 is a λ -element.

Theorem

A κ -quantale is λ -free iff it is λ -coherent.

Theorem

A κ -quantale is λ -free iff it is λ -coherent. More precisely, we have the following.

- 1 For any λ -quantale L , $\mathfrak{F}_\kappa^\lambda L$ is λ -coherent and

$$\mathfrak{E}_\lambda^\kappa \mathfrak{F}_\kappa^\lambda L = \{\downarrow a : a \in L\}.$$

Theorem

A κ -quantale is λ -free iff it is λ -coherent. More precisely, we have the following.

- 1 For any λ -quantale L , $\mathfrak{F}_\kappa^\lambda L$ is λ -coherent and

$$\mathfrak{E}_\lambda^\kappa \mathfrak{F}_\kappa^\lambda L = \{\downarrow a : a \in L\}.$$

- 2 For any λ -coherent κ -frame L , the inclusion $\mathfrak{E}_\lambda^\kappa L \rightarrow L$ lifts to an isomorphism $\mathfrak{F}_\kappa^\lambda \mathfrak{E}_\lambda^\kappa L \rightarrow L$.

Theorem

A κ -quantale is λ -free iff it is λ -coherent. More precisely, we have the following.

- 1 For any λ -quantale L , $\mathfrak{F}_\kappa^\lambda L$ is λ -coherent and

$$\mathfrak{E}_\lambda^\kappa \mathfrak{F}_\kappa^\lambda L = \{\downarrow a : a \in L\}.$$

- 2 For any λ -coherent κ -frame L , the inclusion $\mathfrak{E}_\lambda^\kappa L \rightarrow L$ lifts to an isomorphism $\mathfrak{F}_\kappa^\lambda \mathfrak{E}_\lambda^\kappa L \rightarrow L$.

This result directly generalizes to κ -quantales Madden's corresponding result for κ -frames.

κ -quantale quotients

κ -quantale quotients

We can now handle quotients of κ -quantales by a simple application of the functor \mathfrak{F}^κ .

κ -quantale quotients

We can now handle quotients of κ -quantales by a simple application of the functor \mathfrak{F}^κ . Given a binary relation R on a κ -quantale L ,

κ -quantale quotients

We can now handle quotients of κ -quantales by a simple application of the functor \mathfrak{F}^κ . Given a binary relation R on a κ -quantale L ,

- first pass to $\mathfrak{F}^\kappa L$, the free quantale over L ,

κ -quantale quotients

We can now handle quotients of κ -quantales by a simple application of the functor \mathfrak{F}^κ . Given a binary relation R on a κ -quantale L ,

- first pass to $\mathfrak{F}^\kappa L$, the free quantale over L ,
- factor $\mathfrak{F}^\kappa L$ by the relation $\tilde{R} \equiv \{(\downarrow a, \downarrow b) : aRb\}$, getting quotient map $\mu : \mathfrak{F}^\kappa L \rightarrow \mathfrak{F}^\kappa L / \tilde{R}$,

We can now handle quotients of κ -quantales by a simple application of the functor \mathfrak{F}^κ . Given a binary relation R on a κ -quantale L ,

- first pass to $\mathfrak{F}^\kappa L$, the free quantale over L ,
- factor $\mathfrak{F}^\kappa L$ by the relation $\tilde{R} \equiv \{(\downarrow a, \downarrow b) : aRb\}$, getting quotient map $\mu : \mathfrak{F}^\kappa L \rightarrow \mathfrak{F}^\kappa L / \tilde{R}$,
- and the desired quotient map is the range restriction of $\mu\rho_L^\kappa : L \rightarrow \mathfrak{F}^\kappa L / \tilde{R}$.

κ -quantale quotients

We can now handle quotients of κ -quantales by a simple application of the functor \mathfrak{F}^κ . Given a binary relation R on a κ -quantale L ,

- first pass to $\mathfrak{F}^\kappa L$, the free quantale over L ,
- factor $\mathfrak{F}^\kappa L$ by the relation $\tilde{R} \equiv \{(\downarrow a, \downarrow b) : aRb\}$, getting quotient map $\mu : \mathfrak{F}^\kappa L \rightarrow \mathfrak{F}^\kappa L / \tilde{R}$,
- and the desired quotient map is the range restriction of $\mu\rho_L^\kappa : L \rightarrow \mathfrak{F}^\kappa L / \tilde{R}$.

Definition

A κ -ideal U on L is R -saturated iff

$$\forall a, b, c \in L \quad (aRb \implies (ac \in U \iff bc \in U)).$$

We denote by $\langle A \rangle_R$ the smallest R -saturated κ -ideal containing a subset $A \subseteq L$.

Theorem

Let L be a κ -quantale, $\kappa > 0$, and let R be a binary relation on L . Then the R -saturated κ -ideals form a quantale in the order inherited from $\mathfrak{F}^\kappa L$, and the map

$$(a \mapsto \langle a \rangle_R) : L \rightarrow \{\langle a \rangle_R : a \in L\}$$

is a κ **Qnt**-quotient of L by the smallest κ -congruence containing R .

Colimits

Let $D = (L_i, \phi_{ij})_I$ be a diagram in $\kappa\mathbf{Qnt}$.

Colimits

Let $D = (L_i, \phi_{ij})_I$ be a diagram in $\kappa\mathbf{Qnt}$. Consider the colimit $(\delta_i : L_i \rightarrow S)_I$ in \mathbf{CMon} ,

Colimits

Let $D = (L_i, \phi_{ij})_I$ be a diagram in $\kappa\mathbf{Qnt}$. Consider the colimit $(\delta_i : L_i \rightarrow S)_I$ in \mathbf{CMon} , embed S in $\mathfrak{F}_\kappa^0 S$ via $\rho_{\kappa S}^0$,

Colimits

Let $D = (L_i, \phi_{ij})_I$ be a diagram in $\kappa\mathbf{Qnt}$. Consider the colimit $(\delta_i : L_i \rightarrow S)_I$ in \mathbf{CMon} , embed S in $\mathfrak{F}_\kappa^0 S$ via $\rho_{\kappa S}^0$, and then factor $\mathfrak{F}_\kappa^0 S$ by the smallest congruence which forces the $\rho_\kappa^0 \delta_i$'s to respect κ -joins.

Let $D = (L_i, \phi_{ij})_I$ be a diagram in $\kappa\mathbf{Qnt}$. Consider the colimit $(\delta_i : L_i \rightarrow S)_I$ in \mathbf{CMon} , embed S in $\mathfrak{F}_\kappa^0 S$ via $\rho_{\kappa S}^0$, and then factor $\mathfrak{F}_\kappa^0 S$ by the smallest congruence which forces the $\rho_\kappa^0 \delta_i$'s to respect κ -joins.

Definition

A pre-ideal $U \subseteq S$ is **saturated** iff it satisfies the following conditions.

Let $D = (L_i, \phi_{ij})_I$ be a diagram in $\kappa\mathbf{Qnt}$. Consider the colimit $(\delta_i : L_i \rightarrow S)_I$ in \mathbf{CMon} , embed S in $\mathfrak{F}_\kappa^0 S$ via $\rho_{\kappa S}^0$, and then factor $\mathfrak{F}_\kappa^0 S$ by the smallest congruence which forces the $\rho_\kappa^0 \delta_i$'s to respect κ -joins.

Definition

A pre-ideal $U \subseteq S$ is **saturated** iff it satisfies the following conditions.

- 1 For all $i \in I$ and all $a \leq b$ in L_i , and for all $s \in S$, if $\delta_i(b) s \in U$ then $\delta_i(a) s \in U$.

Let $D = (L_i, \phi_{ij})_I$ be a diagram in $\kappa\mathbf{Qnt}$. Consider the colimit $(\delta_i : L_i \rightarrow S)_I$ in \mathbf{CMon} , embed S in $\mathfrak{F}_\kappa^0 S$ via $\rho_{\kappa S}^0$, and then factor $\mathfrak{F}_\kappa^0 S$ by the smallest congruence which forces the $\rho_\kappa^0 \delta_i$'s to respect κ -joins.

Definition

A pre-ideal $U \subseteq S$ is **saturated** iff it satisfies the following conditions.

- 1 For all $i \in I$ and all $a \leq b$ in L_i , and for all $s \in S$, if $\delta_i(b) s \in U$ then $\delta_i(a) s \in U$.
- 2 For all $i \in I$ and $A \subseteq_\kappa L_i$ with $b = \bigvee A$, and for all $s \in S$, if $\delta_i(a) s \in U$ for all $a \in A$ then $\delta_i(b) s \in U$.

Let $D = (L_i, \phi_{ij})_I$ be a diagram in $\kappa\mathbf{Qnt}$. Consider the colimit $(\delta_i : L_i \rightarrow S)_I$ in \mathbf{CMon} , embed S in $\mathfrak{F}_\kappa^0 S$ via $\rho_{\kappa S}^0$, and then factor $\mathfrak{F}_\kappa^0 S$ by the smallest congruence which forces the $\rho_\kappa^0 \delta_i$'s to respect κ -joins.

Definition

A pre-ideal $U \subseteq S$ is **saturated** iff it satisfies the following conditions.

- 1 For all $i \in I$ and all $a \leq b$ in L_i , and for all $s \in S$, if $\delta_i(b) s \in U$ then $\delta_i(a) s \in U$.
- 2 For all $i \in I$ and $A \subseteq_\kappa L_i$ with $b = \bigvee A$, and for all $s \in S$, if $\delta_i(a) s \in U$ for all $a \in A$ then $\delta_i(b) s \in U$.

Let $[A]_R$ designate the smallest R -saturated pre-ideal containing a subset $A \subseteq S$.

Let $D = (L_i, \phi_{ij})_I$ be a diagram in $\kappa\mathbf{Qnt}$. Consider the colimit $(\delta_i : L_i \rightarrow S)_I$ in \mathbf{CMon} , embed S in $\mathfrak{F}_\kappa^0 S$ via $\rho_{\kappa S}^0$, and then factor $\mathfrak{F}_\kappa^0 S$ by the smallest congruence which forces the $\rho_\kappa^0 \delta_i$'s to respect κ -joins.

Definition

A pre-ideal $U \subseteq S$ is **saturated** iff it satisfies the following conditions.

- ① For all $i \in I$ and all $a \leq b$ in L_i , and for all $s \in S$, if $\delta_i(b) s \in U$ then $\delta_i(a) s \in U$.
- ② For all $i \in I$ and $A \subseteq_\kappa L_i$ with $b = \bigvee A$, and for all $s \in S$, if $\delta_i(a) s \in U$ for all $a \in A$ then $\delta_i(b) s \in U$.

Let $[A]_R$ designate the smallest R -saturated pre-ideal containing a subset $A \subseteq S$. Let

$$\tilde{L} \equiv \{[A]_R : A \subseteq_\kappa L\},$$

and let $\gamma_i : L_i \rightarrow \tilde{L} \equiv (a \mapsto [a]_R, a \in A)$.

Theorem

$(\gamma_i : L_i \rightarrow \tilde{L})$ is a $\kappa\mathbf{Qnt}$ colimit of the diagram $D = (L_i, \phi_{ij})_I$.

Theorem

$(\gamma_i : L_i \rightarrow \tilde{L})$ is a $\kappa\mathbf{Qnt}$ colimit of the diagram $D = (L_i, \phi_{ij})_I$.

This directly generalizes to κ -quantales Johnstone's construction of the frame colimit.

Coproducts

This description of the colimit specializes to coproducts.

Coproducts

This description of the colimit specializes to coproducts. The simplification occurs because the underlying **CMon** coproduct S is transparent.

Coproducts

This description of the colimit specializes to coproducts. The simplification occurs because the underlying **CMon** coproduct S is transparent. The resulting $\kappa\mathbf{Qnt}$ coproduct construction is intuitive and explicit; and lends itself to a characterization in general terms.

This description of the colimit specializes to coproducts. The simplification occurs because the underlying **CMon** coproduct S is transparent. The resulting $\kappa\mathbf{Qnt}$ coproduct construction is intuitive and explicit; and lends itself to a characterization in general terms.

Theorem

Let $\kappa > 0$. A family $(v_i : L_i \rightarrow L)_J$ of κ -morphisms is a $\kappa\mathbf{Qnt}$ coproduct of the family $(L_i)_J$ iff it has these properties.

This description of the colimit specializes to coproducts. The simplification occurs because the underlying **CMon** coproduct S is transparent. The resulting $\kappa\mathbf{Qnt}$ coproduct construction is intuitive and explicit; and lends itself to a characterization in general terms.

Theorem

Let $\kappa > 0$. A family $(v_i : L_i \rightarrow L)_J$ of κ -morphisms is a $\kappa\mathbf{Qnt}$ coproduct of the family $(L_i)_J$ iff it has these properties.

- 1 $\bigcup_J v_i [L_i]$ generates L .

This description of the colimit specializes to coproducts. The simplification occurs because the underlying **CMon** coproduct S is transparent. The resulting $\kappa\mathbf{Qnt}$ coproduct construction is intuitive and explicit; and lends itself to a characterization in general terms.

Theorem

Let $\kappa > 0$. A family $(v_i : L_i \rightarrow L)_J$ of κ -morphisms is a $\kappa\mathbf{Qnt}$ coproduct of the family $(L_i)_J$ iff it has these properties.

- 1 $\bigcup_J v_i [L_i]$ generates L .
- 2 For any $I_0 \subseteq_\omega J$ and $I_1 \subseteq_\kappa J$, and for any choice of $a_i \in L_i$, $i \in I_0$, and $b_j \in L_j$, $j \in I_1$,

$$\bigwedge_{I_0} v_i (a_i) \leq \bigvee_{I_1} v_j (b_j) \implies \exists i \in I_0 \cap I_1 (a_i \leq b_i).$$

Free κ -quantales (over sets)

This description of coproducts specializes to free κ -quantales.

Free κ -quantales (over sets)

This description of coproducts specializes to free κ -quantales. This is the generalization to $\kappa\mathbf{Qnt}$ of Whitman's condition for distributive lattices.

Free κ -quantales (over sets)

This description of coproducts specializes to free κ -quantales. This is the generalization to $\kappa\mathbf{Qnt}$ of Whitman's condition for distributive lattices.

Theorem

Let L be a κ -quantale, $\kappa > 0$, generated by a subset X . Then L is freely generated by X iff for any $X_0 \subseteq_{\omega} X$ and $Y \subseteq_{\kappa} X$, and for any choice of integers $n_x, m_y \in \mathbb{Z}^+$, $x \in X_0, y \in Y$,

$$\prod_{X_0} x^{n_x} \leq \bigvee_Y y^{m_y} \implies \exists x \in X_0 \cap Y \quad (n_x \geq m_y).$$