


Identifying groups of finite Morley rank with a split BN -pair of rank 1

Josh Wiscons


University of Colorado, Boulder

Structures of fMr




Intro: groups of finite Morley rank (fMr)

Structures of fMr




Intro: groups of finite Morley rank (fMr)

Structures of fMr



Intro: groups of finite Morley rank (fMr)

Structures of fMr



Fix a first order language \mathcal{L} and a (ω -saturated) \mathcal{L} -structure \mathcal{M} . Morley rank assigns to each definable set, X , either -1 , an **ordinal**, or ∞ denoted $\text{RM}(X)$:


Fix a first order language \mathcal{L} and a (ω -saturated) \mathcal{L} -structure \mathcal{M} . Morley rank assigns to each definable set, X , either -1 , an **ordinal**, or ∞ denoted $\text{RM}(X)$:

- $\text{RM}(X) \geq 0$ iff X is non-empty;

Morley rank

Fix a first order language \mathcal{L} and a (ω -saturated) \mathcal{L} -structure \mathcal{M} . Morley rank assigns to each definable set, X , either -1 , an **ordinal**, or ∞ denoted $\text{RM}(X)$:


- $\text{RM}(X) \geq 0$ iff X is non-empty;
- $\text{RM}(X) \geq \alpha + 1$ iff there exists definable Y_i , $1 \leq i < \omega$, such that



Morley rank

Fix a first order language \mathcal{L} and a (ω -saturated) \mathcal{L} -structure \mathcal{M} . Morley rank assigns to each definable set, X , either -1 , an ordinal, or ∞ denoted $\text{RM}(X)$:

- $\text{RM}(X) \geq 0$ iff X is non-empty;
- $\text{RM}(X) \geq \alpha + 1$ iff there exists definable Y_i , $1 \leq i < \omega$, such that




- $\text{RM}(X) \geq \lambda$ for λ a limit ordinal iff $\text{RM}(X) \geq \alpha$ for all $\alpha < \lambda$;

Morley rank

Fix a first order language \mathcal{L} and a (ω -saturated) \mathcal{L} -structure \mathcal{M} . Morley rank assigns to each definable set, X , either -1 , an ordinal, or ∞ denoted $\text{RM}(X)$:

- $\text{RM}(X) \geq 0$ iff X is non-empty;
- $\text{RM}(X) \geq \alpha + 1$ iff there exists definable Y_i , $1 \leq i < \omega$, such that




- $\text{RM}(X) \geq \lambda$ for λ a limit ordinal iff $\text{RM}(X) \geq \alpha$ for all $\alpha < \lambda$;
- $\text{RM}(X) = \infty$ iff $\text{RM}(X) \geq \alpha$ for all ordinals α .

Morley rank

Fix a first order language \mathcal{L} and a (ω -saturated) \mathcal{L} -structure \mathcal{M} . Morley rank assigns to each definable set, X , either -1 , an ordinal, or ∞ denoted $\text{RM}(X)$:

- $\text{RM}(X) \geq 0$ iff X is non-empty;
- $\text{RM}(X) \geq \alpha + 1$ iff there exists definable Y_i , $1 \leq i < \omega$, such that



- $\text{RM}(X) \geq \lambda$ for λ a limit ordinal iff $\text{RM}(X) \geq \alpha$ for all $\alpha < \lambda$;
- $\text{RM}(X) = \infty$ iff $\text{RM}(X) \geq \alpha$ for all ordinals α .

$\text{RM}(\mathcal{M})$ is defined to be $\text{RM}(M)$.

Groups of finite Morley rank

Definition

A group of finite Morley rank is an expansion of a “pure group”, i.e. $(G; \cdot, ^{-1}, 1, \dots)$, that has finite Morley rank (fMr).

Groups of finite Morley rank

Definition

A group of finite Morley rank is an expansion of a “pure group”, i.e. $(G; \cdot, ^{-1}, 1, \dots)$, that has finite Morley rank (fMr).

- RM (together with Morley degree) gives DCC on definable subgroups.
 - This only requires $\text{RM}(G) \neq \infty$.

Groups of finite Morley rank

Definition

A group of finite Morley rank is an expansion of a “pure group”, i.e. $(G; \cdot, ^{-1}, 1, \dots)$, that has finite Morley rank (fMr).

- RM (together with Morley degree) gives DCC on definable subgroups.
 - This only requires $\text{RM}(G) \neq \infty$.
- (Cherlin-Macintyre) An infinite division ring has fMr iff it is an algebraically closed field.

Groups of finite Morley rank

Definition

A group of finite Morley rank is an expansion of a “pure group”, i.e. $(G; \cdot, ^{-1}, 1, \dots)$, that has finite Morley rank (fMr).

- RM (together with Morley degree) gives DCC on definable subgroups.
 - This only requires $\text{RM}(G) \neq \infty$.
- (Cherlin-Macintyre) An infinite division ring has fMr iff it is an algebraically closed field.
- Structures interpretable in structures of fMr have fMr.

Groups of finite Morley rank

Definition

A group of finite Morley rank is an expansion of a “pure group”, i.e. $(G; \cdot, ^{-1}, 1, \dots)$, that has finite Morley rank (fMr).

- RM (together with Morley degree) gives DCC on definable subgroups.
 - This only requires $\text{RM}(G) \neq \infty$.
- (Cherlin-Macintyre) An infinite division ring has fMr iff it is an algebraically closed field.
- Structures interpretable in structures of fMr have fMr.

Examples

- 1 Affine algebraic groups over an algebraically closed field, K , have fMr: $\text{GL}_n(K)$, $\text{PSL}_n(K)$, finite groups, \dots

Groups of finite Morley rank

Definition

A **group of finite Morley rank** is an expansion of a “pure group”, i.e. $(G; \cdot, ^{-1}, 1, \dots)$, that has finite Morley rank (fMr).

- RM (together with Morley degree) gives DCC on definable subgroups.
 - This only requires $\text{RM}(G) \neq \infty$.
- (Cherlin-Macintyre) An infinite division ring has fMr iff it is an algebraically closed field.
- Structures interpretable in structures of fMr have fMr.

Examples

- 1 Affine algebraic groups over an algebraically closed field, K , have fMr: $\text{GL}_n(K)$, $\text{PSL}_n(K)$, finite groups, \dots
- 2 \mathbb{Z} does **NOT** have fMr because $\mathbb{Z} \not\cong 2\mathbb{Z} \not\cong 4\mathbb{Z} \not\cong 8\mathbb{Z} \not\cong \dots$

Cherlin-Zil'ber Conjecture

Conjecture (Cherlin-Zil'ber)

An infinite simple group of finite Morley rank is isomorphic to an (affine) algebraic group over an algebraically closed field.

Cherlin-Zil'ber Conjecture

Conjecture (Cherlin-Zil'ber)

An infinite simple group of finite Morley rank is isomorphic to an (affine) algebraic group over an algebraically closed field.

The analysis breaks into 4 types based on the structure of a Sylow 2-subgroup, S . The types are

Degenerate: $S^\circ = 1$,

Odd: S° is nontrivial, divisible, and abelian (S° is a 2-torus),

Even: S° is nontrivial, nilpotent, and of bounded exponent (S° is 2-unipotent), and

Mixed: S° contains a 2-torus and a 2-unipotent subgroup.

Cherlin-Zil'ber Conjecture

Conjecture (Cherlin-Zil'ber)

An infinite simple group of finite Morley rank is isomorphic to an (affine) algebraic group over an algebraically closed field.

The analysis breaks into 4 types based on the structure of a Sylow 2-subgroup, S . The types are

Degenerate: $S^\circ = 1$,

Odd: S° is nontrivial, divisible, and abelian (S° is a 2-torus),


Even: S° is nontrivial, nilpotent, and of bounded exponent (S° is 2-unipotent), and

Mixed: S° contains a 2-torus and a 2-unipotent subgroup.


Theorem (Altinel-Borovik-Cherlin)


There are no infinite simple groups of finite Morley rank of mixed type and those of even type are indeed algebraic.

Theorem of Altinel, Borovik, and Cherlin




Theorem of Altinel, Borovik, and Cherlin





- We consider the following inductive frameworks.
 - K^* -groups: groups in which every proper definable simple section is an algebraic group.



- We consider the following inductive frameworks.
 - K^* -groups: groups in which every proper definable simple section is an algebraic group.
 - L^* -groups: groups in which every proper definable simple section of **odd type** is an algebraic group.

Recognizing algebraic groups

Recognizing algebraic groups

Theorem (Kramer-Tent-Van Maldeghem)

*An infinite simple group of finite Morley rank with an irreducible BN-pair of **Tits rank at least 3** is isomorphic to an algebraic group over an algebraically closed field. The same holds for **Tits rank 2** if the associated polygon is Moufang.*

Recognizing algebraic groups

Theorem (Kramer-Tent-Van Maldeghem)

*An infinite simple group of finite Morley rank with an irreducible BN-pair of **Tits rank at least 3** is isomorphic to an algebraic group over an algebraically closed field. The same holds for **Tits rank 2** if the associated polygon is Moufang.*

Question

What can be said about the Tits rank 1 case?

Recognizing algebraic groups

Theorem (Kramer-Tent-Van Maldeghem)

*An infinite simple group of finite Morley rank with an irreducible BN-pair of **Tits rank at least 3** is isomorphic to an algebraic group over an algebraically closed field. The same holds for **Tits rank 2** if the associated polygon is Moufang.*

Question

What can be said about the Tits rank 1 case?

- *BN*-pairs for a group give rise to a geometry on which the group acts.

Recognizing algebraic groups

Theorem (Kramer-Tent-Van Maldeghem)

An infinite simple group of finite Morley rank with an irreducible BN-pair of Tits rank at least 3 is isomorphic to an algebraic group over an algebraically closed field. The same holds for Tits rank 2 if the associated polygon is Moufang.

Question

What can be said about the Tits rank 1 case?

- *BN*-pairs for a group give rise to a geometry on which the group acts.
- In Tits rank 1, the geometry degenerates, and we simply have a group acting 2-transitively on a set.

Recognizing algebraic groups

Theorem (Kramer-Tent-Van Maldeghem)

An infinite simple group of finite Morley rank with an irreducible BN-pair of Tits rank at least 3 is isomorphic to an algebraic group over an algebraically closed field. The same holds for Tits rank 2 if the associated polygon is Moufang.


Question

What can be said about the Tits rank 1 case?

- *BN*-pairs for a group give rise to a geometry on which the group acts.
- In Tits rank 1, the geometry degenerates, and we simply have a group acting 2-transitively on a set.
- We focus on groups with a **split** *BN*-pair of Tits rank 1.

Split BN -pairs of Tits rank 1


General picture:



- $G = B \sqcup BnB$ for some $n \in N - H$


Split BN -pairs of Tits rank 1

General picture:



- $G = B \sqcup BnB$ for some $n \in N - H$


$\mathrm{PSL}_2(K)$:



- $\mathrm{PSL}_2(K) = B \sqcup B \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} B$


Split BN -pairs of Tits rank 1

General picture:



- $G = B \sqcup BnB$ for some $n \in N - H$

$\mathrm{PSL}_2(K)$:



- $\mathrm{PSL}_2(K) = B \sqcup B \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} B$
- $\begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix} \cong K^+$ and $\begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} \cong K^\times$

The main result

Theorem (W)

Let G be an infinite simple L^ -group of finite Morley rank with a split BN -pair of Tits rank 1. If U is infinite and abelian, then $G \cong \mathrm{PSL}_2(K)$ for K an algebraically closed field.*

The main result

Theorem (W)

Let G be an infinite simple L^ -group of finite Morley rank with a split BN -pair of Tits rank 1. If U is infinite and abelian, then $G \cong \mathrm{PSL}_2(K)$ for K an algebraically closed field.*

An outline.

The main result

Theorem (W)

Let G be an infinite simple L^ -group of finite Morley rank with a split BN -pair of Tits rank 1. If U is infinite and abelian, then $G \cong \mathrm{PSL}_2(K)$ for K an algebraically closed field.*

An outline.

- ① The focal points are $B = U \rtimes H$ and $N - H$.
 - We hope $U \cong K^+$, $H \cong K^\times$, and $N - H$ consists entirely of involutions.

The main result

Theorem (W)

Let G be an infinite simple L^ -group of finite Morley rank with a split BN -pair of Tits rank 1. If U is infinite and abelian, then $G \cong \mathrm{PSL}_2(K)$ for K an algebraically closed field.*

An outline.

- 0 The focal points are $B = U \rtimes H$ and $N - H$.
 - We hope $U \cong K^+$, $H \cong K^\times$, and $N - H$ consists entirely of involutions.
- 1 U has the structure of a vector space.
 - May assume that $\mathrm{char}(U) > 2$ (using De Medts and Tent; Wiscons).

The main result

Theorem (W)

Let G be an infinite simple L^ -group of finite Morley rank with a split BN -pair of Tits rank 1. If U is infinite and abelian, then $G \cong \mathrm{PSL}_2(K)$ for K an algebraically closed field.*

An outline.

- 0 The focal points are $B = U \rtimes H$ and $N - H$.
 - We hope $U \cong K^+$, $H \cong K^\times$, and $N - H$ consists entirely of involutions.
- 1 U has the structure of a vector space.
 - May assume that $\mathrm{char}(U) > 2$ (using De Medts and Tent; Wiscons).
- 2 In a minimal counterexample:

The main result

Theorem (W)

Let G be an infinite simple L^ -group of finite Morley rank with a split BN -pair of Tits rank 1. If U is infinite and abelian, then $G \cong \mathrm{PSL}_2(K)$ for K an algebraically closed field.*

An outline.

- 0 The focal points are $B = U \rtimes H$ and $N - H$.
 - We hope $U \cong K^+$, $H \cong K^\times$, and $N - H$ consists entirely of involutions.
- 1 U has the structure of a vector space.
 - May assume that $\mathrm{char}(U) > 2$ (using De Medts and Tent; Wiscons).
- 2 In a minimal counterexample:
 - i. H has no infinite normal solvable subgroups (using De Medts and Tent);

The main result

Theorem (W)

Let G be an infinite simple L^ -group of finite Morley rank with a split BN -pair of Tits rank 1. If U is infinite and abelian, then $G \cong \mathrm{PSL}_2(K)$ for K an algebraically closed field.*

An outline.

- 0 The focal points are $B = U \rtimes H$ and $N - H$.
 - We hope $U \cong K^+$, $H \cong K^\times$, and $N - H$ consists entirely of involutions.
- 1 U has the structure of a vector space.
 - May assume that $\mathrm{char}(U) > 2$ (using De Medts and Tent; Wiscons).
- 2 In a minimal counterexample:
 - i. H has no infinite normal solvable subgroups (using De Medts and Tent);
 - ii. H has a definable normal quasisimple subgroup Q ;

The main result

Theorem (W)

Let G be an infinite simple L^ -group of finite Morley rank with a split BN -pair of Tits rank 1. If U is infinite and abelian, then $G \cong \mathrm{PSL}_2(K)$ for K an algebraically closed field.*

An outline.

- 0 The focal points are $B = U \rtimes H$ and $N - H$.
 - We hope $U \cong K^+$, $H \cong K^\times$, and $N - H$ consists entirely of involutions.
- 1 U has the structure of a vector space.
 - May assume that $\mathrm{char}(U) > 2$ (using De Medts and Tent; Wiscons).
- 2 In a minimal counterexample:
 - i. H has no infinite normal solvable subgroups (using De Medts and Tent);
 - ii. H has a definable normal quasisimple subgroup Q ;
 - iii. Q has infinite 2-subgroups, so Q is algebraic by the L^* -hypothesis;

The main result

Theorem (W)

Let G be an infinite simple L^ -group of finite Morley rank with a split BN -pair of Tits rank 1. If U is infinite and abelian, then $G \cong \mathrm{PSL}_2(K)$ for K an algebraically closed field.*

An outline.

- 0 The focal points are $B = U \rtimes H$ and $N - H$.
 - We hope $U \cong K^+$, $H \cong K^\times$, and $N - H$ consists entirely of involutions.
- 1 U has the structure of a vector space.
 - May assume that $\mathrm{char}(U) > 2$ (using De Medts and Tent; Wiscons).
- 2 In a minimal counterexample:
 - i. H has no infinite normal solvable subgroups (using De Medts and Tent);
 - ii. H has a definable normal quasisimple subgroup Q ;
 - iii. Q has infinite 2-subgroups, so Q is algebraic by the L^* -hypothesis;
 - iv. $H = Q$, so H is algebraic;

The main result

Theorem (W)

Let G be an infinite simple L^ -group of finite Morley rank with a split BN -pair of Tits rank 1. If U is infinite and abelian, then $G \cong \mathrm{PSL}_2(K)$ for K an algebraically closed field.*

An outline.


- 0 The focal points are $B = U \rtimes H$ and $N = H$.
 - We hope $U \cong K^+$, $H \cong K^\times$, and $N = H$ consists entirely of involutions.
- 1 U has the structure of a vector space.
 - May assume that $\mathrm{char}(U) > 2$ (using De Medts and Tent; Wiscons).
- 2 In a minimal counterexample:
 - i. H has no infinite normal solvable subgroups (using De Medts and Tent);
 - ii. H has a definable normal quasisimple subgroup Q ;
 - iii. Q has infinite 2-subgroups, so Q is algebraic by the L^* -hypothesis;
 - iv. $H = Q$, so H is algebraic;
 - v. a Borel subgroup of H has an infinite centralizer CONTRADICTING (i).

Summary and questions

We have that for G an infinite simple L^* -group of fMr. . .

Summary and questions

We have that for G an infinite simple L^* -group of fMr. . .




+

- U is abelian
- H is arbitrary

Summary and questions

We have that for G an infinite simple L^* -group of fMr...




+

- U is abelian
- H is arbitrary

 $\implies G \cong \text{PSL}_2(K)$

Summary and questions


We have that for G an infinite simple L^* -group of fMr...



Questions

Summary and questions

We have that for G an infinite simple L^* -group of fMr...




Questions

Q1: Can we generalize to U solvable (or nilpotent) and H arbitrary?

Summary and questions

We have that for G an infinite simple L^* -group of fMr...



$$+ \begin{array}{l} \bullet U \text{ is } \text{arbitrary} \\ \quad \quad \quad \text{abelian} \\ \bullet H \text{ is } \text{arbitrary} \\ \quad \quad \quad \text{abelian} \end{array} \stackrel{?}{\implies} G \cong \text{PSL}_2(K)$$


Questions

Q1: Can we generalize to U solvable (or nilpotent) and H arbitrary?

Q2: Can we address when U is arbitrary and H is abelian?

Summary and questions

We have that for G an infinite simple L^* -group of fMr...



Questions

Q1: Can we generalize to U solvable (or nilpotent) and H arbitrary?

Q2: Can we address when U is arbitrary and H is abelian?

Q3: Can we apply the main result to simple groups with Prüfer 2-rank 1?