

Relational Tame Congruence Theory

Mike Behrisch

Boulder, CO, 6 June 2010

Outline

- 1 Preliminaries and notations
- 2 “Relational TCT” as a localisation theory
- 3 Examples of irreducible algebras / neighbourhoods

Outline

- 1 Preliminaries and notations
- 2 “Relational TCT” as a localisation theory
- 3 Examples of irreducible algebras / neighbourhoods

Polymorphisms and invariant relations

Let A be a set, $n \in \mathbb{N}$ and $m \in \mathbb{N}_+$.

$$O_A^{(n)} := A^{A^n}$$

$$R_A^{(m)} := \mathcal{P}(A^m)$$

$$O_A := \bigcup_{n \in \mathbb{N}} O_A^{(n)}$$

$$R_A := \bigcup_{m \in \mathbb{N}_+} R_A^{(m)}$$

For $f \in O_A^{(n)}$ and $S \subseteq A^m$ (i.e. $S \in R_A^{(m)}$):

$$f \triangleright S \iff S \in \text{Sub}(\langle A; f \rangle^m)$$

Polymorphisms and invariant relations

Let A be a set, $n \in \mathbb{N}$ and $m \in \mathbb{N}_+$.

$$O_A^{(n)} := A^{A^n}$$

$$R_A^{(m)} := \mathcal{P}(A^m)$$

$$O_A := \bigcup_{n \in \mathbb{N}} O_A^{(n)}$$

$$R_A := \bigcup_{m \in \mathbb{N}_+} R_A^{(m)}$$

For $f \in O_A^{(n)}$ and $S \subseteq A^m$ (i.e. $S \in R_A^{(m)}$):

$$f \triangleright S \iff S \in \text{Sub}(\langle A; f \rangle^m)$$

$$\iff f \in \text{Hom}(\langle A; S \rangle^n; \langle A; S \rangle)$$

Polymorphisms and invariant relations

For $F \subseteq O_A$ and $Q \subseteq R_A$:

$$\begin{aligned} \text{Inv} \langle A; F \rangle &:= \text{Inv}_A F := \{ S \in R_A \mid \forall f \in F : f \triangleright S \} \\ &= \bigcup_{m \in \mathbb{N}_+} \text{Sub}(\langle A; F \rangle^m) \end{aligned}$$

$$\begin{aligned} \text{Pol} \langle A; Q \rangle &:= \text{Pol}_A Q := \{ f \in O_A \mid \forall S \in Q : f \triangleright S \} \\ &= \bigcup_{n \in \mathbb{N}} \text{Hom}(\langle A; Q \rangle^n; \langle A; Q \rangle) \end{aligned}$$

$$\text{Clo}(\mathbf{A}) := \text{Pol Inv } \mathbf{A} = T(\mathbf{A}) \quad (\text{term operations}).$$


Outline

- 1 Preliminaries and notations
- 2 “Relational TCT” as a localisation theory
- 3 Examples of irreducible algebras / neighbourhoods


Restricting algebras to neighbourhoods

A


Restricting algebras to neighbourhoods




Restricting algebras to neighbourhoods




Restricting algebras to neighbourhoods




Restricting algebras to neighbourhoods




Restricting algebras to neighbourhoods




Restricting algebras to neighbourhoods




Restricting algebras to neighbourhoods



Restricting algebras to neighbourhoods




Restricting algebras to neighbourhoods



Definition (neighbourhood)

$U \in \text{Neigh } \mathbf{A} : \iff$
 $U = e[A]$ for some
 $e \in \text{Idem } \mathbf{A} :=$
 $\{ g \in \text{Clo}^{(1)}(\mathbf{A}) \mid g^2 = g \}$


Restricting algebras to neighbourhoods



Definition (neighbourhood)

$U \in \text{Neigh } \mathbf{A} : \iff$
 $U = e[A]$ for some
 $e \in \text{Idem } \mathbf{A} :=$
 $\{ g \in \text{Clo}^{(1)}(\mathbf{A}) \mid g^2 = g \}$


Restricting algebras to neighbourhoods



Definition (neighbourhood)

$U \in \text{Neigh } \mathbf{A} : \iff$
 $U = e[A]$ for some
 $e \in \text{Idem } \mathbf{A} :=$
 $\{ g \in \text{Clo}^{(1)}(\mathbf{A}) \mid g^2 = g \}$

Restricting algebras to neighbourhoods




Definition (neighbourhood)


$U \in \text{Neigh } \mathbf{A} : \iff$
 $U = e[A]$ for some
 $e \in \text{Idem } \mathbf{A} :=$
 $\{g \in \text{Clo}^{(1)}(\mathbf{A}) \mid g^2 = g\}$

$$\begin{aligned}
 \mathbf{A}|_U &= \langle U; \{ (e \circ f) \uparrow_{U^{ar f}}^U \mid f \in \text{Clo}(\mathbf{A}) \} \rangle \\
 &= \langle U; \{ f \uparrow_{U^{ar f}}^U \mid f \in \text{Clo}(\mathbf{A}) \wedge f \triangleright U \} \rangle
 \end{aligned}$$


Distinguishing invariant relations




Distinguishing invariant relations




Distinguishing invariant relations




Distinguishing invariant relations




Distinguishing invariant relations




Distinguishing invariant relations



Distinguishing invariant relations



Distinguishing invariant relations



Definition (cover)

$\mathcal{U} \subseteq \text{Neigh } \mathbf{A}$ **cover of \mathbf{A}** iff for all $m \in \mathbb{N}_+$ and $S, T \in \text{Inv}^{(m)} \mathbf{A}$ holds

$$S \neq T \implies \exists U \in \mathcal{U} : S|_U \neq T|_U.$$

Characterisation of covers



Characterisation of covers



Theorem (Kearnes, A. Szendrei, 2001)

For an algebra \mathbf{A} and a collection $\mathcal{U} \subseteq \text{Neigh } \mathbf{A}$ of neighbourhoods of \mathbf{A} (where each $U \in \mathcal{U}$ satisfies $U = e_U[\mathbf{A}]$ for some fixed $e_U \in \text{Idem } \mathbf{A}$) t.f.a.e.:

- 1 \mathcal{U} is a **cover** of \mathbf{A} .

Characterisation of covers



Theorem (Kearnes, A. Szendrei, 2001)

For an algebra \mathbf{A} and a collection $\mathcal{U} \subseteq \text{Neigh } \mathbf{A}$ of neighbourhoods of \mathbf{A} (where each $U \in \mathcal{U}$ satisfies $U = e_U[A]$ for some fixed $e_U \in \text{Idem } \mathbf{A}$) t.f.a.e.:

- 1 \mathcal{U} is a **cover** of \mathbf{A} .
- 2 There is some $q \in \mathbb{N}_+$, tuples $(U_1, \dots, U_q) \in \mathcal{U}^q$ and $(f_1, \dots, f_q) \in (\text{Clo}^{(1)}(\mathbf{A}))^q$ and a term operation $\lambda \in \text{Clo}^{(q)}(\mathbf{A})$ such that for all $x \in A$ holds

$$\lambda(e_{U_1} \circ f_1(x), \dots, e_{U_q} \circ f_q(x)) = x.$$

Characterisation of covers



Theorem (Kearnes, A. Szendrei, 2001)

For an algebra \mathbf{A} and a collection $\mathcal{U} \subseteq \text{Neigh } \mathbf{A}$ of neighbourhoods of \mathbf{A} (where each $U \in \mathcal{U}$ satisfies $U = e_U[A]$ for some fixed $e_U \in \text{Idem } \mathbf{A}$) t.f.a.e.:

- 1 \mathcal{U} is a **cover** of \mathbf{A} .
- 2 There is some $q \in \mathbb{N}_+$, tuples $(U_1, \dots, U_q) \in \mathcal{U}^q$ and $(f_1, \dots, f_q) \in (\text{Clo}^{(1)}(\mathbf{A}))^q$ and a term operation $\lambda \in \text{Clo}^{(q)}(\mathbf{A})$ such that for all $x \in A$ holds

$$\lambda(e_{U_1} \circ f_1(x), \dots, e_{U_q} \circ f_q(x)) = x.$$

- 3 There is some $q \in \mathbb{N}_+$ and a tuple $(U_1, \dots, U_q) \in \mathcal{U}^q$ such that \mathbf{A} is a retract of $\mathbf{A}|_{U_1} \times \dots \times \mathbf{A}|_{U_q}$

Characterisation of covers


Theorem (Kearnes, A. Szendrei, 2001)

For an algebra \mathbf{A} and a collection $\mathcal{U} \subseteq \text{Neigh } \mathbf{A}$ of neighbourhoods of \mathbf{A} (where each $U \in \mathcal{U}$ satisfies $U = e_U[\mathbf{A}]$ for some fixed $e_U \in \text{Idem } \mathbf{A}$) t.f.a.e.:

- 1 \mathcal{U} is a **cover** of \mathbf{A} .
- 2 There is some $q \in \mathbb{N}_+$, tuples $(U_1, \dots, U_q) \in \mathcal{U}^q$ and $(f_1, \dots, f_q) \in (\text{Clo}^{(1)}(\mathbf{A}))^q$ and a term operation $\lambda \in \text{Clo}^{(q)}(\mathbf{A})$ such that for all $x \in A$ holds

$$\lambda(e_{U_1} \circ f_1(x), \dots, e_{U_q} \circ f_q(x)) = x.$$

- 3 There is some $q \in \mathbb{N}_+$ and a tuple $(U_1, \dots, U_q) \in \mathcal{U}^q$ such that $\widetilde{\mathbf{A}}$ is a retract of $\widetilde{\mathbf{A}}|_{U_1} \times \dots \times \widetilde{\mathbf{A}}|_{U_q}$, i.e. there are relational morphisms such that



When localisation is useless . . .

Definition (Irreducibility)

An algebra \mathbf{A} is called **irreducible**, iff every cover $\mathcal{U} \subseteq \text{Neigh } \mathbf{A}$ necessarily contains the neighbourhood $A \in \mathcal{U}$.

When localisation is useless . . .

Definition (Irreducibility)

An algebra \mathbf{A} is called **irreducible**, iff every cover $\mathcal{U} \subseteq \text{Neigh } \mathbf{A}$ necessarily contains the neighbourhood $A \in \mathcal{U}$.

Definition (Irreducible neighbourhood)

$U \in \text{Neigh } \mathbf{A}$ **irreducible** : $\iff \mathbf{A}|_U$ irreducible.

When localisation is useless . . .

Definition (Irreducibility)

An algebra \mathbf{A} is called **irreducible**, iff every cover $\mathcal{U} \subseteq \text{Neigh } \mathbf{A}$ necessarily contains the neighbourhood $A \in \mathcal{U}$.

Definition (Irreducible neighbourhood)

$U \in \text{Neigh } \mathbf{A}$ **irreducible** : $\iff \mathbf{A}|_U$ irreducible.

Theorem (Kearnes, A. Szendrei, 2001)

A finite algebra \mathbf{A} is irreducible iff the set of all **unary non-bijective term operations** is an **invariant** relation,
i.e. $\text{Clo}^{(1)}(\mathbf{A}) \setminus \text{Sym } A \in \text{Sub}(\mathbf{A}^A)$.


Refinement of covers

$$\mathcal{V} \leq_{\text{ref}} \mathcal{U}$$

quasiorder

idea: smaller neighbour-
hoods

Refinement of covers




$$v \leq_{\text{ref}} u$$

quasiorder

idea: smaller neighbourhoods

Refinement of covers




$$v \leq_{\text{ref}} u$$

quasiorder

idea: smaller neighbourhoods

Refinement of covers



$$v \leq_{\text{ref}} u$$

quasiorder

idea: smaller neighbourhoods

Existence and uniqueness of covers

Theorem (Kearnes, A. Szendrei, 2001)

Every finite algebra \mathbf{A} has got **exactly one nonrefinable, irredundant cover \mathcal{U} up to isomorphism**. All neighbourhoods in \mathcal{U} are **irreducible**.

Existence and uniqueness of covers

Theorem (Kearnes, A. Szendrei, 2001)

Every finite algebra \mathbf{A} has got **exactly one nonrefinable, irredundant cover** \mathcal{U} **up to isomorphism**. All neighbourhoods in \mathcal{U} are **irreducible**.

Consequence:

cover Decomposition of algebras in small parts (up to term equivalence)

uniqueness exactly one distinguished cover up to isomorphism consisting of irreducible neighbourhoods

irreducible algebras = basic building blocks of finite algebras

check by $\text{Clo}^{(1)}(\mathbf{A}) \setminus \text{Sym } A \in \text{Sub}(\mathbf{A}^A)$

Outline

- 1 Preliminaries and notations
- 2 “Relational TCT” as a localisation theory
- 3 Examples of irreducible algebras / neighbourhoods

Irreducible classes of algebras

i.e. all member are irreducible algebras

- ① $|A| = 1$: one-element algebras without nullary constants

Irreducible classes of algebras

i.e. all member are irreducible algebras

- 1 $|A| = 1$: one-element algebras without nullary constants
- 2 $|A| = 2$: all two-element algebras

Irreducible classes of algebras

i.e. all member are irreducible algebras

- 1 $|A| = 1$: one-element algebras without nullary constants
- 2 $|A| = 2$: all two-element algebras
- 3 essentially unary algebras $\text{Clo}(\mathbf{A}) = \langle \text{Clo}^{(1)}(\mathbf{A}) \rangle_{O_A}$

Irreducible classes of algebras

i.e. all member are irreducible algebras

- 1 $|A| = 1$: one-element algebras without nullary constants
- 2 $|A| = 2$: all two-element algebras
- 3 essentially unary algebras $\text{Clo}(\mathbf{A}) = \langle \text{Clo}^{(1)}(\mathbf{A}) \rangle_{O_A}$
- 4 all idempotent algebras ($f(x, \dots, x) \approx x$)

Irreducible classes of algebras

i.e. all member are irreducible algebras

- 1 $|A| = 1$: one-element algebras without nullary constants
- 2 $|A| = 2$: all two-element algebras
- 3 essentially unary algebras $\text{Clo}(\mathbf{A}) = \langle \text{Clo}^{(1)}(\mathbf{A}) \rangle_{O_A}$
- 4 all idempotent algebras ($f(x, \dots, x) \approx x$)
especially all semilattices, lattices, Boolean algebras

Irreducible classes of algebras

i.e. all member are irreducible algebras

- 1 $|A| = 1$: one-element algebras without nullary constants
- 2 $|A| = 2$: all two-element algebras
- 3 essentially unary algebras $\text{Clo}(\mathbf{A}) = \langle \text{Clo}^{(1)}(\mathbf{A}) \rangle_{O_A}$
- 4 all idempotent algebras ($f(x, \dots, x) \approx x$)
especially all semilattices, lattices, Boolean algebras
- 5 (partially) bounded lattices, or semilattices resp. ($|A| > 1$)

Irreducible classes of algebras

i.e. all member are irreducible algebras

- 1 $|A| = 1$: one-element algebras without nullary constants
- 2 $|A| = 2$: all two-element algebras
- 3 essentially unary algebras $\text{Clo}(\mathbf{A}) = \langle \text{Clo}^{(1)}(\mathbf{A}) \rangle_{O_A}$
- 4 all idempotent algebras ($f(x, \dots, x) \approx x$)
especially all semilattices, lattices, Boolean algebras
- 5 (partially) bounded lattices, or semilattices resp. ($|A| > 1$)
- 6 orthocomplemented lattices, Boolean algebras ($|A| > 1$)

Irreducible classes of algebras

i.e. all member are irreducible algebras

- 1 $|A| = 1$: one-element algebras without nullary constants
- 2 $|A| = 2$: all two-element algebras
- 3 essentially unary algebras $\text{Clo}(\mathbf{A}) = \langle \text{Clo}^{(1)}(\mathbf{A}) \rangle_{O_A}$
- 4 all idempotent algebras ($f(x, \dots, x) \approx x$)
especially all semilattices, lattices, Boolean algebras
- 5 (partially) bounded lattices, or semilattices resp. ($|A| > 1$)
- 6 orthocomplemented lattices, Boolean algebras ($|A| > 1$)
- 7 vector spaces $\neq \{0\}$ over finite fields

Groups

Result

Let $\mathbf{G} = \langle G; \cdot, {}^{-1}, e \rangle$ be a finite group and $\exp \mathbf{G} = \text{lcm} \{ \text{ord } x \mid x \in G \} = \prod_{i=1}^k \underbrace{p_i^{k_i}}_{:=q_i}$.

Groups

Result

Let $\mathbf{G} = \langle G; \cdot, {}^{-1}, e \rangle$ be a finite group and $\exp \mathbf{G} = \text{lcm} \{ \text{ord } x \mid x \in G \} = \prod_{i=1}^k \underbrace{p_i^{k_i}}_{:=q_i}$.

Then

$$\begin{array}{ccc}
 U : \mathcal{P}(\{q_1, \dots, q_k\}) & \longrightarrow & \text{Neigh } \mathbf{G} \\
 S & \longmapsto & U(S) := \left\{ x \in G \mid x^{\prod S} = e \right\}
 \end{array}$$

is a **lattice isomorphism**.

Groups

Result

Let $\mathbf{G} = \langle G; \cdot, {}^{-1}, e \rangle$ be a finite group and $\exp \mathbf{G} = \text{lcm} \{ \text{ord } x \mid x \in G \} = \prod_{i=1}^k \underbrace{p_i^{k_i}}_{:=q_i}$.

Then

$$U: \mathcal{P}(\{q_1, \dots, q_k\}) \longrightarrow \text{Neigh } \mathbf{G}$$

$$S \longmapsto U(S) := \left\{ x \in G \mid x^{\prod S} = e \right\}$$

is a **lattice isomorphism**. The **irreducible neighbourhoods** of \mathbf{G} are exactly the **atoms** of this lattice, that is the neighbourhoods of the form

$$U_i = \{ x \in G \mid x^{q_i} = e \} \quad (1 \leq i \leq k)$$

Groups

Result

Let $\mathbf{G} = \langle G; \cdot, {}^{-1}, e \rangle$ be a finite group and $\exp \mathbf{G} = \text{lcm} \{ \text{ord } x \mid x \in G \} = \prod_{i=1}^k \underbrace{p_i^{k_i}}_{:=q_i}$.

Then

$$U: \mathcal{P}(\{q_1, \dots, q_k\}) \longrightarrow \text{Neigh } \mathbf{G}$$

$$S \longmapsto U(S) := \left\{ x \in G \mid x^{\prod S} = e \right\}$$

is a **lattice isomorphism**. The **irreducible neighbourhoods** of \mathbf{G} are exactly the **atoms** of this lattice, that is the neighbourhoods of the form

$$U_i = \{ x \in G \mid x^{q_i} = e \} \quad (1 \leq i \leq k)$$

$$= \bigcup \text{Syl}_{p_i} \mathbf{G}.$$

Semigroups

Result (Waldhauser, 2009)

A finite semigroup $\mathbf{S} = \langle S; \cdot \rangle$ is irreducible iff

- it is either **not completely regular**

Semigroups

Result (Waldhauser, 2009)

A finite semigroup $\mathbf{S} = \langle S; \cdot \rangle$ is irreducible iff

- it is either **not completely regular**
i.e. there is one $x \in S$ that is not contained in a subgroup

Semigroups

Result (Waldhauser, 2009)

A finite semigroup $\mathbf{S} = \langle S; \cdot \rangle$ is irreducible iff

- it is either **not completely regular**
- or it is a **union of subgroups** whose sizes are all **powers** of a common **prime**.

Semigroups

Result (Waldhauser, 2009)

A finite semigroup $\mathbf{S} = \langle S; \cdot \rangle$ is irreducible iff

- it is either **not completely regular**
- or it is a **union of subgroups** whose sizes are all **powers** of a common **prime**.

$\mathbf{S} = \langle S; \cdot \rangle$ has the following irreducible neighbourhoods:

Semigroups

Result (Waldhauser, 2009)

A finite semigroup $\mathbf{S} = \langle S; \cdot \rangle$ is irreducible iff

- it is either **not completely regular**
- or it is a **union of subgroups** whose sizes are all **powers** of a common **prime**.

$\mathbf{S} = \langle S; \cdot \rangle$ has the following irreducible neighbourhoods:

- S if it is **not completely regular**

Semigroups

Result (Waldhauser, 2009)

A finite semigroup $\mathbf{S} = \langle S; \cdot \rangle$ is irreducible iff

- it is either **not completely regular**
- or it is a **union of subgroups** whose sizes are all **powers** of a common **prime**.

$\mathbf{S} = \langle S; \cdot \rangle$ has the following irreducible neighbourhoods:

- S if it is **not completely regular**
- set of **idempotents** $\{x \in S \mid x^2 = x\}$

Semigroups

Result (Waldhauser, 2009)

A finite semigroup $\mathbf{S} = \langle S; \cdot \rangle$ is irreducible iff

- it is either **not completely regular**
- or it is a **union of subgroups** whose sizes are all **powers** of a common **prime**.

$\mathbf{S} = \langle S; \cdot \rangle$ has the following irreducible neighbourhoods:

- S if it is **not completely regular**
- set of **idempotents** $\{x \in S \mid x^2 = x\}$
- for every prime divisor p of $\exp \text{Gr } \mathbf{S} := \text{lcm} \{ \text{ord } x \mid x \in \text{Gr } \mathbf{S} \}$
the set

$$\{x \in \text{Gr } \mathbf{S} \mid \text{ord } x \text{ is a power of } p\}.$$

Subalgebra primal algebras

A subalgebra primal $:\iff \exists Q \subseteq R_A^{(1)} : \text{Clo}(\mathbf{A}) = \text{Pol}_A Q$

Subalgebra primal algebras

$$\begin{aligned} \mathbf{A} \text{ subalgebra primal} &: \iff \exists Q \subseteq R_A^{(1)} : \text{Clo}(\mathbf{A}) = \text{Pol}_A Q \\ &\iff \text{Clo}(\mathbf{A}) = \text{Pol}_A \text{Inv}^{(1)} \mathbf{A} \end{aligned}$$

Subalgebra primal algebras

$$\mathbf{A} \text{ subalgebra primal} : \iff \exists Q \subseteq R_A^{(1)} : \text{Clo}(\mathbf{A}) = \text{Pol}_A Q$$
$$\iff \text{Clo}(\mathbf{A}) = \text{Pol}_A \text{Inv}^{(1)} \mathbf{A}$$

\implies unary relational structures $\underline{\mathbf{A}} = \langle A; Q \rangle$, where $Q \subseteq R_A^{(1)}$.

Subalgebra primal algebras

$$\begin{aligned} \mathbf{A} \text{ subalgebra primal} &: \iff \exists Q \subseteq R_A^{(1)} : \text{Clo}(\mathbf{A}) = \text{Pol}_A Q \\ &\iff \text{Clo}(\mathbf{A}) = \text{Pol}_A \text{Inv}^{(1)} \mathbf{A} \end{aligned}$$

\implies unary relational structures $\underline{\mathbf{A}} = \langle A; Q \rangle$, where $Q \subseteq R_A^{(1)}$.

$$\underline{\mathbf{A}} \text{ irreducible} \iff \mathbf{A} = \langle A; \text{Pol}_A Q \rangle \text{ is irreducible}$$

Subalgebra primal algebras

$$\mathbf{A} \text{ subalgebra primal} : \iff \exists Q \subseteq R_A^{(1)} : \text{Clo}(\mathbf{A}) = \text{Pol}_A Q$$

$$\iff \text{Clo}(\mathbf{A}) = \text{Pol}_A \text{Inv}^{(1)} \mathbf{A}$$

\implies unary relational structures $\underline{\mathbf{A}} = \langle A; Q \rangle$, where $Q \subseteq R_A^{(1)}$.

$$\underline{\mathbf{A}} \text{ irreducible} \iff \mathbf{A} = \langle A; \text{Pol}_A Q \rangle \text{ is irreducible}$$

$$\iff \underline{\mathbf{A}} \rightsquigarrow \text{type } 1 / 2$$


Irreducible subalgebra primal algebras

Let $\underline{\mathbf{A}} = \langle A; Q \rangle$ be a unary relational structure where $Q \leq \langle \mathcal{P}(A), A, \cap \rangle$ as follows. $\implies \underline{\mathbf{A}}$ **irreducible**.

Irreducible subalgebra primal algebras

Let $\mathbf{A} = \langle A; Q \rangle$ be a unary relational structure where $Q \leq \langle \mathcal{P}(A), A, \cap \rangle$ as follows. $\implies \mathbf{A}$ **irreducible**.


$$\textcircled{1} \{B, B \cup \{a\}\} \cup \{\{x\} \mid x \in A \setminus \{a\}\} \subseteq Q$$




Irreducible subalgebra primal algebras

Let $\underline{\mathbf{A}} = \langle A; Q \rangle$ be a unary relational structure where $Q \leq \langle \mathcal{P}(A), A, \cap \rangle$ as follows. $\implies \underline{\mathbf{A}}$ **irreducible**.

$$\textcircled{1} \quad \{B, B \cup \{a\}\} \cup \{\{x\} \mid x \in A \setminus \{a\}\} \subseteq Q$$



$$\textcircled{2} \quad \{a, b\} \cup \{\{x\} \mid x \in A \setminus \{a, b\}\} \subseteq Q$$



Problems / remarks

- 1 Characterisation of irreducible neighbourhoods of certain finite algebras!!!

Problems / remarks

- 1 Characterisation of irreducible neighbourhoods of certain finite algebras!!!
- 2 Well-behaved algebras and difficult ones

Problems / remarks

- 1 Characterisation of irreducible neighbourhoods of certain finite algebras!!!
- 2 Well-behaved algebras and difficult ones
- 3 Considering polynomial clones (algebras \mathbf{A}_A)

Problems / remarks

- 1 Characterisation of irreducible neighbourhoods of certain finite algebras!!!
- 2 Well-behaved algebras and difficult ones
- 3 Considering polynomial clones (algebras \mathbf{A}_A)
- 4 Structure of the clone lattice

Problems / remarks

- 1 Characterisation of irreducible neighbourhoods of certain finite algebras!!!
- 2 Well-behaved algebras and difficult ones
- 3 Considering polynomial clones (algebras \mathbf{A}_A)
- 4 Structure of the clone lattice
- 5 Other structures, e.g. congruence primal algebras, quasi primal algebras...

Problems / remarks

- 1 Characterisation of irreducible neighbourhoods of certain finite algebras!!!
- 2 Well-behaved algebras and difficult ones
- 3 Considering polynomial clones (algebras \mathbf{A}_A)
- 4 Structure of the clone lattice
- 5 Other structures, e.g. congruence primal algebras, quasi primal algebras...
- 6 Your ideas ...

References:



Keith A. Kearnes.

Tame Congruence Theory is a localization theory.

Lecture Notes from “A Course in Tame Congruence Theory” Workshop, Budapest, 2001, available at

<http://www.math.u-szeged.hu/confer/algebra/2001/lec+ex1.ps>, 2001.



Mike Behrisch.

Relational Tame Congruence Theory and subalgebra primal algebras.

Master's thesis, TU Dresden [Dresden University of Technology], available at
http://www.math.tu-dresden.de/~mbehri/documents/subalgebraPrimal_links.pdf, September 2009.



Tamás Waldhauser.

Irreducible semigroups.

personal communication, 2009.